

Journal of Degraded and Mining Lands Management

<http://www.jdmlm.ub.ac.id>

DOAJ
DIRECTORY OF
OPEN ACCESS
JOURNALS

Crossref

ProQuest

CABI

ASEAN
CITATION
INDEX

S1
sinta

Scopus

Q3
1st 2024
0.28

p-ISSN: 2339-076X

e-ISSN: 2502-2458

Vol 13 No 1 (January 2026)



Research Centre for the Management of Degraded and Mining Lands,
and Soil Department, Faculty of Agriculture, Brawijaya University
<http://ircmedmind.ub.ac.id>, <http://tanah.ub.ac.id>





[Home](#) / [Editorial Team](#)

Editorial Team

Editor-in-Chief

Eko Handayanto, Research Centre for the Management of Degraded and Mining Lands, Brawijaya University, Indonesia

Associate Editor-in-Chief

Christopher W Anderson, School of Agriculture and Environment, Massey University, New Zealand

Reni Ustiatik, Research Centre for the Management of Degraded and Mining Lands, Brawijaya University, Indonesia

Sri Rahayu Utami, Soil Department, Faculty of Agriculture, Brawijaya University, Indonesia

Wani Hadi Utomo, Soil Department, Faculty of Agriculture, Brawijaya University, Indonesia

Editorial Board

Anizan Isahak, Faculty of Science and Technology, School of Environmental and Natural Resource Sciences, Universiti Kebangsaan Malaysia, Malaysia

Baiq Dewi Krisnayanti, United Nations Development Programme, Jakarta, Indonesia

B M Kumar, College of Forestry, Kerala Agricultural University, India

Constantinos Ehalotis, Department of Natural Resources and Agricultural Engineering, Agricultural University of Athens, Greece

Gusti Irya Ichriani, Department of Soil Science, Faculty of Agriculture, Lambung Mangkurat University, South Kalimantan, Indonesia

Hamdan Jol, Department of Land Management, Faculty of Agriculture, Universiti Putra Malaysia, Malaysia

Jianxu Wang, State Key Laboratory of Environmental Geochemistry, Chinese Academy of Sciences, Guiyang, China

Suhartini S, Department of Agricultural Economics, Faculty of Agriculture, Brawijaya University, Indonesia

Syahrul Kurniawan, Soil Department, Faculty of Agriculture, Brawijaya University, Indonesia

Widowati, Department of Agrotechnology, Faculty of Agriculture, Tribhuwana Tunggaladewi University, Indonesia

Yasuyuki Okimori, Japan Biochar Research Center, Ritsumeikan University, Japan

Technical Editor


Achmad Riyanto, Soil Department, Faculty of Agriculture, Brawijaya University, Indonesia

Christanti Agustina, Soil Department, Faculty of Agriculture, Brawijaya University, Indonesia

Nina Dwi Lestari, Soil Department, Faculty of Agriculture, Brawijaya University, Indonesia

Rizki Trisnadi, Brawijaya University, Indonesia

Vol. 13 No. 1 (2026)



DOI: <https://doi.org/10.15243/jdmlm.2026.131>

Published: 01-01-2026

Research Article

Geochemical stability of reclaimed coal mine waste dump: Implications for pit lake water quality and post-mining landscape utilization

Didik Triwibowo, Muthia Elma, Eko Suhartono, Rony Riduan, Ihsan Noor

9059-9070



DOI : [10.15243/jdmlm.2026.131.9059](https://doi.org/10.15243/jdmlm.2026.131.9059)

Spatial correlation between coastal distance and pyrite pH in acid sulfate soils of Paser, East Kalimantan

Titin Eka Setianingsih, Bagus Adi Nugroho, Raden Mas Nurhartanto, Zulkarnain Zulkarnain

9071-9080



DOI : [10.15243/jdmlm.2026.131.9071](https://doi.org/10.15243/jdmlm.2026.131.9071)

Abstract View : 172

pdf downloads: 190

Geospatial analysis of land degradation and settlement in Barru Regency: Implications for sustainable land management

Hämman Badruttamanan Amiruddin, Samsu Arif, Sakka Sakka

9081-9094



DOI : [10.15243/jdmlm.2026.131.9081](https://doi.org/10.15243/jdmlm.2026.131.9081)

Abstract View : 141

pdf downloads: 162

Groundwater quality assessment and suitability for drinking and irrigation purposes in the Jassan District, eastern Iraq

Sattar Obaid Maiws Al-Mayyahi, Hussein Ali Awadh Al-Zamili

9095-9109



DOI : [10.15243/jdmlm.2026.131.9095](https://doi.org/10.15243/jdmlm.2026.131.9095)

Abstract View : 165

pdf downloads: 141

Modification of contour planting system and its impact on improving infiltration characteristics of suboptimal land and soybean yields

 Sunarti Sunarti , Endriani Endriani , Najla Anwar Fuadi

 9111-9120



 DOI : 10.15243/jdmim.2026.131.9111

 Abstract View : 80

 pdf downloads: 80

Evaluation of soil and water properties in relation to indicative rice productivity in the coastal fields of Pangandaran, West Java, Indonesia

 Nasrudin Nasrudin , Budiastuti Kurniasih , Eka Tarwaca Susila Putra , Eko Hanudin

 9121-9130



 DOI : 10.15243/jdmim.2026.131.9121

 Abstract View : 142

 pdf downloads: 100

Assessing long-term land use/land cover changes in Dong Thap Province, Upper Vietnamese Mekong Delta: A 33-year retrospective using satellite data

 Ho Nguyen , Ta Hoang Trung , Tran The Dinh

 9131-9140



 DOI : 10.15243/jdmim.2026.131.9131

 Abstract View : 146

 pdf downloads: 78

Granulometric assessment of sediment accumulation for desilting the Toukomtoré agricultural water reservoir in Burkina Faso

 Adama Ilboudo , P. Louis Yameogo , Joseph Nomwindé Kabore , Daouda Guébré , Harouna Ouédraogo , Edmond Hien

 9141-9151



 DOI : 10.15243/jdmim.2026.131.9141

 Abstract View : 76

 pdf downloads: 53

The potential of fungi isolated from palm oil mill effluent (POME) for mycoremediation of water contaminated with pome waste

 Zakaria Zakaria , Miftahul Ilmi , Rina Sri Kasiamdari

 9153-9162



 DOI : 10.15243/jdmim.2026.131.9153

 Abstract View : 201

 pdf downloads: 148

Hydrological modeling of small coastal peat island in degraded peatlands of Bengkalis Island, Riau Province

 Sigit Sutikno , Muhamad Yusa , Rinaldi Rinaldi , Ahmad Muhammad , Hendra Saputra , Khusnul Setia Wardani , Koichi Yamamoto

 9163-9175



 DOI : 10.15243/jdmim.2026.131.9163

 Abstract View : 170

 pdf downloads: 87

Application of organic matter-enriched biochar for improving degraded acid sulfate soil properties in rice (*Oryza sativa* L.) cultivation

 Agusalim Masulili , Sri Rahayu , Ida Ayu Suci , Edi Kurniadi

 9177-9190



 DOI : 10.15243/jdmim.2026.131.9177

 Abstract View : 128

 pdf downloads: 96

Nutrient and productivity indices as tools for soil fertility assessment in Chebli, Mitidja Plain, Algeria

 Abdelkader Laribi , Amel Krireche , Zahida Dehnoun

 9191-9198



 DOI : 10.15243/jdmim.2026.131.9191

 Abstract View : 74

 pdf downloads: 66

Sustainable management strategies for degraded subak rice fields in South Denpasar, Bali

 Lina Anandari Dwi Ambarini , Evi Frimawaty , Herdis Herdiansyah

 9199-9209



 DOI : 10.15243/jdmim.2026.131.9199

 Abstract View : 117

 pdf downloads: 101

The impact of nickel mining on vegetation cover in a small island: a case study in Manuran Island, Raja Ampat, Indonesia

 Mohammad Afdhal Adidharma , Nurul Mu'min Z , Adam Santrio , Andi Azwar Anas Idris , Anggun Rahmi Diah Fatmi , Awaluddin Awaluddin , Nanang Pindasari , Elly Ermawati , Andhy Rahmat Padyawan , Sirniawan Sirniawan

 9211-9223



 DOI : 10.15243/jdmim.2026.131.9211

 Abstract View : 232

 pdf downloads: 120

Predicting landscape transitions with machine learning: a case study of the Nagavalli Basin (2000-2030)

 Ruba Maarouf , Vazeer Mahamood , P. Jagadeeswara Rao

 9225-9237



 DOI : 10.15243/jdmim.2026.131.9225

 Abstract View : 39

 pdf downloads: 47

After mining catastrophe: the language of economic survival of the Sipalay mine people at Sipalay, Negros Occidental, Philippines

 Rubie Lyn C. Barrientos , John Gerald A. Pilar

 9239-9257



 DOI : 10.15243/jdmim.2026.131.9239

 Abstract View : 151

 pdf downloads: 69

Rehabilitation of disturbed land in bauxite mining: Challenges and prospects in Sanggau, West Kalimantan, Indonesia

Wanda Wanda , Aji Ali Akbar , Erisa Ayu Waspadi Putri , Aji Suhartoyo

9259-9269



DOI : 10.15243/jdmim.2026.131.9259

Abstract View : 234

pdf downloads: 144

Land cover and socio-economic dynamics of coffee to oil palm land conversion in Way Kanan, Indonesia

Tiara Aprilia Putri Hernanda , Akhmad Fauzi , Baba Barus , Bustanul Arifin

9271-9291



DOI : 10.15243/jdmim.2026.131.9271

Abstract View : 89

pdf downloads: 109

Improving substrate properties and foxtail millet growth in gold mine tailings using Azotobacter consortium

Reginawanti Hindersah , Aliya Zahrah Adawiah , Pujawati Suryatmana , Marenda Ishak Sonjaya Sule , Fiky Yulianto Wicaksono , Triyani Dewi

9293-9302



DOI : 10.15243/jdmim.2026.131.9293

Abstract View : 55

pdf downloads: 43

Modeling of groundwater conservation zone using the modified GOD method on part of the southern slope of Merapi Volcano, Yogyakarta, Indonesia

Sadewa Purba Sejati , Rivi Neritarani

9303-9317



DOI : 10.15243/jdmim.2026.131.9303

Abstract View : 61

pdf downloads: 67

GIS-based landslide susceptibility mapping in IV Koto, Matur, and Tanjung Raya Subdistricts of Agam Regency, West Sumatra

Riko Maiyudi , Adree Octova , Yoszi Mingsi Anaperta , Mulya Gusman , Dedi Yulhendra , Exwil Naldo , Adhitya Baron

9319-9328



DOI : 10.15243/jdmim.2026.131.9319

Abstract View : 131

pdf downloads: 76

Predictive model for California Bearing Ratio (CBR) in expansive coastal subgrades: a rapid geotechnical assessment for degraded and marginal lowland areas

Soewignjo Agus Nugroho , Syawal Satibi , Agus Ika Putra , Zulkifli Zulkifli , Sigit Sutikno , Muhamad Yusa , Rinaldi Rinaldi , Koichi Yamamoto

9329-9342



DOI : 10.15243/jdmim.2026.131.9329

Abstract View : 84

pdf downloads: 50

From resources to reserves: a real option approach to quarry mining in the Aimas River, West Papua

Arif Setiawan , Ricardo O. M. Hutapea

9343-9354



DOI : 10.15243/jdmlm.2026.131.9343

Abstract View : 74

pdf downloads: 43

Landslide susceptibility mapping based on K-Means and Self-Organizing Map clustering with Geographic Information System in Tasikmalaya, West Java

Mimin Iryanti , Nanang Dwi Ardi , Riska Siti Nurjanah

9355-9369



DOI : 10.15243/jdmlm.2026.131.9355

Abstract View : 105

pdf downloads: 78

Characterization of suspended sediment dynamics in Tanjung Pasir, Banten, Indonesia: Implications for the National Capital Integrated Coastal Development (NICCD) program's coastal reservoir

Reno Arief Rachman , Hamzah Haru Radityo Suharyanto , Aloysius Bagyo Widagdo , Maria Noosa Airawati , Ibnu Fauzi , Nofika Cahyani Putri , Fajar Yulianto , Dhedy Husada Fadjar Perdana , Mardi Wibowo

9371-9383



DOI : 10.15243/jdmlm.2026.131.9371

Abstract View : 83

pdf downloads: 50

Land suitability for chili on the slopes of Raung Indonesia, towards sustainable agriculture

Basuki Basuki , Tri Candra Setiawati , Bambang Hermiyanto , Subhan Arif Budiman , Vega Kartika Sari

9385-9398



DOI : 10.15243/jdmlm.2026.131.9385

Abstract View : 105

pdf downloads: 98

Study of seawater intrusion in the coastal alluvial aquifer, West Semarang, Central Java, Indonesia

Rizq Fajrianto , Wahyu Wilopo , Pulung Arya Pranantya , Pipat Laowattanabandit

9399-9413



DOI : 10.15243/jdmlm.2026.131.9399

Abstract View : 84

pdf downloads: 88

Assessment of the soil chemical properties on the intensive coffee plantations in the Central Highlands, Vietnam

Nguyen Vu Ky , Ta Hoang Trung , Pham Lan Anh , Hoang Hai Long , Dinh Thi Nha Truc , Nguyen Huy Anh

9415-9425



DOI : 10.15243/jdmlm.2026.131.9415

Abstract View : 81

pdf downloads: 92

Examining the several vegetation indices values to estimate carbon stock at the mining reclamation area of PT. Adaro Indonesia, South Kalimantan

 Dony Rachmanadi , Luthfan Meilana Nugraha , Dodik Choiron , Mekka Riadhah , Rusdi Husin , Didik Triwibowo , Ajiddinnor Ajiddinnor

 9427-9442



 DOI : 10.15243/jdmim.2026.131.9427

 Abstract View : 85

 pdf downloads: 80

Spatial assessment of erosion and landslide hazards as indicators of land degradation in Solok Regency, West Sumatra

 Dasrizal Dasrizal , Erna Juita , Ratna Wilis , Febriandi Febriandi , Trina Febriani

 9443-9453



 DOI : 10.15243/jdmim.2026.131.9443

 Abstract View : 164

 pdf downloads: 49

Synergistic effects of potassium humate and clay-rich soil on improving the chemical and physical properties of tin tailing

 Lambang Praja Panghingguan , Arief Hartono , Anwar Syaiful

 9455-9462



 DOI : 10.15243/jdmim.2026.131.9455

 Abstract View : 107

 pdf downloads: 49

Phosphorus adsorption and release dynamics in calcareous soil as affected by humic and fulvic acids

 Aswad H. Aswad , Mohammed M. Hamid , Mohammad Tarkhan Abo Almeekh , Mohammed R. Ajeel

 9463-9472



 DOI : 10.15243/jdmim.2026.131.9463

 Abstract View : 75

 pdf downloads: 53

Spatial modeling of soil erosion in the Teesta River Basin in Bangladesh using RUSLE and remote sensing data in Google Earth Engine

 Erni Saurmalinda Butar Butar , Jedtavong Thepvongsa

 9473-9486



 DOI : 10.15243/jdmim.2026.131.9473

 Abstract View : 130

 pdf downloads: 102

The forest farmers' mindsets contribution to ecosystem degradation in Meru Betiri National Park, Indonesia: towards sustainable conservation through a translation practices approach

 Fahrudi Ahwan Ikhsan , Sumarmi Sumarmi , Sugeng Utaya , Agus Sugiarto , Eni Heldayani

 9487-9504



 DOI : 10.15243/jdmim.2026.131.9487

 Abstract View : 79

 pdf downloads: 86

Sustainable small-scale mining through geological and resource estimation: a holistic, society-centred approach

 Ntiyiso Ally Mahtlaule , Fhatuwani Sengani , Maropene Tebello Dina Rapholo

 9505-9530



 DOI : 10.15243/jdmlm.2026.131.9505

 Abstract View : 118

 pdf downloads: 76

Assessment of water erosion risk in the Fergoug watershed in Algeria using a GIS-based RUSLE approach

 Ahmed Tewfik Oualid , Abderrahmane Hamimed , Benyekhlef Benamina , Kaddour Miloudi , Mohammed Benslimane

 9531-9544



 DOI : 10.15243/jdmlm.2026.131.9531

 Abstract View : 161

 pdf downloads: 117

Native arbuscular mycorrhizal fungi and Albizia saponaria for potential revegetation of tropical asphalt and nickel post-mining areas

 Faisal Danu Tuheeru , Husna Husna , Wiwin Rahmawati Nurdin , Surya Edy Wibowo , Edy Jamal Tuheeru , Albasri Albasri , Asrianti Arif

 9545-9554



 DOI : 10.15243/jdmlm.2026.131.9545

 Abstract View : 58

 pdf downloads: 64

Block compost for acid mine drainage remediation and as a growing medium for Melaleuca cajuputi seedlings

 Nisa Daniswati , Irdika Mansur , Elis Nina Herliyana

 9555-9565



 DOI : 10.15243/jdmlm.2026.131.9555

 Abstract View : 74

 pdf downloads: 52

Geotechnical zonation and bearing capacity assessment for risk-informed foundation design in the heterogeneous soils of Girsang Sipangan Bolon, Indonesia

 Refky Adi Nata , Gaofeng Ren , Jukepsa Andas , Yudi Arista Yulanda , Hisni Rahmi , Verra Syahmer

 9567-9585



 DOI : 10.15243/jdmlm.2026.131.9567

 Abstract View : 144

 pdf downloads: 78

The potential of coffee husk biochar for chromium removal from acid mine drainage using constructed wetlands

 Opi Mida Titania , Fahrudin Fahrudin , Andi Ilham Latunra

 9587-9596



 DOI : 10.15243/jdmlm.2026.131.9587

 Abstract View : 98

 pdf downloads: 94

Effect of the combined application of rhizobacteria and compost on the ability of *Pteris vittata* L. as a phytoremediator for mercury in gold mine tailing

Tien Aminatun , Budiwati Budiwati , Suhartini Suhartini , Rahmanto Raharjo

9597-9606



DOI : 10.15243/jdmim.2026.131.9597

Abstract View : 201

pdf downloads: 118

Soil deterioration assessment as an indicator of land degradation in dry land ecosystems based on a multi-criteria approach, North Halmahera, Indonesia

Adnan Sofyan , Tri Mulya Hartati , Lily Ishak , Krishna Aji , Gunawan Hartono , Nurul Ainun Tangge , Fitriani H. Achmad

9607-9621



DOI : 10.15243/jdmim.2026.131.9607

Abstract View : 125

pdf downloads: 73

Assessment of landslides in the Banjarsari area, Ciamis Regency, Indonesia: A morphographic and morphometric approach

Mochammad Malik Ibrahim , Jasmin Arya Alfalah , Salsabila Az Zahra

9623-9634



DOI : 10.15243/jdmim.2026.131.9623

Abstract View : 45

pdf downloads: 55

Agricultural land, coal mining, and urbanisation: A GIS-based analysis of land cover changes in Kosovo

Besarta Ibrahim , Valbon Bytyqi , Tropikë Agaj , Ibrahim Ramadani

9635-9644



DOI : 10.15243/jdmim.2026.131.9635

Abstract View : 179

pdf downloads: 90

Dehydrogenase enzyme activity and lead (Pb) availability due to the application of biochar and *Bacillus* sp. in heavy metal-contaminated soil

Bella Agit Solekhah , Rossyda Priyadarshini , Maroeto Maroeto

9645-9657



DOI : 10.15243/jdmim.2026.131.9645

Abstract View : 108

pdf downloads: 58

Bioremediation of the Lapindo mud contaminated soil using eco-enzyme: Case of a total petroleum hydrocarbon removal

Dwi Rakhmatun Sulistiyaningrum , Barlah Rumhayati , Arie Srihardyastutie

9659-9668



DOI : 10.15243/jdmim.2026.131.9659

Abstract View : 123

pdf downloads: 86

Path analysis of "new" humic acid derived from water hyacinth on phosphorus dynamics in P-deficient Inceptisols and chili response under intensive cultivation

Emma Trinurani Sofyan , Mega Kartika Hermawan , Rija Sudirja , Anne Nurbaiti , Betty Natalie Fitriatin , Oviyanti Mulyani , Mieke Rochimi Setiawati , Muhamad Rahman Djuansah

9669-9687



DOI : 10.15243/jdmlm.2026.131.9669

Abstract View : 107

pdf downloads: 75

Research Article

Native arbuscular mycorrhizal fungi and *Albizia saponaria* for potential revegetation of tropical post-asphalt mining and post-nickel mining areas

**Faisal Danu Tuheteru^{1*}, Husna¹, Wiwin Rahmawati Nurdin¹, Surya Edy Wibowo¹,
Edy Jamal Tuheteru², Albasri¹, Asrianti Arif¹**

¹ Department of Forestry, Faculty of Forestry and Environmental Science, Universitas Halu Oleo, Kendari 93121, Indonesia

² Department of Mining Engineering, Faculty of Earth and Energy Technology, Trisakti University, Jakarta, Indonesia

*corresponding authors: fdtuheteru1978@gmail.com

Abstract

Article history:

Received 8 November 2025

Revised 5 December 2025

Accepted 11 December 2025

Keywords:

Glomus intraradices

legume trees

restoration

tropical region

Native arbuscular mycorrhizal fungi (AMF) have been shown to effectively enhance the growth and nutrient uptake of *Vitex cofassus* grown on post-asphalt mining soils. However, their potential needs to be further evaluated across various mining substrates and different tree species. This study aimed to assess the effects of native AMF inoculation on the early growth performance of *Albizia saponaria* seedlings cultivated on post-asphalt mining and post-nickel mining media under greenhouse conditions. A completely randomized design was employed with seven treatments: control, *Racocetra crispi*, *Glomus intraradices*, *Glomus* sp., *Glomus* sp.-LW10, *Glomus* sp.-SW10, and Mycofer IPB or AMF mixture. Parameters observed after three months of growth included plant height, AMF root colonization percentage, total dry weight, mycorrhizal inoculation effect (MIE), and the uptake of phosphorus (P), nitrogen (N), calcium (Ca), and nickel (Ni). The results indicated that native AMF inoculation significantly improved early growth and total dry weight of *A. saponaria* seedlings. The highest MIE values were observed in the *G. intraradices* treatment, reaching 92.73% and 82.9%. Native AMF inoculation also enhanced the uptake of P, N, Ca, and Ni in plant tissues. These findings suggest that local AMF isolates can be effectively utilized to improve seedling quality and accelerate the success of revegetation programs on degraded post-mining lands in Indonesia.

To cite this article: Tuheteru, F.D., Husna, Nurdin, W.R., Wibowo, S.E., Tuheteru, E.J., Albasri, and Arif, A. 2026. Native arbuscular mycorrhizal fungi and *Albizia saponaria* for potential revegetation of tropical post-asphalt mining and post-nickel mining areas. Journal of Degraded and Mining Lands Management 13(1):9545-9554, doi:10.15243/jdmlm.2026.131.9545.

Introduction

Reclamation of post-mining land poses a significant environmental management challenge in tropical regions, particularly in Indonesia, where extensive areas of degraded land result from mineral and asphalt mining activities (Pratiwi et al., 2021; Albasri et al., 2023; Miswanto et al., 2025). Mining operations lead to the loss of topsoil layers, depletion of organic matter, and disruption of soil structure, all of which reduce the soil's ability to support plant growth (Nasution et al., 2024). In post-nickel mining areas,

soils are characterized by high concentrations of heavy metals (Ni, Cr, Co), a low Ca/Mg ratio, and deficiencies in essential nutrients such as phosphorus (P), potassium (K), and nitrogen (N) (Prematuri et al., 2020). In contrast, post-asphalt mining soils are dominated by clay and clay loam fractions, have slightly alkaline pH, and exhibit low nutrient availability (Albasri et al., 2023; Tuheteru et al., 2025). Therefore, soil biotechnology-based restoration approaches are required to improve soil fertility and facilitate early plant growth at both the nursery and field stages. One promising soil biotechnological

approach to address these challenges is the inoculation of arbuscular mycorrhizal fungi (AMF). Arbuscular mycorrhizal fungi (AMF) are mutualistic symbionts of most terrestrial plants that play a key role in enhancing nutrient uptake, particularly phosphorus (P), and improving plant tolerance to various abiotic stresses such as drought, salinity, and heavy metal toxicity (Bi et al., 2018; Begum et al., 2019).

In degraded lands resulting from mining activities, AMF function as natural bio-ameliorants by forming extensive hyphal networks that enhance soil aggregate stability, improve soil structure, and support root colonization (Begum et al., 2019; de Moura et al., 2022). The effectiveness of AMF varies among species; therefore, screening is necessary to identify the most effective isolates for land reclamation (Wulandari et al., 2024). Native AMF inoculation is considered more adaptive than commercial isolates, as native strains possess greater resilience and colonization ability under extreme environmental conditions (Brundrett and Tedersoo, 2018).

Studies conducted on post-mining lands in Sulawesi have shown that local arbuscular mycorrhizal fungi (AMF) can enhance the growth of several tree species, such as *Pericopsis mooniana*, *Pterocarpus indicus*, and *Kalappia celebica*, on nutrient-poor substrates (Husna et al., 2021a,b; 2025). In addition, the use of AMF originating from post-asphalt mining soils has been proven to improve growth, root biomass, and phosphorus uptake efficiency in *Vitex cofassus* seedlings grown on post-asphalt mining media (Tuheteru et al., 2025). However, no studies have yet examined the effects of local AMF on the early growth of native Sulawesi forest tree seedlings grown in soil media with contrasting characteristics, such as post-asphalt and post-nickel mining substrates.

One leguminous tree species from Sulawesi selected to evaluate its growth response and dependency on native AMF inoculation was *Albizia saponaria* (Lour.) Blume. *A. saponaria* is a multipurpose tree species in the Fabaceae family, with significant ecological and economic value in tropical regions of Southeast Asia. This species is characterized by rapid growth, the ability to fix atmospheric nitrogen through symbiosis with rhizobia, and tolerance to nutrient-poor soils (Fern, 2024; Hikmat et al., 2025). Furthermore, *A. saponaria* has been reported to exhibit a strong response and high dependency on local AMF inoculation when grown in Ultisol soil media (Tuheteru et al., 2021). These characteristics make *A. saponaria* a promising candidate for revegetation of post-asphalt mining and post-nickel mining areas, with support from AMF inoculation.

The differences in substrate characteristics are expected to influence the effectiveness of AMF colonization and the physiological performance of inoculated plants. Therefore, understanding the early growth response of *A. saponaria* to native AMF inoculation from both substrates will provide valuable

insights into the potential use of local fungi as natural biofertilizer agents for mining ecosystem restoration.

This study aimed to evaluate the early growth of *A. saponaria* seedlings inoculated with several native AMF species originating from post-asphalt mining soils. The results of this research are expected to provide a scientific basis for the application of mycorrhizal inoculation technology in forest seedling production, particularly for the reclamation and restoration of post-mining lands in Indonesia.

Materials and Methods

Post-mining soils

The post-asphalt mining and post-nickel mining soils were obtained from the disposal site of PT Wika Bitumen in Buton District and PT Pernick Sultra in North Konawe District, Southeast Sulawesi, Indonesia. The soils were stored in a greenhouse before use. The physical and chemical properties of the soils were analyzed at the Biotechnology and Environmental Laboratory, Faculty of Mathematics and Natural Sciences, Halu Oleo University, Kendari, and the results are presented in Table 1.

Preparation of AMF inoculum and inoculation

AMF inocula used were *Racocetra crispera*, *Glomus intraradices*, *Glomus* sp., *Glomus* sp.-LW10, and *Glomus* sp.-SW1, which were isolated from the rhizosphere of plants in post-asphalt mining soil (Tuheteru et al., 2022) and Mycofer IPB. The AMF inoculum was propagated in zeolite media using *Pueraria javanica*, which was grown for three months under greenhouse conditions at the Indonesian Mycorrhiza Association (AMI), Southeast Sulawesi Branch (6°38'07.35" S dan 106°49'31.72" E), Kendari, Southeast Sulawesi, Indonesia.

Polyethylene pots (15 × 20 cm) were filled with 1 kg of sterile planting media (a mixture of post-asphalt mining soil or post-nickel mining soil, river sand, and husk charcoals having a proportion of 3:1:1). AMF inoculum was produced by placing 20 g of inoculum of each treatment at 1-3 cm underneath the seedlings. Two new leaves of respective *A. saponaria* seedlings were transplanted into pots, followed by seedling maintenance, watering, and observation for three months (10 July-11 October 2024 and 09 July-10 October 2025). The seedlings were watered daily to field capacity with tap water. The daytime temperature in the nursery during the trial was 25 to 35 °C, with 80-89% relative humidity and a 12-hour photoperiod.

Seed germination and AMF treatment

A. saponaria seeds were collected from their trees at the Arboretum of the Botanical Garden of Halu Oleo University, Kendari, Southeast Sulawesi. The seeds were soaked in hot water at 50-60 °C for 24 hours to cool gradually, then germinated in a plastic sprout

maker (20 × 20 × 5 cm) which has been perforated and contains sterile sand media at the Plastic House of the

Indonesian Mycorrhizal Association (IMA), Southeast Sulawesi Branch.

Table 1. The physical and chemical properties of post-asphalt mining and post-nickel mining soils.

No	Parameter	Post-asphalt mining soil		Post-nickel mining soil	
		Value	Criteria ^a	Value	Criteria ^b
1	pH H ₂ O	7.2	Neutral	7.84	
2	Organic C (Walkley and Black) (%)	7.56	Very high	4.30	High
3	Total N (Kjeldahl) (%)	0.13	Low	0.27	Moderate
4	C/N ratio	58	Very high	16	High
5	P ₂ O ₅ (HCl 25%) (mg 100 g ⁻¹)	55	High	26.41	Moderate
6	K ₂ O (HCl 25%) (mg 100 g ⁻¹)	7	Very low	22.60	Moderate
7	P ₂ O ₅ (Olsen) (ppm)	8	Low	-	-
8	K ₂ O (Morgan) (ppm)	49	-	-	-
9	Ca (NH ₄ -Acetate 1N pH 7) (cmol kg ⁻¹)	18.58	High	13.70	High
10	Mg (NH ₄ -Acetate 1N pH 7) (cmol kg ⁻¹)	1.47	Moderate	4.90	High
11	K (NH ₄ -Acetate 1N pH 7) (cmol kg ⁻¹)	0.10	Low	-	-
12	Na (NH ₄ -Acetate 1N pH 7) (cmol kg ⁻¹)	0.23	Low	-	-
13	CEC (NH ₄ -Acetate 1N pH 7) (cmol kg ⁻¹)	23.89	Moderate	18	Moderate
14	Base saturation (NH ₄ -Acetate 1N pH 7) (%)	85	Very high	-	-
15	Al ³⁺ (KCl 1N) (cmol kg ⁻¹)	0.00	-	-	-
16	H ⁺ (KCl 1N) (cmol kg ⁻¹)	0.07	-	-	-
17	Texture (pipet)				
18	Sand (%)	10		-	
19	Silt (%)	40	Clay	95.65	Silt
20	Clay (%)	50		4.35	
21	Fe (DTPA) (ppm)	7.3	Sufficient	128.72	Sufficient
22	Mn (DTPA) (ppm)	9.5	Sufficient	-	-
23	Cu (DTPA) (ppm)	1.2	Sufficient	-	-
24	Zn (DTPA) (ppm)	0.3	Deficient	-	-
25	Pb (Morgan Wolf) (ppm)	0.4	Normal		
26	Cd (Morgan Wolf) (ppm)	0.2	Normal		
27	CaCO ₃ (Titrimetric) (%)	2.1	-		
28	Ni (AAS flame) (mg kg ⁻¹)			0.0047	
29	Cr (AAS flame) (mg L ⁻¹)			0.0262	

Notes: ^aSoil Research Center (2009). ^bTuheteru et al. (2025).

The treatments tested in this experiment were (A) control, (B) *Racocetra crispera*, (C) *Glomus intraradices*, (D) *Glomus sp.*, (E) *Glomus sp.*-LW10, (F) *Glomus sp.*-SW10, and (G) Mycofer IPB (post-asphalt mining media) or AMF Mixed (post-nickel mining media). The seven treatments were arranged in a completely randomized design with three replications.

Growth, dry weight, and nutrient determination

Seedlings were harvested after three months of growth. The samples were then oven-dried at 70 °C for 48 hours to determine their shoot and root dry weight. Roots were stained using the modified protocol from Vierheilig et al. (2005). A total of 30 pieces of fresh roots (± 1 cm long) were randomly taken from the plant roots. The origins were cleaned in 10% KOH for 2 days, then immersed in H₂O₂ solution for 10-20 minutes, and rinsed thoroughly. The roots were then soaked in a 0.2% HCl solution for 20 minutes and placed in a 0.05% Trypan Blue solution. Ten root

samples were placed on a glass slide, a cover glass was applied, and the slide was observed under a microscope. Available P was measured in sodium bicarbonate extracts at pH 8.5 and quantified according to the Olsen method at 660 nm using a flow-injection automated ion analyzer. Available Ca was extracted with 1 N NH₄-Acetate, pH 7, and determined by atomic absorption. The Zn content was analyzed by digesting the sample with an acid mixture (HNO₃-H₂O₂) and measuring it using an Atomic Absorption Spectrophotometer (AAS). The Ni content was determined after digesting the sample with a mixture of HNO₃ and HCl at high temperature, followed by AAS analysis. Parameters and calculation methods are presented in Table 2.

Statistical analysis

The data were analyzed using analysis of variance (F test), followed by Duncan's Multiple Range Test (DMRT) at the 95% confidence level when the F test indicated a significant effect.

Table 2. Parameters and calculation methods.

Parameter	Formula	Reference
Plant growth	Plant height and stem diameter were measured 1 cm above the soil medium at three months after transplantation. At the end of the study, the number of leaves was counted.	Tuheteru et al. (2025)
Dry weight	Drying all plant parts in an oven at 70 °C until a constant weight was reached, then weighing them to obtain the root, shoot, and total dry weight.	Husna et al. (2025)
Nodule	Number of nodules per root.	Husna et al. (2025)
The Mycorrhizal Growth Responses (MGR)	[total dry weight of mycorrhizal plant-total dry weight of non-mycorrhizal plant/total dry weight of non-mycorrhizal plant] × 100%.	Cavagnaro et al. (2003)
AMF colonization	[Σ number of fields of view colonized/Σ total observed field of view] × 100.	Vierheilig et al. (2005)
Mycorrhizal dependency (%)	$(1-[b/a]) \times 100$; where b is the mean plant dry biomass of non-AMF treatments, and a is the mean plant dry biomass in treatments with AMF presence. Therefore, an MD>0 indicates plant biomass enhancement by AMF, while MD<0 indicates the opposite.	van der Heijden et al. (2003)
The content and uptake of N, P, Ca, and Ni	Content: the amount of nutrients contained in plant tissue and uptake: Multiplying the nutrient concentrations by the dry weights of the plants.	Wang et al. (2005)
The increase/decrease in nutrient uptake of AMF-treated seedlings relative to the controls	[nutrient absorption of AMF plant - nutrient absorption of non-mycorrhizal plant/nutrient absorption of non-mycorrhizal plant] × 100%.	Wang et al. (2005)

Results and Discussion

Colonization of AMF and MD

A. saponaria seedlings on post-asphalt mining media inoculated with Mycofer IPB had the highest colonization rate (97%), followed by *Glomus* sp.-SW10 (90.1%), *Glomus* sp.-LW10 (88.8%), and *Glomus* sp. (81.0%) (Table 1, $p < 0.01$). On post-nickel mining media, the highest colonization rate was observed in the *Glomus intraradices* treatment (52.83%) (Table 1, $p < 0.01$). The MD value for *G. intraradices* on both media was higher at 92.73% and 82.9%, respectively.

Plant growth

On post-asphalt mining land media, inoculation with *G. intraradices* significantly increased plant height and differed significantly from other treatments (Table 3). Three-month-old *A. saponaria* seedlings inoculated with *G. intraradices* had higher stem diameters and differed significantly from other treatments except for *Glomus* sp.-SW10. All AMF treatments significantly increased the number of plant leaves compared to the control. *A. saponaria* seedlings inoculated with *Glomus* sp.-SW10 had the highest number of root nodules and did not differ significantly from other treatments except for the control and *Glomus* sp.-LW10 treatments. In post-nickel mining media inoculation with *G. intraradices* significantly increased the height, diameter, and leaf number of

three-month-old plants. It differed significantly from other treatments, except for AMF Mixed (Table 3). *G. crista* significantly increased the number of root nodules and did not differ significantly from *G. intraradices*. The performance of the *A. saponaria* seedlings is presented in Figure 1.

Plant dry weight

On post-asphalt mining media, inoculation with *G. intraradices* significantly increased the dry weight of three-month-old *A. saponaria* and differed significantly from other treatments (Table 4). The increase in mycorrhizal growth (MGR) ranged from 376.9 to 1,523.1%. The SRR value ranged from 1.65 to 5.95. In post-nickel mining media, inoculation with *G. intraradices* significantly increased the dry weight of *A. saponaria* plants. It differed significantly from other treatments, except for AMF mixed in the root dry weight. The increase in mycorrhizal plant growth (MGR) ranged from 35.7 to 1,485.7%. The SRR value ranged from 0.94 to 1.35. There were no differences between treatments in the SQI variable.

Nutrient content and uptake

Inoculation with *G. intraradices* significantly increased the N and P levels of three-month-old *A. saponaria* plants in post-asphalt mining media (Table 5). In terms of P content, the *G. intraradices* treatment did not differ significantly from *R. crista* and differed significantly from the other treatments.

There were no differences between treatments in terms of plant Zn levels (Table 5). Inoculation with *G. intraradices* significantly increased n and p uptake in three-month-old *A. saponaria* plants (Table 5).

There were no differences between treatments in plant Zn uptake (Table 5). There were no differences between treatments in plant N and P content (Table 6).

Table 3. AMF colonization and growth of three-month-old *A. saponaria* seedlings grown with or without mycorrhizal fungi under greenhouse conditions.

Treatment	Mycorrhizal Colonization (%)*	Height (cm)	Stem Diameter (mm)	Leaf Number	Nodule
Post-asphalt mining media					
Control	11.2±2.72 c	6.90±0.46 d	0.40±0.10 c	5.7±0.67 b	0.00±0.00 c
<i>Gigaspora crispera</i>	65.9±3.74 b	29.17±1.53 b	1.10±0.12 abc	15.3±1.33 a	1.00±0.58 bc
<i>Glomus intraradices</i>	65.0±5.75 b	41.23±3.39 a	1.70±0.15 a	19.3±2.03 a	2.00±0.58 ab
<i>Glomus</i> sp.	81.0±10.14 ab	22.37±1.73 c	0.90±0.44 bc	16.7±0.33 a	2.33±0.33 ab
<i>Glomus</i> sp.-LW10	88.8±4.42 a	20.87±0.61 c	1.00±0.10 abc	16.7±1.45 a	0.00±0.00 c
<i>Glomus</i> sp.-SW10	90.1±6.05 a	22.43±1.85 c	1.27±0.33 ab	15.3±1.86 a	3.00±0.58 a
Mycofer IPB	97.0±0.12 a	19.60±1.15 c	0.87±0.03 bc	15.0±1.53 a	2.33±0.88 ab
Pr>F	<.0001	<.0001	0.0372	0.0004	0.0047
Post-nickel mining media					
Control	0.80±0.81 c	3.63±0.22 d	0.53±2.47 c	8.00±0.58 c	0.67±0.33 c
<i>Gigaspora crispera</i>	32.27±3.45 b	8.07±0.47 bc	0.76±0.02 bc	8.00±0.58 c	4.00±0.58 a
<i>Glomus intraradices</i>	52.83±3.96 a	12.10±0.81 a	1.21±0.11 a	11.3±0.67 a	3.67±0.67 ab
<i>Glomus</i> sp.	30.47±3.33 b	7.03±1.11 c	0.90±0.11 b	8.67±0.88 bc	0.67±0.33 c
<i>Glomus</i> sp.-LW10	31.93±2.78 b	7.97±0.71 bc	0.79±0.15 bc	9.00±0.58 bc	0.33±0.33 c
<i>Glomus</i> sp.-SW10	14.87±1.24 bc	6.17±0.35 cd	0.89±0.08 b	8.67±0.33 bc	0.00±0.00 d
AMF Mixed	35.20±1.82 ab	10.63±1.79 ab	0.99±0.18a b	10.3±0.67 ab	2.67±0.33 b
Pr>F	0.0014	0.0003	0.0088	0.0167	0.0047

Notes: Average values followed by different letters in the same column are significantly different at DMRT ($p < 0.05$); *Mean±SE.



Figure 1. Performance of three-month-old *A. saponaria* seedlings (left) and roots (right).

Notes: A (control), B (*Racocetra crispera*), C (*Glomus intraradices*), D (*Glomus* sp.), E (*Glomus* sp.-LW10), F (*Glomus* sp.-SW10), and G (Mycofer IPB).

Table 4. Dry weight, mycorrhizal dependency (MD), the mycorrhizal growth responses (MGR), and shoot root ratio (SRR), of *A. saponaria* seedlings three months old, grown with or without mycorrhizal fungi under greenhouse conditions.

Treatment	Dry weight (g)			MD (%)	MGR (%)	SRR
	Roots	Shoots	Total			
Post-asphalt mining media						
Control	0.04±0.004 b	0.09±0.012 d	0.13±0.016 c	-	-	2.54±0.104 cd
<i>Gigaspora crispera</i>	0.15±0.01 b	0.69±0.08 bc	0,85±0.09 b	84.78±6.29	553.8±34.5	4.53±0.36 b
<i>Glomus intraradices</i>	0.93±0.39 a	1.18±0.05 a	2,11±0.43 a	92.73±3.96	1,523.1±81.3	1.65±0.45 d
<i>Glomus</i> sp.	0.24±0.06 b	0.59±0.09 bc	0,83±0.15 b	80.53±7.35	538.5±17.7	2.62±0.24 c d
<i>Glomus</i> sp.-LW10	0.14±0.02 b	0.51±0.09 c	0,65±0.10 bc	80.30±7.84	376.9±21.3	3.53±0.34 c
<i>Glomus</i> sp.-SW10	0.27±0.05 b	0.88±0.19 b	1,15±0.24 b	77.29±17.22	784.6±30.7	3.16±0.11 c
Mycofer IPB	0.09±0.01 b	0.54±0.06 c	0,61±0.07 bc	73.83±10.68	369.2±58.7	5.95±0.40 a
Pr>F	0.0134	<.0001	0.0003	-	-	<.0001
Post-nickel mining media						
Control	0.07±0.01 c	0.07±0.01 d	0.14±0.01 d	-	-	0.96±0.04 b
<i>Gigaspora crispera</i>	0.90±0.01 bc	0.10±0.02 cd	0.19±0.03 cd	26.3±12.1	35.7±20.6	1.10±0.05 ab
<i>Glomus intraradices</i>	0.35±0.04 a	0.47±0.02 a	0.82±0.06 a	82.9±1.00	485.7±35.6	1.35±0.10 a
<i>Glomus</i> sp.	0.14±0.02 b	0.13±0.02 cd	0.27±0.03 c	48.1±10.2	92.8±41.3	0.95±0.02 b
<i>Glomus</i> sp.-LW10	0.12±0.01 bc	0.16±0.02 c	0.27±0.03 c	48.1±3.90	92.8±15.2	1.35±0.08 a
<i>Glomus</i> sp.-SW10	0.14±0.01 b	0.16±0.02 c	0.30±0.02 c	53.3±4.30	114.3±17.5	1.19±0.19 ab
AMF Mixed	0.33±0.01 a	0.31±0.04 b	0.64±0.05 b	78.1±3.00	357.1±70.1	0.94±0.10 b
Pr>F	<.0001	<.0001	<.0001	-	-	0.0270

Notes: Average values followed by different letters in the same column are significantly different at DMRT (p<0.05); *Mean±SE.

Table 5. N, P, and Zn content and uptake of *A. saponaria* plant cultivated with or without AMF in post-asphalt mining media at nursery conditions after three months.

Treatment	Content (mg g ⁻¹)			Uptake (mg plant ⁻¹ x 10 ⁻³)		
	N	P	Zn	N	P	Zn
Control	2.33±0.067 b	0.00027±0.000009 c	0.00047±0.000268	290±44.3 c	0.037±0.005 b	0.033±0.023
<i>Racocetra crispera</i>	2.93±0.133 b	0.00052±0.000036 ab	0.00100±0.000202	2,510±379.0 b	0.447±0.078 b	0.867±0.273
<i>Glomus intraradices</i>	4.17±0.819 a	0.00069±0.000113 a	0.00093±0.000140	9,430±790.7 a	1.560±0.586 a	1.967±0.235
<i>Glomus</i> sp.	2.33±0.088 b	0.00046±0.000142 bc	0.00113±0.000230	1,960±435.7 b	0.420±0.208 b	1.033±0.395
<i>Glomus</i> sp.-LW10	2.13±0.033 b	0.00032±0.000013 bc	0.00146±0.000286	1,390±241.1 bc	0.213±0.043 b	1.00±0.355
<i>Glomus</i> sp.-SW10	2.30±0.058 b	0.00037±0.000038 bc	0.00176±0.000878	2,670±603.4 b	0.440±0.123 b	2.300±1.347
Mycofer IPB	2.23±0.033 b	0.00033±0.000018 bc	0.00086±0.000150	1,390±138.3 bc	0.210±0.032 b	0.5667±0.162
Pr>F	0.0053	0.0142	0.3959	<.0001	0.0116	0.1551

Table 6. N, P, and Ni content and uptake of *A. saponaria* plant cultivated with or without AMF in post-nickel mining media at nursery conditions after three months.

Treatment	Content (mg g ⁻¹)			Uptake (mg plant ⁻¹)		
	N	P	Ni	N	P	Ni
Control	0.28±0.0088	0.56±0.0491	0.053±0.002 4c	0.040±0.0048 c	0.080±0.0121 b	0.0076±0.0007 c
<i>Racocetra crispera</i>	0.29±0.0033	0.73±0.1665	0.066±0.0030 c	0.055±0.0075 c	0.147±0.0533 b	0.0126±0.0016 c
<i>Glomus intraradices</i>	0.30±0.0176	0.99±0.2734	0.191±0.0426 a	0.252±0.0315 a	0.840±0.2786 a	0.1620±0.0483 a
<i>Glomus</i> sp.	0.29±0.0088	0.72±0.0907	0.067±0.0120 c	0.077±0.0113 c	0.187±0.0101 b	0.0186±0.0054 c
<i>Glomus</i> sp.-LW10	0.28±0.0088	0.83±0.0404	0.078±0.0212 c	0.078±0.0095 c	0.229±0.0359 b	0.0221±0.0078 c
<i>Glomus</i> sp.-SW10	0.29±0.0033	0.71±0.2184	0.103±0.0023 bc	0.085±0.0067 c	0.204±0.0479 b	0.0306±0.0025 c
AMF Mixed	0.30±0.0088	1.12±0.1660	0.155±0.0175 ab	0.191±0.0143 b	0.713±0.0986 a	0.1015±0.0188 b
Pr>F	0.5601	0.2991	0.0015	<.0001	0.0015	0.0005

Table 7. The increase/decrease of nutrient uptake in *A. saponaria* seedlings.

Treatment	Increase/decrease of nutrient uptake (%)			Treatment	Increase/decrease of nutrient uptake (%)		
	N	P	Zn		N	P	Ni
Post-asphalt mining media				Post-nickel mining media			
Control	-	-	-	Control	-	-	-
<i>Racocetra crispera</i>	765.5	1,108.1	2,527.3	<i>Racocetra crispera</i>	37.5	83.8	65.8
<i>Glomus intraradices</i>	3,151.7	4,116.2	5,860.6	<i>Glomus intraradices</i>	530.0	950.0	2,031.6
<i>Glomus</i> sp.	575.9	1,035.1	3,030.3	<i>Glomus</i> sp.	92.5	133.8	144.7
<i>Glomus</i> sp.-LW10	379.3	475.7	2,930.3	<i>Glomus</i> sp.-LW10	95.0	186.3	190.8
<i>Glomus</i> sp.-SW10	820.7	1,089.2	6,869.7	<i>Glomus</i> sp.-SW10	112.5	155.0	302.6
Mycofer IPB	379.3	467.6	1,617.3	AMF Mixed	377.5	791.3	1,235.5

For the variables of Ni content and N, P, and Ni uptake, the *G. intraradices* treatment did not differ significantly from the other treatments except for the AMF mixed treatment for the variables of Ni content and p uptake (Table 6). AMF inoculation increased the uptake of P, N, Zn, and Ni in *A. saponaria* seedlings under nursery conditions (Table 7). In post-asphalt mining media, the increase in N, P, and Zn uptake ranged from 379.3-3,151.7%, 467.6-4,116.3%, and 1,517.3-5,860.6%, respectively. The increase in N, P, and Ni uptake by *A. saponaria* plants on post-nickel mining media was 37.5-530%, 83.8-950%, and 65.8-2,031.6%, respectively.

The results of this study indicate that AMF inoculation, especially *G. intraradices*, can improve the quality of *A. saponaria* seedlings in degraded soils, both in post-asphalt mining and post-nickel mining media. The effectiveness of *G. intraradices* is likely due to its ability to expand plant root networks via external hyphae, thereby increasing the efficiency of water and nutrient absorption, particularly phosphorus and nitrogen (Tian et al., 2023). Research by Barea et al. (2016) shows that *G. intraradices* has a high capacity to increase phosphorus availability in nutrient-poor soils. In addition, *G. intraradices* has been shown to increase plant resistance to environmental stress, with high adaptability and sporulation (Jansa et al., 2008; Bending et al., 2014; Ouhaddou et al., 2025). This study also supports these findings by showing significant increases in growth, dry weight, and nutrient uptake in plants inoculated with *G. intraradices* in post-mining media.

A. saponaria seedlings inoculated with AMF on post-asphalt mining and post-nickel mining media had dry weights approximately 3-15 and 0.35-4 times greater than the control, respectively. These results are consistent with those of Amir et al. (2018), who found that *Metrosideros laurifolia* mycorrhizal seedlings had dry weights approximately 4 times those of the control on serpentine soil media. Increased dry weight of mycorrhizal plants on serpentine press has also been reported in *Sorghum vulgare* (Amir et al., 2012). Tuheteru et al. (2021) reported that the application of local FMA increased the total dry weight of three-month-old *A. saponaria* seedlings grown on red-yellow podzolic soil (Ultisol) by 11 times compared to the control.

Increased growth, root nodules, and dry weight of plants are closely related to increased absorption of phosphorus and other nutrients N and Zn in mycorrhizal plant tissues (Table 5). *A. saponaria* seedlings inoculated with AMF on post-asphalt mining and post-nickel mining media had increased nutrient uptake of N by 0.35-4, P (3-31), and Zn (16-68), as well as N (0.37-5.3) and P 0.83-9.5 times greater than the control. One mechanism of P uptake is through higher production of phosphatase enzymes in the rhizosphere, thereby mobilizing inorganic P into a form available to plants. The detailed mechanism of P uptake by AMF has been reviewed by Wang (2017).

This research agrees with several previous research results on various media conditions of post-mining land, such as high P uptake in *Pterocarpus indicus* and *Vitex cofassus* plants in post-asphalt mining media conditions (Husna et al., 2025; Tuheteru et al., 2025), *K. celebica* plants in gold tailings media (Husna et al., 2021), *P. mooniana* in post-nickel mining media conditions (Husna et al., 2016), and *Amygdalus pedunculata* in coal mining (Bi et al., 2018). In addition, FMA reduces metal toxicity by immobilizing metals on hyphal walls and vesicles and by increasing plant antioxidant synthesis (Begum et al., 2019; Wang et al., 2024). This explains the ability of *A. saponaria* mycorrhizal seedlings to grow better even in nutrient-poor media. In addition to improving growth, AMF colonization also increases *A. saponaria* plants' dependence on local FMA inoculation.

In this study, the MD values ranged from 73.83% to 92.73% (post-asphalt mining media) and 26% to 82% (post-nickel mining media), with the highest MD values generally observed in the *G. intraradices* treatment. The results of this study indicate that locally isolated AMF from post-asphalt mining land can enhance the growth of *A. saponaria* plants under less favorable conditions. Compatibility of native AMF with *A. saponaria* is possible due to several factors, namely 1) native AMF is suitable for media conditions; 2) local AMF is compatible with root exudates produced by *A. saponaria* roots; and 3) genotypically, local AMF can absorb and conduct water and nutrients to the host plant (Tuheteru et al., 2025). The results of this study indicate that *A. saponaria* is highly dependent on *G. intraradices*. High dependence (MIE 91%) of *A. saponaria* was also reported by Tuheteru et al. (2021).

Based on these results, *A. saponaria* nurseries at the nursery scale need to be equipped with *G. intraradices* or other AMF. Several similar studies report that inoculation with local AMF increases photosynthesis rates, available P, and seedling survival when transferred to the field (Husna et al., 2021; Wulandari et al., 2024). Thus, the application of local AMF at the nursery stage is an efficient bioamelioration approach to improve seedling quality and accelerate revegetation success on post-mining degraded land. The results of this study also emphasize the importance of selecting local AMF isolates from extreme ecosystems, as these isolates have adaptive capabilities to the physical and chemical conditions of the local soil. Native AMF from post-asphalt mining land, specifically *G. intraradices*, has been shown to increase nutrient uptake and growth of *A. saponaria* compared to non-mycorrhizal plants. This supports the concept that specific symbiosis between fungi and host plants plays a key role in colonization effectiveness (Brundrett and Tedersoo, 2018). Therefore, integrating native AMF inoculation technology into forestry seedling programs is a strategic step to strengthen the sustainability of mine ecosystem restoration in Indonesia.

Conclusion

Native AMF inoculation has been shown to increase initial growth, dry weight, and uptake of N, P, Zn, and Ni in *Albizia saponaria* seedlings planted in post-asphalt mining and post-nickel mining media. *Glomus intraradices* isolates had the most significant effect on growth and nutrient uptake efficiency compared to other treatments. The application of native AMF from post-asphalt mining land has been proven effective in improving seedling quality and resistance to degraded environmental conditions. These findings support the use of native AMF and *A. saponaria* as an essential component in sustainable post-mining land restoration strategies in Indonesia.

Acknowledgements

This research was financially supported by the Ministry of Education, Culture, Research, and Technology of the Republic of Indonesia (No. 08/UN29.20/PG/2024) and the Ministry of Higher Education, Science, and Technology of the Republic of Indonesia (No. 24/UN29.20/PG/2025). They also extend their thanks to the Head of Mining Engineering of PT. WIKA Bitumen, Buton, Southeast Sulawesi Province, and to the Head of Mining Engineering of PT. Pernick Sultra, Konawe Utara, Indonesia.

References

- Albasri, Tuheteru, F.D., Husna, Budiharta, S., Tuheteru, E.J., Mulyono, S., Yunanto, T. and Karlin, V. 2023. Composition, structure and diversity of vegetation on asphalt post-mining sites in Buton Island, Indonesia. *Floresta Ambient* 30(3):e20230022, doi:10.1590/2179-8087-FLORAM-2023-0022.
- Amir, H., Cavaloc, Y. and Lambers, H. 2018. Arbuscular mycorrhizal fungi and sewage sludge enhance growth and adaptation of *Metrosideros laurifolia* on ultramafic soil in New Caledonia: A field experiment. *Mycorrhiza* 28(7):629-639, doi:10.1007/s00572-018-0852-7.
- Amir, H., Ducouso, M. and Cavaloc, Y. 2012. Arbuscular mycorrhizal fungi from New Caledonian ultramafic soils improve tolerance to nickel of endemic plant species. *Mycorrhiza* 22(8):631-642, doi:10.1007/s00572-018-0852-7.
- Barea, J.M., Pozo, M.J. and Azcón-Aguilar, C. 2016. Microbial management of rhizosphere and mycorrhizosphere: Exploring the role of beneficial microorganisms in improving soil quality and plant growth. *Soil Biology and Biochemistry* 103:73-85, doi:10.1016/j.soilbio.2016.08.015.
- Begum, N., Qin, C., Ahanger, M.A., Raza, S., Khan, M.I., Ashraf, M., Ahmed, N. and Zhang, L. 2019. Role of arbuscular mycorrhizal fungi in plant growth regulation: Implications in abiotic stress tolerance. *Frontiers in Plant Science* 10:1068, doi:10.3389/fpls.2019.01068.
- Bending, G.D., Turner, M.K. and Rayns, F. 2014. Effects of mycorrhizal fungi on plant growth under stress conditions. *Plant and Soil* 376(1-2):119-130, doi:10.1007/s11104-014-2136-9.
- Bi, Y., Zhang, Y. and Zou, H. 2018. Plant growth and their root development after inoculation of arbuscular mycorrhizal fungi in coal mine subsided areas. *International Journal of Coal Science and Technology* 5:47-53, doi:10.1007/s40789-018-0201-x.
- Brundrett, M.C. and Tedersoo, L. 2018. Evolutionary history of mycorrhizal symbioses and global host plant diversity. *New Phytologist* 220(4):1108-1115, doi:10.1111/nph.14976.
- Cavagnaro, T.R., Smith, F.A., Ayling, S.M. and Smith SE. 2003. Growth and phosphorus nutrition of a Paris-type arbuscular mycorrhizal symbiosis. *New Phytologist* 157(1):127, doi:10.1046/j.1469-8137.2003.00654.x.
- de Moura, M.A., Oki, Y., Arantes-Garcia, L., Cornelissen, T., Nunes, Y.R.F. and Fernandes, G.W. 2022. Mycorrhiza fungi application as a successful tool for worldwide mine land restoration: current state of knowledge and the way forward. *Ecological Engineering* 178, doi:10.1016/j.ecoeng.2022.106580.
- Fern, K. 2024. *Albizia saponaria*. Useful Tropical Plants Database. Retrieved from <https://tropical.theferns.info/viewtropical.php?id=Albizia+saponaria>
- Hikmat, A., Zuhud, E.A.M., Al Manar, P., Wilyan, R., Munggaran, I.K., Hidayat, S. and Robika. 2025. Diversity and utilization of plants in kelekak agroforestry in Air Mesu Village, Central Bangka District, Indonesia. *Biodiversitas Journal of Biological Diversity* 26(3):1105-1113, doi:10.13057/biodiv/d260309.
- Husna, Tuheteru, F.D. and Arif, A. 2021a. Arbuscular mycorrhizal fungi to enhance the growth of tropical endangered species *Pterocarpus indicus* and *Pericopsis mooniana* in post-gold mine field in Southeast Sulawesi, Indonesia. *Biodiversitas Journal of Biological Diversity* 22(9):3844-3853, doi:10.13057/biodiv/d220930.
- Husna., Tuheteru, F.D. and Arif, A. 2021b. The potential of arbuscular mycorrhizal fungi to conserve *Kalappia celebica*, an endangered endemic legume on gold mine tailings in Sulawesi, Indonesia. *Journal of Forestry Research* 32: 675-682, doi:10.1007/s11676-020-01097-8.
- Husna., Tuheteru, F.D., Nurdin, W.R., Ramadin. and Arif A. 2025. Enhanced growth and nutrient uptake of *Pterocarpus indicus* in asphalt post-mining soil media using local arbuscular mycorrhizal fungi. *Biodiversitas Journal of Biological Diversity* 26:1956-1963, doi:10.13057/biodiv/d260444.
- Jansa, J., Kolar, J. and Stribny, J. 2008. Effects of mycorrhizae on soil nutrient uptake in heavy metal contaminated soils. *Plant and Soil* 302(1-2):53-64, doi:10.1007/s11104-007-9491-2.
- Miswanto, A., Rochani, S., Dani, U., Haryadi, H., Suherman, I., Suseno, T., Astuti, W., Wijayanti, R., Prakosa, A., Prakosa, A., Hartono, H., Prameswara, G. and Petru, H.T.B.M. 2025. Buton asphalt: prospect and strategy for asphalt import substitution in Indonesia. *Mineral Economics* 38:119, doi:10.1007/s13563-024-00459-6.
- Nasution, M.J., Tugiyono, Bakri, S., Setiawan, A., Murhadi, Wulandari, C. and Wahono, E.P. 2024. The impact of increasing nickel production on forest and environment in Indonesia: A review. *Jurnal Sylva Lestari* 12(3):549-579, doi:10.23960/jsl.v12i3.847.
- Ouhaddou, R., Amrani, S., Fikri, M. and El Hajjami, M. 2025. The importance of the *Glomus* genus as a potential bioremediation tool in heavy metal stress. *Agronomy* 16(1):32, doi:10.3390/agronomy16010032.
- Pratiwi, Narendra, B.H., Siregar, C.A., Turjaman, M., Hidayat, A., Rachmat, H.H., Mulyanto, B., Suwardi, Iskandar, Maharani, R., Rayadin, Y., Prayudyarningsih,

- R., Yuwati, T.W., Prematuri, R. and Susilowati, A. 2021. Managing and reforesting degraded post-mining landscape in Indonesia: A Review. *Land* 10(6):658, doi:10.3390/land10060658.
- Prematuri, R., Turjaman, M., Sato, T. and Tawaraya, K. 2020. The impact of nickel mining on soil properties and growth of two fast-growing tropical tree species. *International Journal of Forestry Research* 2020(19):1-9, doi:10.1155/2020/883759.
- Soil Research Center. 2009. *Technical Guidelines for Chemical Analysis of Soil, Plants, Water, and Fertilizers*. Second edition. Bogor (in Indonesian).
- Tian, Y., Xu, J., Lian, X., Wei, B., Ma, X. and Wu, P. 2023. Effect of *Glomus* intraradices on root morphology, biomass production and phosphorus use efficiency of Chinese fir seedlings under low phosphorus stress. *Frontiers in Plant Science* 13:1095772, doi:10.3389/fpls.2022.1095772.
- Tuheteru, F.D., Husna, Albasri, Effendy, H.M., Arif, A., Basrudin, Tuheteru, E.J., Mulyono, S. and Irianto, R.S. 2022. Diversity of arbuscular mycorrhizal fungi in asphalt post-mining land in Buton Island, Indonesia. *Biodiversitas Journal of Biological Diversity* 23(12):5290-5297, doi:10.13057/biodiv/d231038.
- Tuheteru, F.D., Husna, and Alimuddin, L.D. 2021. Response of growth and dependency level of *Albizia saponaria* (Lour.) local arbuscular mycorrhizae fungi from Southeast Sulawesi. *Biota: Jurnal Ilmiah Ilmu-Ilmu Hayati* 16(2):252-261, doi:10.24002/biota.v16i2.107 (in Indonesian).
- Tuheteru, F.D., Husna, Nurdin, W.R., Himawan, A., Tuheteru, E.J., Albasri, Mulyono, S. and Arif, A. 2025. Native arbuscular mycorrhizal fungi promote the growth of *Vitex cofassus* seedlings in post-asphalt mining soil media. *Journal of Degraded and Mining Lands Management* 12(2):7225-7233, doi:10.15243/jdmlm.2025.122.7225.
- van der Heijden, M.G.A., Wiemken, A. and Sanders, I.R. 2003. Different arbuscular mycorrhizal fungi alter coexistence and resource distribution between co-occurring plants. *New Phytologist* 157(3):569-578, doi:10.1046/j.1469-8137.2003.00688.x.
- Vierheilig, H., Schweiger, P. and Brundrett, M. 2005. An overview of methods for the detection and observation of arbuscular mycorrhizal fungi in roots. *Physiologia Plantarum* 125:393-404, doi:10.1111/j.1399-3054.2005.00564.x.
- Wang, F., Lin, X. and Yin, R. 2005. Heavy metal uptake by arbuscular mycorrhizas of *Elsholtzia splendens* and the potential for phytoremediation of contaminated soil. *Plant and Soil* 269:225-23, doi:10.1007/s11104-004-0517-8.
- Wang, F. 2017. Arbuscular mycorrhizas and ecosystem restoration. In: Wu, Q.S. (ed.), *Arbuscular Mycorrhizas and Stress Tolerance of Plants*. (pp 245-291). Springer Nature Link, Singapore, doi:10.1007/978-981-10-4115-0_11.
- Wang, Q., Liu, M., Wang, Z., Li, J., Liu, K. and Huang, D. 2024. The role of arbuscular mycorrhizal symbiosis in plant abiotic stress. *Frontiers in Microbiology* 14:1323881, doi:10.3389/fmicb.2023.1323881.
- Wulandari, D., Nufus, M., Faridah, E., Maulana, A.F. and Tawaraya, K. 2024. Native arbuscular mycorrhizal fungi and *Nauclea orientalis* for potential reclamation of tropical coal mining areas. *Environmental Advances* 15:100462, doi:10.1016/j.envadv.2023.100462.