	Journal Title	: <u>Jurnal Multidisiplin Madani (MUDIMA)</u>
<section-header></section-header>	Scope	: Multidisciplinary
	Language	: Indonesia and English
	Editor in Chief	: <u>Professor Jen Peng Huang</u>
	e-ISSN	: <u>2808-5639 (Online)</u>
	Frequency	: Monthly
	DOI	: Prefix 10.55927
	Office	: Jl. Ir. H. Juanda No.56B, Sukaraja, Kec. Medan Maimun, Kota Medan, Sumatera Utara 22159
	Email	: admin@formosapublisher.org
	Publisher	: Formosa Publisher
	Citation Analysis	: Google Scholar, Scopus, Dimension

Jurnal Multidisiplin Madani (MUDIMA) is dedicated to exploring and disseminating the results of various creative and innovative thoughts based on scientific research and thought processes. This Journal is a double blind review journal focus on the empirical research that provide novelty of the research with the latest issues and contributions to the scientific research.

Jurnal Multidisiplin Madani (MUDIMA) has been widely indexed by:

1. EBSCO, 2. Google Scholar, 3. Copernicus International (ICV 2021:70,71), 4. Open Aire 5. GARUDA, 6. <u>ScienceGate</u> 7. <u>Directory of Research Journals Indexing</u> , 8. <u>SHERPA/Romeo</u> , 9. <u>RESEARCH BIBLE</u> , 10. DIMENSIONS , 11. WorldCat , 12. SciLit , 13. CiteFactor, 14. Semantic Scholar , 15. Eurasian Scientific Journal Index , 16. IP Indexing 17. Root Indexing , 18. IUPUI Libraries , 19. Scite , 20. Neliti , 21. Advanced Sciences

Index (ASI), 22. Harvard Library, 23. IP Indexing 24. Indonesia OneSearch 25. Base (bielefeld academic search engine) 26. Fatcat , 27. Exaly, 28. Stanford Libraries

Other Libraries:



Complete indexing and libraries information can be checked: https://journal.formosapublisher.org/index.php/mudima/indexing

Vol. 3 No. 9 (2023): September, 2023



Published: 2023-09-29

Editorial Team

EDITOR IN CHIEF:



Professor Jen Peng Huang, Ph.D

Email: jenpeng.huang@gmail.com Department of Information Management, Southern Taiwan University of Science and Technology

Editorial Board





Dr. Aaron Raymond See

Lusius Sinurat, SS, M.Hum Pena Sinergi, Indonesia

Southern Taiwan University of Science and Technology, Taiwan



Graduate school, Dusit Thani College, Bangkok, Thailand



Lê Thị Bích Ngọc, Ph.D



Ari Kristanta Tarigan S.Sn., M.Sn

Formosa Publisher

Editorial Members

Dr. Uma Shankar Yadav

Allahabad Pryagraj, India

Dr. Cris Norman P. Olipas

Nueva Ecija University of Science and Technology, Philippines

Dr. Kamran Abdullayev

Formosa Publisher

Irwan Ginting S.Th., M.Pd.K

Motilal Nehru National Institute of Technology Institute of Economics of Azerbaijan National Academy of Sciences, Azerbaijan

Vinay Singh

UPES, India

Assistant Editor

Marisa Yunita br Depari, S.Pd

E-Mail: journalmudima@gmail.com

WhatsApp: <u>+6287837338057</u>

Sisnawati Br Sitepu, S.Pd

E-Mail: admin@formosapublisher.org

WhatsApp: +6281360721819



Jurnal Multidisiplin Madani (MUDIMA)

Homepage: https://journal.formosapublisher.org/index.php/mudima

ISSN: 2808-5639 (Online) Research Article



Volume 3, No 9, September (2023)

DOI: https://doi.org/10.55927/mudima.v3i9.3384

Effect of Coffee Consumption with Muscle Strength in Male Adolescents

Futsal Team

Ayu Astianing Tiyas¹, Eveline Margo^{2*} Universitas Trisakti, Jakarta **Corresponding Author**: Eveline Margo eveline-margo@trisakti.ac.id

ARTICLEINFO

Keywords: Caffeine, Muscle Strength, Vertical Jump Test, Male Adolescents

Received: 2 AugustRevised: 19 AugustAccepted: 20 September

©2023 Tiyas, Margo: This is an openaccess article distributed under the terms of the <u>Creative Commons Atribusi 4.0</u>



ABSTRACT

In the realm of sports, caffeine has become a popular choice for athletes seeking to enhance their strength and maintain peak performance levels. Research has demonstrated that caffeine can facilitate the conversion of fat into energy and increase calcium levels within muscle cells, thereby improving muscle performance and reducing fatigue. This study was conducted to examine the impact of coffee consumption on muscle strength in a male high school futsal team. Employing an experimental method involving a one-group pre-post test design, the study included 22 teenagers. This research was conducted at SMA 5 Tambun Selatan, Bekasi. The researchers utilized the vertical jump test to gauge leg muscle strength both before and after administering coffee to participants and analysed use paired t-test. The result is increased muscle strength after giving coffee at a dose of 2mg CAF/kgBW (51.45 ± 8.92) and 3mg CAF/kgBW (54.05 ± 8.81) compared to before giving coffee (46.68 ± 9.99). There is relationship between coffee consumption and increased muscle strength (p=0.00)

INTRODUCTION

Adolescence is a transition from childhood to adulthood which is marked by physical growth and development as well as psychological changes. World Health Organization says adolescents are population with an age range from 10 to 19 years old.(Success Stories South-East Asia Region, 2021) Based on Indonesian Republic Health Rules number 25, 2014, adolescents are citizens in the group of age 10 to 18 years, whereas according to BKKBN, the age range for adolescents is 10 to 24 years old and married.(Kementrian aren't Kesehatan RI, 2014). The transition from childhood to adulthood involves a lot of changes, including in growth and development both physical, sexual, mental, and social. (Mushaddik et al., 2022; Success Stories South-East Asia Region, 2021; Suharyanti & Suharyanti, 2021) Adolescents's growth and development are triggered by several factors, one of which is physical activity. Sport has become an important factor that is expected to improve physical and spiritual fitness in adolescents.(Abdulaziz et al., 2016)

Physical fitness can be improved through physical activity by doing sports. (Nandatama et al., 2017) Alvara Research Center national survey in Jakarta, 2018, said that of 1.200 millennial generation in Indonesia, 72.9% liked sports, of which 44.5% chosed football or futsal followed by running at 18.6%, badminton at 18.1%, cycling at 13.9% and the rest for other sports.(Databoks, 2018)

Futsal is a sport that played in closed field and requires 2 x 20 minutes of game. Because of that, athletes need good physical fitness. Good physical training must include flexibility, strength, power and endurance. It is necessary to be given several training maintain programs to or improve player performance, for example increasing muscle strength. Muscle strength can be built through progressive weight training, such as push-ups, situps, squat jumps .(Tang et al., 2020)

Around 80% of the world's population consumes caffeine every day. Caffeine is easy to find in the market in different forms, such as tea, coffee, cocoa, energy drinks, pills, and many other beverages. Recent research studies showed that caffeine has positive effects on sport performance. The main component of coffee is caffeine, which is an ergogenic substance that can work as stimulant so it can improve several aspects of performance and prevent fatigue.(Burke, 2008; Nandatama et al., 2017; Samoggia & Rezzaghi, 2021)

Based on studies, coffee drinks that contain caffeine with 4.5-5.5 mg/kgBW can help to improve football player's physical performance, including the strength and muscle endurance.(Mielgo-Ayuso et al., 2019) The International Olympic Committee claims that the safe dose for caffeine intake was 3 mg up to 6 mg CAF/kgBW to obtain ergogenic effects.(Maughan et al., 2018) The current study by Tangen DS et al. declared that giving 3 mg/kgBW of caffeine can increase the height of vertical jump at rest state, (Tangen et al., 2020) and according to Nandatama R et al., giving coffee at dose 2 gram/150 ml which consumed one hour before exercise can increase the strength and endurance football athletes.(Nandatama et al., 2017) Contrarily, research conducted by Wilk M et al., administering high doses of caffeine (9 mg/kgBW and 11 mg/kgBW) was unable to provide ergogenic effects on individuals that addicted to coffee, even though the dose was given exceeded the average daily dose.(Wilk et al., 2019) The study that conducted by Desbrow B et al., also declare that caffeine administration did not have a significant effect on endurance.(Desbrow et al., 2012) Based on the variation results above, this study aims to search the effect between coffee consumption and muscle strength in male high school futsal teams.

METHODS

This experimental study is a one-group prepost-test design which was held in December 2021 in SMA 5 Tambun Selatan, Bekasi. The subjects were 22 male teenagers which was selected by nonrandom sampling. The inclusion for this study were:1) male teenager who studied in SMA 5 Tambun Selatan, Bekasi. 2) aged 15-18 years. 3) played futsal. 4) have a healthy body with 36.50C-37.50C temperature body, blood pressure $\leq 120/80$ mmHg, pulse 60-100 x/minute, oxygen saturation 95-100% 5) be willing to participate in this study and sign informed consent. Meanwhile, the exclusion criteria were: 1) having a history of gastritis, ulcus pepticum, heart and lung disease 2) taking food or drinks containing caffeine at least 5 days before this study, such as coffee, tea, chocolate, fizzy drinks, and foods that containing coffee, tea, cocoa and so on 3) taking tobacco consumption 4) suffered injury.

Subjects were interviewed, and after that, a psyhical examination was carried out including body weight, body temperature, blood pressure, oxygen saturation using oxymeter and pulse rate. Body weight is measured using digital scales to determine the amount of coffee consumption. Vertical jump test was used to measure leg muscle strength. This study uses Nescafe Classic®, which contains 100% pure robusta coffee with one caffeine content range from 80 to 90 mg/2gr Nescafe Classic®. (Nestle, 2022)

Subjects will be tested for muscle strength before and after being given coffee. Before being given a coffee drink, subjects will take the vertical jump test 3 times and the highest score will be taken. After that, subjects will be given Nescafe classic® with a dose of 2mg CAF/kgBW on the first day and 3mg CAF/kgBW on the sixth day, which each will be dissolved in 150 ml of water. The two main tests were separated by 6 days and the subjects were not allowed take a food or drinks that containing coffee also no exercise was allowed. Caffeine has an elimination half-life, which is around 2 to 12 hours, with an average of 5 hours in the body, and according to Republic Indonesia Food and Drug Control Agency (BPOM RI), the washout period is usually more than five times of the terminal half-life of the drug, so the researchers carried out washout period for 5 days. (Mandal, 2023; Sampurno, 2005)

After drinking the coffee, the subjects will do the vertical jump test 3 times again at the 30th, 45th and 60th minutes, and the highest score will be taken. The assessment is done by calculating the difference between the subject's hand reach while standing upright and when jumping. In the previous study, there was an increase in leg muscle strength with vertical jump test if the jump height difference before and after drinking coffee was $\geq 2.2 \pm 0.5$ cm. (Wilk et al., 2019)

RESULTS AND DISCUSSION

Variable	n (%)	Mean ± SD	
Age		16.55 ± 1.01	
15 years	3 (13.6%)		
16 years	9 (40.9%)		
17 years	5 (22.7%)		
18 years	5 (22.7%)		
Leg muscle strengt	h		
Before drinking co	ffee	46.68 ± 9.99	
After drinking coff	èe,		
the dose:			
2 mg CAF	/kgBW	51.45 ± 8.92	
3 mg CAF	/kgBW	54.05 ± 8.81	

Table 1. Research Subjects Characteristics (n=22)

Data table 1 shows the age mean subjects was 16.55 ± 1.01 years with the largest age distribution was 16 years old (40.9%). The averange leg muscle strength before dringking coffee was 46.68 ± 9.99 , whereas after drinking coffee with a dose 2mgCAF/kgBW was 51.45 ± 8.92 and at a dose 3mg CAF/kgBW was 54.05 ± 8.81 .

A go	Leg Muscle Strength		р-
Age	Increase (n)	Not Increase (n)	value
15-16 years	11	1	0.000*
17-18 years	9	1	0.892*

Table 2 Polationship Age and Musels Strength

Table 3. Relationship Coffee Consumption with Muscle Strength

Caffaina dasa	Leg Muse	n valua	
Callelle uose	Before	After	- p-value
2mg CAF/kgBW	46.68±9.99	51.45±8.92	0.00*
3mg CAF/kgBW	46.68±9.99	54.05 ± 8.81	0.00*
*: Paired t-test			

The analysis using paired t-test showed that there was a significant relationship between coffee consumption with muscle strength with a p-value 0.00 (p < 0.05).



Figure 1. Leg Muscle Strength Before and After Coffee Consumption

Based on figure 1, there was a significant increase in muscle strength after giving coffee at dose 2mg CAF/kgBW or dose 3mg CAF/kgBW which was measured 3 times at the 30th, 45th and 60th minutes. This increase of leg muscle strength can be seen continuing at every minute. At dose 2mg CAF/kgBW, it's shown that the average at the 30th minute was 47.95, at the 45th minute was 49.63 and the 60th minute was 51.45. Meanwhile at dose 3mg CAF/kgBW, the average at the 30th minute was 48.54, at the 45th minute was 50.95 and at the 60th minute was 54.05. In this study, all subjects were male, because there were no females in the futsal extracurricular at SMA 5 Tambun, Bekasi. The same results were also obtained according to another study. It was said that female participation is low because of the stigma that futsal is an activity that has violent and courageous characteristics that are identical to the male's characteristics. (Akurat & Maksum, 2021)

In our results, there was no relationship between age and muscle strength because all of the subjects were in the teenage age category. One of the risk factors that can influence muscle strength is age. Physicologically, peak mass growth and muscle strength are gained in late adolescence. Meanwhile, muscle mass decreases around 3 to 8% at the age of 30 years and this decreases even higher above the age of 60 years old. (Ng et al., 2019; Volpi et al., 2004)

There was a relationship between coffee consumption and leg muscle strength in both doses of caffeine p=0.00 (p<0.05). This study results showed an increase in the total leg muscle strength average before consuming coffee (46.68±9.66) and after consuming coffee with a dose of 2mg CAF/kgBW and 3mg CAF/kgBW (54.05±8.81) using vertical jump test. Coffee contains purine alkaloids called caffeine (1,3,7-trimethylxanthine) which bind and regulate several protein that has physiological effects and stimulates sympathetic nervous system as an adenosine receptor antagonist.(Ludwig et al., 2014; Wishart et al., 2017)

Caffeine is one of the ergogenic substances that can increase muscle strength. Caffeine also can help improve sport performance, and muscle performance especially to increase aerobic activity endurance as well as increasing the ability to repetitions or repeat movements in muscle training. This is due to the effects of caffeine which can inhibit the muscle Ca2+ reuptake by sarcoplasmic reticulum thus reducing excitability threshold value and making muscle contraction last longer. Caffeine increases C-AMP by inhibiting the activity of cyclic nucleotide phosphodiesterase, so the result is increasing free fatty acids (FFA) that can be used as energy and causes glycogen saving effects for endurance performance in the long term. Caffeine has a structure similar to adenosine, acting as a competitive antagonist at the adenosine receptor causing low adenosine concentrations in the body, resulting in increased motor activity and lipolysis inhibition to increases FFA production so it can be used as energy for muscle contraction. In addition, caffeine causes the epinephrine release from the adrenal gland causing lipolysis in muscle and fat tissues as an energy sources. As a results of the increasing FFA, the use of glycogen at the beginning of exercise decreases. (Hayati, 2012; Samoggia & Rezzaghi, 2021; Spriet, 2014)

Caffeine as an adenosine receptor antagonist (A1 and A2) on the skeletal muscle membrane make increases exicitation-contraction coupling through the release of Ca2+ from the sarcoplasma reticulum and improves Na+-K+ATPase pump activity which helps increase muscle performance. Adenosine has the effect to inhibiting lipolysis and stimulating insulin to glucose uptake which causes skeletal muscle contraction. In addition, because it is accompanied by an increase in adenosine consentration during exercise, it can increase glycogenolysis in both working and non working tissue also skeletal adipose tissue during physical activity. Futhermore, caffeine has been proven to improve the force velocity relationship and the fibre's muscle conduction velocity. (Hodgson et al., 2013; Tangen et al., 2020)

In this study, it was found that muscle power performance increased every minutes in both doses, 2mg CAF/kgBW and 3mg CAF/kgBW, which the height occurring at the 60th minute. Recently, some studies said the low caffeine dose administration (3mg/kgBW) also can produce an ergogenic effect, with no changes in heart rate, catecholamines level, lactate, FFA and glycerol during exercise. Every minute the muscle strength increases because of the associated caffeine absorption by the small intestine. Approximately 0.5-2% of the caffeine is excreted by urine and the rest will be absorbed by the body through the small intestine. After oral caffeine consumption, the working time will be reached within 45-60 minutes, where the maximum concentration in plasma will be reached between 30-120 minutes. The increase in muscle strength for each respondent was different due to each individual's physiological digestive system differences which depended on the time of gastric emptying and food consumption. The elimination of caffeine half-life was around 2 to 12 hours, with an average 5 hours. It's been reported that women had shorter half-life compared to men. (Grzegorzewski et al., 2022; Mandal, 2023; Spriet, 2014; Wishart et al., 2017)

Studies conducted by Nandatama R et al. and Tangen DS et al. had the same results with this study, which showed that there was an increase in muscle strength after giving coffee, even though it was said that it could increase athlete's endurance.(Nandatama et al., 2017; Tangen et al., 2020)

CONCLUSION

There was a significant relationship between coffee consumption and muscle strength in both doses. The increasing muscle strength reached to the maximum at the 60th minutes in both doses.

REFERENCES

- Abdulaziz, M. F., Dharmawan, D. B., & Putri, D. T. (2016). MOTIVASI MAHASISWA TERHADAP AKTIVITAS OLAHRAGA PADA SORE HARI DI SEKITAR AREA TAMAN SUTERA UNIVERSITAS NEGERI SEMARANG. Journal of Physical Education.
- Akurat, Y., & Maksum, A. (2021). FAKTOR-FAKTOR PENYEBAB RENDAHNYA PARTISIPASI SISWA PUTRI DALAM EKSTRAKURIKULER FUTSAL DI SMAN 18 SURABAYA. 09.
- Burke, L. M. (2008). Caffeine and sports performance. Applied Physiology, Nutrition, and Metabolism, 33(6), 1319–1334. https://doi.org/10.1139/H08-130
- Databoks, databoks. (2018, July 27). Semakin muda usia kian senang olahraga. https://databoks.katadata.co.id/datapublish/20 18/07/27/semakin-muda-usia-kian-senangolahraga
- Desbrow, B., Biddulph, C., Devlin, B., Grant, G. D., Anoopkumar-Dukie, S., & Leveritt, M. D. (2012). The effects of different doses of caffeine on endurance cycling time trial performance. Journal of Sports Sciences, 30(2), 115–120. https://doi.org/10.1080/02640414.2011.6324 31

- Grzegorzewski, J., Bartsch, F., Köller, A., & König,
 M. (2022). Pharmacokinetics of Caffeine: A
 Systematic Analysis of Reported Data for
 Application in Metabolic Phenotyping and
 Liver Function Testing. Frontiers in
 Pharmacology, 12, 752826.
 https://doi.org/10.3389/fphar.2021.752826
- Hayati, H. (2012). Penggunaan kafein dalam dunia olahraga dan efeknya sebagai ergogenik.
 WAHANA, 58(1). https://doi.org/10.36456/wahana.v58i1.1296
- Hodgson, A. B., Randell, R. K., & Jeukendrup, A. E.
 (2013). The Metabolic and Performance Effects of Caffeine Compared to Coffee during Endurance Exercise. PLoS ONE, 8(4), e59561. https://doi.org/10.1371/journal.pone.0059561
- Kementrian Kesehatan RI. (2014). Infodatin _situs reproduksi remaja. https://pusdatin.kemkes.go.id/resources/down load/pusdatin/inhttps://pusdatin.kemkes.go.id /resources/download/pusdatin/infodatin/infod atin-reproduksi-remaja.pdffodatin/infodatinreproduksi-remaja.pdf.
- Ludwig, I. A., Mena, P., Calani, L., Cid, C., Del Rio,
 D., Lean, M. E. J., & Crozier, A. (2014).
 Variations in caffeine and chlorogenic acid
 contents of coffees: What are we drinking?
 Food Funct., 5(8), 1718–1726.
 https://doi.org/10.1039/C4FO00290C
- Mandal, A. (2023, June 19). Caffeine Pharmacology. https://www.newsmedical.net/health/Caffeine-Pharmacology.aspx
- Maughan, R. J., Burke, L. M., Dvorak, J., Larson-Meyer, D. E., Peeling, P., Phillips, S. M., Rawson, E. S., Walsh, N. P., Garthe, I., Geyer, H., Meeusen, R., Van Loon, L. J. C., Shirreffs, S. M., Spriet, L. L., Stuart, M., Vernec, A., Currell, K., Ali, V. M., Budgett, R. G., ... Engebretsen, L. (2018). IOC consensus

statement: Dietary supplements and the high-
performance athlete. British Journal of SportsMedicine,52(7),439–455.https://doi.org/10.1136/bjsports-2018-099027

- Mielgo-Ayuso, J., Calleja-Gonzalez, J., Del Coso, J., Urdampilleta, A., León-Guereño, P., & Fernández-Lázaro, D. (2019). Caffeine Supplementation and Physical Performance, Muscle Damage and Perception of Fatigue in Soccer Players: A Systematic Review. Nutrients, 11(2), 440. https://doi.org/10.3390/nu11020440
- Mushaddik, I. L., Khalid, K., Anuar, A., Che Hat, S. Z., & Jamaluddin, R. (2022). Lifestyle Practices and Mental Health in Adolescents: Explorative Analysis from Malaysian Health and Morbidity Survey 2017. Adolescents, 2(4), 459–465. https://doi.org/10.3390/adolescents2040036
- Nandatama, S. R., Rosidi, A., & Ulvie, Y. N. S. (2017). Minuman Kopi (Coffea)Terhadap Kekuatan Otot dan Ketahanan Otot Atlet Sepak Bola Usia Remaja di SSB PERSISAC. JurnalGizi, 29–34.
- Nestle. (2022, January 7). Rahasia Nescafe classic. Rahasia NESCAFÉ Classic, Produk Nescafé Kopi Hitam yang Mudah Larut. https://www.nescafe.com/id/artikel/nescafeclassic-kopi-hitamlarut#:~:text=Terbuat%20dari%20Biji%20Ko pi%20Berkualitas,wilayah%20perkebunan%2 0kopi%20yang%20berkualitas
- Ng, A. K., Hairi, N. N., Jalaludin, M. Y., & Majid, H. A. (2019). Dietary intake, physical activity and muscle strength among adolescents: The Malaysian Health and Adolescents Longitudinal Research Team (MyHeART) study. BMJ Open, 9(6), e026275. https://doi.org/10.1136/bmjopen-2018-026275

- Samoggia, A., & Rezzaghi, T. (2021). The Consumption of Caffeine-Containing Products to Enhance Sports Performance: An Application of an Extended Model of the Theory of Planned Behavior. Nutrients, 13(2), 344. https://doi.org/10.3390/nu13020344
- Sampurno, H. (2005). BADAN PENGAWAS OBAT DAN MAKANAN. https://jdihn.go.id/files/491/PER%20KBPOM _NO.HK.00.05.3.1818%20TH%202005_Ten tang%20PEDOMAN%20UJI%20BIOEKIV_ 2005.pdf
- Spriet, L. L. (2014). Exercise and Sport Performance with Low Doses of Caffeine. Sports Medicine, 44(S2), 175–184. https://doi.org/10.1007/s40279-014-0257-8
- Success stories South-East Asia region: Reproductive, maternal, newborn, child and adolescent health. (2021). World Health Organization. Regional Office for South-East Asia.
- Suharyanti, S., & Suharyanti, S. (2021). Kampanye Generasi Berencana (GENRE), Sikap Generasi Z di Jakarta, dan Penetrasi Media Sosial Selama Pandemi COVID-19. Jurnal Studi Komunikasi dan Media, 25(2), 111. https://doi.org/10.31445/jskm.2021.3762
- Tang, A., Hendrik, H., & Nining Wean, I. G. (2020). PENGARUH LATIHAN FISIK TERHADAP PERUBAHAN **KEKUATAN** OTOT PEMAIN FUTSAL **CHERUBIM** FC. COMPETITOR: Jurnal Pendidikan Kepelatihan Olahraga, 12(1), 1. https://doi.org/10.26858/com.v12i1.13531
- Tangen, D. S., Nielsen, S. R., Kolnes, K. J., & Jensen, J. (2020). Caffeine Increases Vertical Jumping Height in Young Trained Males Before But Not After a Maximal Effort Strength Training Session. Journal of Science in Sport and Exercise, 2(2), 145–153. https://doi.org/10.1007/s42978-020-00060-7

- Volpi, E., Nazemi, R., & Fujita, S. (2004). Muscle tissue changes with aging: Current Opinion in Clinical Nutrition and Metabolic Care, 7(4), 405–410.
 https://doi.org/10.1097/01.mco.0000134362.7 6653.b2
- Wilk, M., Krzysztofik, M., Filip, A., Zajac, A., & Del Coso, J. (2019). The Effects of High Doses of Caffeine on Maximal Strength and Muscular Endurance in Athletes Habituated to Caffeine. Nutrients, 11(8), 1912. https://doi.org/10.3390/nu11081912
- Wishart, D., Feunang, Y., Guo, A., Lo, E., Marcu,
 A., Grant, J., Sajed, T., Johnson, D., Li, C.,
 Sayeeda, Z., Assempour, N., lynkkaran, I.,
 Liu, Y., Maciejewski, A., Gale, N., Wilson,
 M., Chin, L., Cummings, R., Le, D., ...
 Wilson, M. (2017). Caffeine (5.0). DrugBank.
 https://go.drugbank.com/drugs/DB00201

Effect of Coffee Consumption with Muscle Strength in Male Adolescents Futsal Team

by admin@formosapublisher.org _

Submission date: 18-Sep-2023 07:17AM (UTC-0400) Submission ID: 2137514603 File name: Eveline_Margo_1.docx (215.96K) Word count: 2473 Character count: 13319

Effect of Coffee Consumption with Muscle Strength in Male Adolescents Futsal Team

Ayu Astianing Tiyas¹ Eveline Margo^{2*} 1. Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia 2*. Departement of Physiology, Faculty of Medicine, Universitas Trisakti, East Jakarta, Indonesia Email: <u>eveline-margo@trisakti.ac.id</u>

ABSTRACT

In the realm of sports, caffeine has become a popular choice for athletes seeking to enhance their strength and maintain peak performance levels. Research has demonstrated that caffeine can facilitate the conversion of fat into energy and increase calcium levels within muscle cells, thereby improving muscle performance and reducing fatigue. This study was conducted to examine the impact of coffee consumption on muscle strength in a male high school futsal team. Employing an experimental method involving a one-group pre-post test design, the study included 22 teenagers. This research was conducted at SMA 5 Tambun Selatan, Bekasi. The researchers utilized the vertical jump test to gauge leg muscle strength both before and after administering coffee to participants and analysed use paired t-test. The result is increased muscle strength after giving coffee at a dose of 2mg CAF/kgBW (51.45±8.92) and 3mg CAF/kgBW (54.05±8.81) compared to before giving coffee (46.68±9.99). There is relationship between coffee consumption and increased muscle strength (p=0.00).

Keywords : caffeine, muscle strength, vertical jump test, male adolescents.

INTRODUCTION

Adolescence is a transition from childhood to adulthood which is marked by physical growth and development as well as psychological changes. World Health Organization says adolescents are population with an age range from 10 to 19 years old. (*Success Stories South-East Asia Region*, 2021) Based on Indonesian Republic Health Rules number 25, 2014, adolescents are citizens in the group of age 10 to 18 years, whereas according to BKKBN, the age range for adolescents is 10 to 24 years old and aren't married. (Kementrian Kesehatan RI, 2014) The transition from childhood to adulthood involves a lot of changes, including in growth and development both physical, sexual, mental, and social. (Mushaddik et al., 2022; *Success Stories South-East Asia Region*, 2021; Suharyanti & Suharyanti, 2021) Adolescents's growth and development are triggered by several factors, one of which is physical activity. Sport has become an important factor that is expected to improve physical and spiritual fitness in adolescents.(Abdulaziz et al., 2016)

Physical fitness can be improved through physical activity by doing sports. (Nandatama et al., 2017) Alvara Research Center national survey in Jakarta, 2018, said that of 1.200 millennial generation in Indonesia, 72.9% liked sports, of which 44.5% chosed football or futsal followed by running at 18.6%, badminton at 18.1%, cycling at 13.9% and the rest for other sports.(Databoks, 2018)

Futsal is a sport that played in closed field and requires 2 x 20 minutes of game. Because of that, athletes need good physical fitness. Good physical training must include flexibility, strength, power and endurance. It is necessary to be given several training programs to maintain or improve player performance, for example increasing muscle strength. Muscle strength can be built through progressive weight training, such as push-ups, sit-ups, squat jumps .(Tang et al., 2020)

Around 80% of the world's population consumes caffeine every day. Caffeine is easy to find in the market in different forms, such as tea, coffee, cocoa, energy drinks, pills, and many other beverages. Recent research studies showed that caffeine has positive effects on sport performance. The main component of coffee is caffeine, which is an ergogenic substance that can work as stimulant so it can improve several aspects of performance and prevent fatigue. (Burke, 2008; Nandatama et al., 2017; Samoggia & Rezzaghi, 2021)

Based on studies, coffee drinks that contain caffeine with 4.5-5.5 mg/kgBW can help to improve football player's physical performance, including the strength and muscle endurance. (Mielgo-Ayuso et al., 2019) The *International Olympic Committee* claims that the safe dose for caffeine intake was 3 mg up to 6 mg CAF/kgBW to obtain ergogenic effects. (Maughan et al., 2018) The current study by Tangen DS et al. declared that giving 3 mg/kgBW of caffeine can increase the height of vertical jump at rest state, (Tangen et al., 2020) and according to Nandatama R et al., giving coffee at dose 2 gram/150 ml which consumed one hour before exercise can increase the strength and endurance football athletes.(Nandatama et al., 2017) Contrarily, research conducted by Wilk M et al., administering high doses of caffeine (9 mg/kgBW and 11 mg/kgBW) was unable to provide ergogenic effects on individuals that addicted to coffee, even though the dose was given exceeded the average daily dose.(Wilk et al., 2019) The study that conducted by Desbrow B et al., also declare that caffeine administration did not have a significant effect on endurance.(Desbrow et al., 2012) Based on the variation results above, this study aims to search the effect between coffee consumption and muscle strength in male high school futsal teams.

METHOD

1

This experimental study is a one-group pre-post-test design which was held in December 2021 in SMA 5 Tambun Selatan, Bekasi. The subjects were 22 male teenagers which was selected by non-random sampling. The inclusion for this study were:1) male teenager who studied in SMA 5 Tambun Selatan, Bekasi. 2) aged 15-18 years. 3) played futsal. 4) have a healthy body with 36.5°C-37.5°C temperature body, blood pressure ≤120/80 mmHg, pulse 60-100 x/minute, oxygen saturation 95-100% 5) be willing to participate in this study and sign informed consent. Meanwhile, the exclusion criteria were: 1) having a history of gastritis, ulcus pepticum, heart and lung disease 2) taking food or drinks containing caffeine at least 5 days before this study, such as coffee, tea, chocolate, fizzy drinks, and foods that containing coffee, tea, cocoa and so on 3) taking tobacco consumption 4) suffered injury.

Subjects were interviewed, and after that, a psyhical examination was carried out including body weight, body temperature, blood pressure, oxygen saturation using oxymeter and pulse rate. Body weight is measured using digital scales to determine the amount of coffee consumption. Vertical jump test was used to measure leg muscle strength. This study uses Nescafe Classic®, which contains 100% pure robusta coffee with one caffeine content range from 80 to 90 mg/2gr Nescafe Classic®. (Nestle, 2022)

Subjects will be tested for muscle strength before and after being given coffee. Before being given a coffee drink, subjects will take the vertical jump test 3 times and the highest score will be taken. After that, subjects will be given Nescafe classic® with a dose of 2mg CAF/kgBW on the first day and 3mg CAF/kgBW on the sixth day, which each will be dissolved in 150 ml of water. The two main tests were separated by 6 days and the subjects were not allowed take a food or drinks that containing coffee also no exercise was allowed. Caffeine has an elimination half-life, which is around 2 to 12 hours, with an average of 5 hours in the body, and according to Republic Indonesia Food and Drug Control Agency (BPOM RI), the washout period is usually more than five times of the terminal half-life of the drug, so the researchers carried out washout period for 5 days. (Mandal, 2023; Sampurno, 2005)

After drinking the coffee, the subjects will do the vertical jump test 3 times again at the 30th, 45th and 60th minutes, and the highest score will be taken. The assessment is done by calculating the difference between the subject's hand reach while standing upright and when jumping. In the previous study, there was an increase in leg muscle strength with vertical jump test if the jump height difference before and after drinking coffee was $\geq 2.2 \pm 0.5$ cm. (Wilk et al., 2019)

ETHICAL CLEARANCE

The ethical approval of this research was obtained from the Research Ethics Commission of the Faculty of Medicine, Universitas Trisakti No 6/KER-FK/IX/2021.

RESULTS

Table 1.	e 1. Research Subjects Characteristics (n=22)		
Variable	n (%)	Mean ± SD	
Age		16.55 ± 1.01	
15 years	3 (13.6%)		
16 years	9 (40.9%)		
17 years	5 (22.7%)		
18 years	5 (22.7%)		
Leg muscle strength			
Before drinking coffee		46.68 ± 9.99	
After drinking coffee,			
the dose:			
2 mg CAF/kgBW		51.45 ± 8.92	
3 mg CAF/kgBW		54.05 ± 8.81	

Data table 1 shows the age mean subjects was 16.55 ± 1.01 years with the largest age distribution was 16 years old (40.9%). The averange leg muscle strength before dringking coffee was 46.68 ± 9.99 , whereas after drinking coffee with a dose 2mgCAF/kgBW was 51.45 ± 8.92 and at a dose 3mgCAF/kgBW was 54.05 ± 8.81 .

Table 2. Relationship age and muscle strength

Age	Leg Mu	n-malue	
nge	Increase (n)	Not Increase (n)	p-ouine
15-16 years	11	1	0 802*
17-18 years	9	1	0.892

*: Chi-square test

Table 2 shows that the majority of both subjects in the age group 15-16 years and 17-18 years experienced increased leg muscle strength, but there was no significant relationship between age and leg muscle strength with a p-value 0.892 (p > 0.05).

There et menere et
--

Coffeine dese	Leg Mus	n-value	
Carrenne uose	Before	After	p-ouiue
2mg CAF/kgBW	46.68±9.99	51.45±8.92	0.00*
3mg CAF/kgBW	46.68±9.99	54.05 ± 8.81	0.00*

*: Paired t-test





Figure 1. Leg muscle strength before and after coffee consumption

Based on figure 1, there was a significant increase in muscle strength after giving coffee at dose 2mg CAF/kgBW or dose 3mg CAF/kgBW which was measured 3 times at the 30th, 45th and 60th minutes. This increase of leg muscle strength can be seen continuing at every minute. At dose 2mg CAF/kgBW, it's shown that the average at the 30th minute was 47.95, at the 45th minute was 49.63 and the 60th minute was 51.45. Meanwhile at dose 3mg CAF/kgBW, the average at the 30th minute was 50.95 and at the 60th minute was 54.05.

DISCUSSION

In this study, all subjects were male, because there were no females in the futsal extracurricular at SMA 5 Tambun, Bekasi. The same results were also obtained according to another study. It was said that female participation is low because of the stigma that futsal is an activity that has violent and courageous characteristics that are identical to the male's characteristics. (Akurat & Maksum, 2021)

In our results, there was no relationship between age and muscle strength because all of the subjects were in the teenage age category. One of the risk factors that can influence muscle strength is age. Physicologically, peak mass growth and muscle strength are gained in late adolescence. Meanwhile, muscle mass decreases around 3 to 8% at the age of 30 years and this decreases even higher above the age of 60 years old. (Ng et al., 2019; Volpi et al., 2004)

There was a relationship between coffee consumption and leg muscle strength in both doses of caffeine p=0.00 (p<0.05). This study results showed an increase in the total leg muscle strength average before consuming coffee (46.68±9.66) and after consuming coffee with a dose of 2mg CAF/kgBW and 3mg CAF/kgBW (54.05±8.81) using vertical jump test. Coffee contains purine alkaloids called caffeine (1,3,7-trimethylxanthine) which bind and regulate several protein that has physiological effects and stimulates sympathetic nervous system as an adenosine receptor antagonist.(Ludwig et al., 2014; Wishart et al., 2017)

Caffeine is one of the ergogenic substances that can increase muscle strength. Caffeine also can help improve sport performance, and muscle performance especially to increase aerobic activity endurance as well as increasing the ability to repetitions or repeat movements in muscle training. This is due to the effects of caffeine which can inhibit the muscle Ca²⁺ reuptake by sarcoplasmic reticulum thus reducing excitability threshold value and making muscle contraction last longer. Caffeine increases C-AMP by inhibiting the activity of cyclic nucleotide phosphodiesterase, so the result is increasing free fatty acids (FFA) that can be used as energy and causes glycogen saving effects for endurance performance in the long term. Caffeine has a structure similar to adenosine, acting as a competitive antagonist at the adenosine receptor causing low adenosine concentrations in the body, resulting in increased motor activity and lipolysis inhibition to increases FFA production so it can be used as energy for muscle contraction. In addition, caffeine causes the epinephrine release from the adrenal gland causing lipolysis in muscle and fat tissues as an energy sources. As a results of the increasing FFA, the use of glycogen at the beginning of exercise decreases. (Hayati, 2012; Samoggia & Rezzaghi, 2021; Spriet, 2014)

Caffeine as an adenosine receptor antagonist (A1 and A2) on the skeletal muscle membrane make increases exicitation-contraction coupling through the release of Ca²⁺ from the sarcoplasma reticulum and improves Na⁺-K⁺ATPase pump activity which helps increase muscle performance. Adenosine has the effect to inhibiting lipolysis and stimulating insulin to glucose uptake which causes skeletal muscle contraction. In addition, because it is accompanied by an increase in adenosine consentration during exercise, it can increase glycogenolysis in both working and non working tissue also skeletal adipose tissue during physical activity. Futhermore, caffeine has been proven to improve the force velocity relationship and the fibre's muscle conduction velocity. (Hodgson et al., 2013; Tangen et al., 2020)

In this study, it was found that muscle power performance increased every minutes in both doses, 2mg CAF/kgBW and 3mg CAF/kgBW, which the height occurring at the 60th minute. Recently, some studies said the low caffeine dose administration (3mg/kgBW) also can produce an ergogenic effect, with no changes in heart rate, catecholamines level, lactate, FFA and glycerol during exercise. Every minute the muscle strength increases because of the associated caffeine absorption by the small intestine. Approximately 0.5-2% of the caffeine is excreted by urine and the rest will be absorbed by the body through the small intestine. After oral caffeine consumption, the working time will be reached within 45-60 minutes, where the maximum concentration in plasma will be reached between 30-120 minutes. The increase in muscle strength for each respondent was different due to each individual's physiological digestive system differences which depended on the time of gastric emptying and food consumption. The elimination of caffeine half-life was around 2 to 12 hours, with an average 5 hours. It's been reported that women had shorter half-life compared to men. (Grzegorzewski et al., 2022; Mandal, 2023; Spriet, 2014; Wishart et al., 2017)

Studies conducted by Nandatama R et al. and Tangen DS et al. had the same results with this study, which showed that there was an increase in muscle strength after giving coffee, even though it was said that it could increase athlete's endurance.(Nandatama et al., 2017; Tangen et al., 2020) **CONCLUSION**

There was a significant relationship between coffee consumption and muscle strength in both doses. The increasing muscle strength reached to the maximum at the 60^{th} minutes in both doses.

CONFLICT OF INTEREST

The authors do not have any conflict of interest to declare in this research.

ACKNOWLEDGMENTS

The researchers would like to thank the cooperation of all subjects at SMA 5 Tambun Selatan, Bekasi in this study.

Effect of Coffee Consumption with Muscle Strength in Male Adolescents Futsal Team

ORIGIN	IALITY REPORT				
8 SIMIL	% ARITY INDEX	6% INTERNET SOURCES	6% PUBLICATIONS	1 % STUDENT PA	PERS
PRIMA	RY SOURCES				
1	jddtonli Internet Sour	ne.info			2%
2	" 1st Anr de Gruy Publication	nual Conference ter GmbH, 2020	e of Midwifery')	', Walter	1%
3	www.ka Internet Sour	ryailmiah.trisak	ti.ac.id		1%
4	Sunutch Montri S meat ac PCR-RFI Science Publication	a Suntrarachun Sumontha. "Ider dulteration in me _P of mitochond and Human We	, Lawan Chan ntification of s eat products u lrial DNA", Foo ellness, 2018	home, ea snake ising od	1 %
5	WWW.W Internet Sour	orld-today-news	s.com		1%
6	Andi Bu Shania I About R Muham	nga, Ranti Salsa Maharani. "Desc Reproductive He madiyah Univer	ibila, Rehan Au cription of Kno alth in Youth a sity Jakarta 20	ula, wledge at 021",	1 %

Muhammadiyah International Public Health and Medicine Proceeding, 2021

7	Ahmad Yudi Satriyono, Hariyani Sulistyoningsih, Wuri Ratna Hidayani. "Knowledge, Attitudes and Behaviors of the Adolescent Reproductive Health Triad", Buletin Ilmu Kebidanan dan Keperawatan, 2022 Publication	<1%
8	repository.unair.ac.id	<1%
9	digitalcommons.andrews.edu	<1%
10	fas.org.in Internet Source	<1%
11	perpusnwu.web.id	<1%
12	Bergita Dumar, Heni Dwi Windarwati, Laily Yuliatun. "Factors Affecting School-Based Mental Health in Adolescents: Scoping Review", Jurnal Aisyah : Jurnal Ilmu Kesehatan, 2023 Publication	<1 %





Exclude quotes Off Exclude bibliography Off Exclude matches Off

AUTHOR'S AGREEMENT

Dear Editor of the **Madani Multidisciplinary Journal (MUDIMA)** Website: <u>https://journal.formosapublisher.org/index.php/mudima</u>

This is to certify that the manuscript,

 Title
 Effect of Coffee Consumption with Muscle Strength in Male Adolescents Futsal Team

Author : Ayu Astianing Tiyas, Eveline Margo*

on behalf of the author(s) we declare:

- 1. The manuscript is original, and its publication does not violate any copyright issues.
- 2. The manuscript has never been published before, in whole or in part, in any scientific journal or publishing company.
- 3. The manuscript has never been/is not included in the journal publishing process.
- 4. The registered author is committed to showing integrity in making this paper.
- 5. All authors agree to publish the final version of the article after the editing process is carried out by the editor.
- 6. All authors agree to publish the manuscript in the **Madani Multidisciplinary Journal** (**MUDIMA**)

Jakarta, 27 September 2023 Corresponding author, Signature and full name and title

dr.Eveline Margo, M. Biomed

Jurnal Multidisiplin Madani (MUDIMA) has been widely indexed by:





Jurnal Multidisiplin Madani (MUDIMA) has been widely indexed by:

Google Scholar, 2. Copernicus International (ICV 2022:69,73), 4. Open Aire 5. GARUDA, 6. ScienceGate
 Directory of Research Journals Indexing, 8.SHERPA/Romee, 9. RESEARCH BIBLE, 10. DIMENSIONS, 11.
 WorldCat, 12. SciLit, 13. CiteFactor, 14. Semantic Scholar, 15. Eurasian Scientific Journal Index, 16. IP
 Indexing 17. Root Indexing, 18. IUPUI Libraries, 19. Scite, 20. Neliti, 21. Advanced Sciences Index (ASI), 22.
 Harvard Library, 23. IP Indexing 24. Indonesia OneSearch 25. Base (bielefeld academic search engine) 26.
 Fatcat, 27. Exaly, 28. Stanford Libraries, 29. Crossref, 30. Russel Sage College Library



Other Libraries:



Complete indexing and libraries information can be checked: https://journal.formosapublisher.org/index.php/mudima/indexing

Formosa Publisher is supported and affiliated with the <u>International Association of Contemporary Researchers</u> (<u>IACR</u>).

Indexed By:				
INDEX COPERNICUS		SHERPA/Romeo	CROSSREE/ORG THE CITATION LINKING BACKBONE	
Google	R GARUDA	ॐ WorldCat [∗]	© Scilit	
Directory of Research Journal Indexing	Dimensions			
BASE Bielefeld Academic Search Engine	scite_	RESEARCHBIB ACADEMIC RESOURCE INDEX	ESJI www.ESJIndex.org	
ROTINDEXING OURNAL ABSTRACTING AND INDEXING SERVICE	ADVANCED SCIENCE INDEX	IPIndexing Indexing Portal	SEMANTIC SCHOLAR	
	(x)	a CP		

Home / Archives / Vol. 3 No. 9 (2023): September, 2023 / Articles

Effect of Coffee Consumption with Muscle Strength in Male Adolescents Futsal Team

Ayu Astianing Tiyas Universitas Trisakti, Jakarta

Eveline Margo Universitas Trisakti, Jakarta

DOI: https://doi.org/10.55927/mudima.v3i9.3384

Keywords: Caffeine, Muscle Strength, Vertical Jump Test, Male Adolescents



ournal.formosapublisher.org/index.php/mudima

Vol. 3 No. 9 (2023): September, 2023



DOI: https://doi.org/10.55927/mudima.v3i9

Published: 2023-09-29

Articles

We Heal as One, We Rise as One: the Rise of Community Pantries in the Midst of COVID-19 Pandemic Northern Philippines Ederlyn Bautista, University of Saint Louis, Tuguegarao City, Philippines Krisha Mae Narag, University of Saint Louis, Tuguegarao City, Philippines Darin Jan Tindowen, University of Saint Louis, Tuguegarao City, Philippines Cirilo Parallag Jr, University of Saint Louis, Tuguegarao City, Philippines	in 1933-1947
Physicochemical and Fat-Soluble Vitamins Constituents of Fresh and Reused Vegetable Oil from Ab Metropolis, Ebonyi State	akaliki
Nwabunma C. Asouzu, Department of Biochemistry, Faculty of Science, Ebonyi State University, Abakaliki, Nigeria Okide Afoma Ifunanya, Department of Nutrition and Dietetics, Alex Ekueme Federal University Teaching Hospital, Nigeria Patrick M. Aja, Department of Biochemistry, Kampala International University, Western-Campus Ishaka, Uganda Okon Michael Ben, Department of Biochemistry, Kampala International University, Western-Campus Ishaka, Uganda Nonso C. Asouzu, Department of Physiotherapy, Alex Ekueme Federal University Teaching Hospital, Abakaliki, Nigeria Aja Lucy, Department of Science Education, Faculty of Education, Ebonyi State University, Nigeria Ejike Daniel Eze, Department of Physiology, School of Medicine, Kabale University, Uganda Afodun Adam Moyosore, Department of Anatomy and Cell Biology, Faculty of Health Sciences, Busitema University, Uganda Awuchi G. Chinaza, School of Natural and Applied Sciences, Kampala International University, Kampala, Uganda	1986-1998
Design Thinking Testing of AR/VR Application for Bali's Lontar Prasi Preservation Christina Purnama Yanti, Institut Bisnis dan Teknologi Indonesia (INSTIKI), Indonesia I Gede Iwan Sudipa, Institut Bisnis dan Teknologi Indonesia (INSTIKI), Indonesia Putu Wirayudi Aditama, Institut Bisnis dan Teknologi Indonesia (INSTIKI), Indonesia	1956-1963
Real Time Face Recognition for Mobile Application Based on Mobilenetv2	1055 1064

I Made Subrata Sandhiyasa, Institut Bisnis dan Teknologi Indonesia (INSTIKI), Indonesia Devi Valentino Waas, Institut Bisnis dan Teknologi Indonesia (INSTIKI), Indonesia

🖾 pdf

inhancing Customer Satisfaction	: Exploring Hotel X'S Facilities and	Service Quality in Bandung City
---------------------------------	--------------------------------------	---------------------------------

Diyan Putranto, Hotel and Tourism Department, Saint Marry Collage, Jakarta, Indonesia Adi Sopyan, Hotel and Tourism Department, Saint Marry Collage, Jakarta, Indonesia Anindita Gayatri, Hotel and Tourism Department, Saint Marry Collage, Jakarta, Indonesia Barliani Dewi Setyawati, Hotel and Tourism Department, Saint Marry Collage, Jakarta, Indonesia Albertus Banundoyo, Hotel and Tourism Department, Saint Marry Collage, Jakarta, Indonesia

Analysis of Financial Distress, Profitability, Liquidity, Leverage on Share Prices of Manufacturing Companies Listed on the IDX

Dahlia Pinem, Universitas Pembangunan Nasional Veteran Jakarta, Indonesia Marlina Tanjung, Universitas Pembangunan Nasional Veteran Jakarta, Indonesia Anggi Angga Resti, Universitas Pembangunan Nasional Veteran Jakarta, Indonesia 1902-1913

1948-1955

B	ndf
<u>~</u>	pui

Benefits of Wayang Kamasan Klungkung Bali in Improving Dignified Cultural Character and Sustainability of **International Tourism** Ni Ketut Wiriani Merta, Universitas Udayana, Denpasar Bali, Indonesia

I Putu Sudana, Universitas Udayana, Denpasar Bali, Indonesia Yohanes Kristianto, Universitas Udayana, Denpasar Bali, Indonesia 1898-1901

🖾 pdf

Geometric Modeling of River Estuary to Support Normalization Program of Veteran River, Banjarmasin City Ferry Sobatnu, Politeknik Negeri Banjarmasin, Indonesia 1999-2010 Nurul Inayah, Politeknik Negeri Banjarmasin, Indonesia

🖾 pdf

🕒 pdf

The Balinese Lontar Manuscript Metadata Model: An Ontology-Based Approach

Ida Bagus Gede Sarasvananda, Institut Bisnis dan Teknologi Indonesia (INSTIKI), Indonesia Putu Gede Surya Cipta Nugraha, Institut Bisnis dan Teknologi Indonesia (INSTIKI), Indonesia Ida Bagus Ary Indra Iswara, Institut Bisnis dan Teknologi Indonesia (INSTIKI), Indonesia

1964-1971

Design of Traffic Lights at the Unsignalized Intersection of Narogong Highway-Pangkalan 5 Road Km	. 13.5
Bekasi City, West Java	

Aulia Choiri, Universitas Global Jakarta, Indonesia
 Ribut Nawang Sari, Universitas Global Jakarta, Indonesia
 Dinar Ali, Universitas Global Jakarta, Indonesia
 Lintang D.A, Universitas Global Jakarta, Indonesia
 Yunan Hanun, Universitas Global Jakarta, Indonesia

A	pdf

Student Perceptions Regarding E-Learning During the Covid-19 Pandemic on the Level of Technostress

2039-2049

2011-2025

Hania Pemilia , Siliwangi University, Indonesia
Yoni Hermawan , Siliwangi University, Indonesia
Bakti Widyaningrum, Siliwangi University, Indonesia

🔎 pdf

Predicting Investment Feasibility in Perumda Establishment Using Payback Period, Net Present Value, and Internal Rate Return

Citra Sarasmitha, University of Merdeka Malang, Indonesia Karina Setia Utami, University of Merdeka Malang, Indonesia

ዾ pdf

Effect of Coffee Consumption with Muscle Strength in Male Adolescents Futsal Team

Ayu Astianing Tiyas, Universitas Trisakti, Jakarta, Indonesia Eveline Margo, Universitas Trisakti, Jakarta, Indonesia 1978-1985

1885-1897

🕒 pdf

Representation of Parental Communication in Nanti Kita Cerita Tentang Hari ini (2020)

Leviana Susanto , Universitas Multimedia Nusantara, Indonesia Tangguh Okta Wibowo, Universitas Multimedia Nusantara, Indonesia 1844-1854

🕒 pdf

🔄 pdf

 How Financial Performance, CSR Disclosure, and GCG Mechanism in Indonesia and Malaysia Banks?

 Brigitta Eva Sinta Amartya, Faculty of Economic and Business, Telkom University, Indonesia

 1820-1837

 Wahdan Arum Inawati, Faculty of Economic and Business, Telkom University, Indonesia

The Relationship Between Parental Support on Children Oral Health and Children Dental Caries	s in
Agroindustrial Area of Jember	

Kiswaluyo, Faculty of Dentistry, Universitas Jember , Indonesia
Roedy Budirahardjo, Faculty of Dentistry, Universitas Jember , Indonesia
Sulistiyani, Faculty of Dentistry, Universitas Jember , Indonesia
Elyda Akhya Afida Misrohmasari, Faculty of Dentistry, Universitas Jember , Indonesia
Ni Luh Putu Maharani Eka Putri, Faculty of Dentistry, Universitas Jember , Indonesia

Implementation of an Accounting Information Systems for Fast Secure Credit

2032-2038

1972-1977

Yuliana, Universitas Hasanuddin, Indonesia Haliah, Universitas Hasanuddin, Indonesia Nirwana, Universitas Hasanuddin, Indonesia

🖾 pdf

Hardiness as the Moderator in the Effect of Mental Workload on Job Stress in Frontline Bank Emp	loyees in the
Pematang Siantar Area	
Rizty Desta Mahestri, Universitas Sumatera Utara, Indonesia	1865-1873

Gustiarti Leila, Universitas Sumatra Utara, Indonesia **Etti Rahmawati,** Universitas Sumatra Utara, Indonesia

🖾 pdf

Saving Behavior of Clove Farmer Households in the Form of Commodities (Case Study in Negeri Ullath Maluku Indonesia)

Jeanee Nikijuluw, Universitas Pattimura, Indonesia

🖾 pdf

Hypnotherapy Intervention on Pain Scale Level of Arteriovenous Fistula Cannulation in Haemodialysis Patients with Chronic Renal Failure

Ahmad Syaripudin, Institut Teknologi dan Kesehatan Mahardika, Indonesia

2026-2031

1914-1920

👌 pdf

The Perspective of Tax Center Digitalization in Strengthening Taxation Socialization: A Case Study at State Polytechnic of Madiun

1874-1884

Tri Septianto, Politeknik Negeri Madiun, Indonesia Koerniawan Dwi Wibawa, Politeknik Negeri Madiun, Indonesia Ikhwan Baidlowi Sumafta, Politeknik Negeri Madiun, Indonesia Imam Mudofir, Politeknik Negeri Madiun, Indonesia Meiliyana Susanti, Politeknik Negeri Madiun, Indonesia



 The Relationship Between Natural Medicine Use and Recovery Time in Early Adult COVID-19 Patients

 Natasya Cindy Edward , Universitas Trisakti, Indonesia
 1921-1932

 Meiyanti, Universitas Trisakti, Indonesia
 1921-1932

🖾 pdf

The Effectiveness of Zeolite-Activated Charcoal in Reducing Iron and Manganese Levels in Well Water of Milala Housing in Pancur Batu District, North Sumatra

Haesti Sembiring, Politeknik Kementerian Kesehatan Medan, Indonesia

1838-1843

ዾ pdf