

## GLOBAL MEDICAL AND HEALTH COMMUNICATION

**Global Medical and Health Communication** is a journal that publishes medical and health articles published since 2013. Articles are original research that needs to be disseminated and written in **English**.

In not so long time, **Global Medical and Health Communication** that managed by the **Faculty of Medicine, Universitas Islam Bandung (Unisba)** and published by **UPT Publikasi Ilmiah Unisba** already accredited by the National Journal Accreditation (Arjuna) managed by the Directorate General of Higher Education, Research, and Technology, Ministry of Education, Culture, Research, and Technology of the Republic of Indonesia with 2<sup>nd</sup> Grade (**Sinta 2**) starting from Volume 7 Number 2 of 2019 to Volume 12 Number 2 of 2024. It's also indexed in the **Directory of Open Access Journals (DOAJ)** on 9<sup>th</sup> May 2017 and **Crossref** on 2<sup>nd</sup> January 2018. With DOAJ and Crossref indexing, this journal can reach international audiences.

This achievement received a positive response from researchers, lecturers, and health observers indicated by the articles submitted, three times the number of articles received initially. The quality of the articles also shows improvement in terms of methodology and writing that is useful for audiences. Research results were best to disseminate as early as possible to good use. Publication of the **Global Medical and Health Communication** every **6 (six) months** in a year will be published every **4 (four) months** starting in 2017.

The news from science and health about challenges and opportunities in Indonesia is still very rare and **Global Medical and Health Communication** hopes to become the best means to support researchers, lecturers, and health practitioners to become the voice of Indonesia, especially in health.

pISSN 2301-9123 | eISSN 2460-5441



VOL 12, NO 2 (2024)

**AUGUST 2024 (ISSUE IN PROGRESS)**

**TABLE OF CONTENTS**

### Articles

Stakeholder Insights on Malaria Elimination Strategies in Pangandaran, West Java: a Qualitative Analysis

PDF

Lia Faridah, Nisa Fauziah, Fayyaza Faiz Adams, Hasna Mufida, Muhamad Lazuardi Akbar, Pricillia Laurenza Salsabila, Wilbert Bernardi Rufinus, Abimanyu Athallah Virajati, Anggisti Nurdinda Chaerany Putri Angga, Khansa Mahira, Ridha Beta Zahra

Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages

PDF

Elsi Septira Wibowo, Eveline Margo

## FOCUS AND SCOPE

**Global Medical and Health Communication** is a journal that publishes research articles on medical and health published every 4 (four) months (April, August, and December). Articles are original research, meta analysis, and systemic review that needs to be disseminated and written in English.

Subjects suitable for publication include but are not limited to the following fields of anesthesiology and intensive care, biochemistry, biomolecular, cardiovascular, child health, dentistry, dermatology and venerology, endocrinology, environmental health, epidemiology, geriatric, health communication, health promotion, hematology, histology, histopathology, immunology, internal medicine, nursing sciences, midwifery, nutrition, nutrition and metabolism, obstetrics and gynecology, occupational health, oncology, ophthalmology, oral biology, orthopedics and traumatology, otorhinolaryngology, pharmacology, pharmacy, preventive medicine, public health, pulmonology, radiology, and reproductive health.

## SECTION POLICIES

### Articles

☒ Open Submissions ☒ Indexed ☒ Peer Reviewed

### Editorial

☐ Open Submissions ☒ Indexed ☐ Peer Reviewed

### Guest Articles

☐ Open Submissions ☒ Indexed ☐ Peer Reviewed

## EDITORIAL TEAM

### Editor in Chief

Santun Bhekti Rahimah, Department of Pharmacology, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

### Managing Editor

Winni Maharani, Department of Microbiology, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

### External Editorial Board

Badrul Hisham Yahaya, Advanced Medical and Dental Institute, Universiti Sains Malaysia, Kepala Batas, Malaysia  
Jerico Franciscus Pardosi, School of Public Health and Social Work, Queensland University of Technology, Kelvin Grove, Queensland, Australia  
Roy Rillera Marzo, Asia Metropolitan University, Johor, Malaysia

### Internal Editorial Board

Ike Rahmawaty Alie, Department of Physiology, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia  
Indri Budiarti, Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia  
Lisa Adhia Garina, Department of Child Health, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia  
Mirasari Putri, Department of Biochemistry, Nutrition, and Biomolecular, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia  
Yuktiana Kharisma, Departement of Pathology Anatomy, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

### Proofreader

Herry Garna, Department of Child Health, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

### Copyeditor

Yudi Feriandi, Department of Medical Education, Bioethics, Humaniora and Islamic Insert, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

### Layout Editor

Agus Chalid, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

### Editorial Assistants

Agus Chalid, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia  
Ihsan Mulyadi Kurniawan, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

pISSN 2301-9123 | eISSN 2460-5441

### Reviewer

Angung Firmansyah Sumantri, Department of Internal Medicine, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

Arief Budi Yulianti, Department of Medical Biology, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

Arif Susanto, Occupational Health, Safety & Environmental (HSE), PT Freeport Indonesia, Tembagapura, Indonesia

Budi Setiabudiawan, Department of Child Health, Faculty of Medicine, Universitas Padjadjaran/Dr. Hasan Sadikin General Hospital, Bandung, Indonesia

Cordia Ming Yeuk Chu, Centre for Environment and Population Health, Griffith University, Brisbane, Australia

Dicky Santosa, Department of Child Health, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

Djatnika Setiabudi, Department of Child Health, Faculty of Medicine, Universitas Padjadjaran, Bandung, Indonesia

Ermi Ndoen, UNICEF Indonesia, Jakarta, Indonesia

Ferry Efendi, Faculty of Nursing, Universitas Airlangga, Surabaya, Indonesia

Hendro Sudjono Yuwono, Department of Surgery, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

[Heni Muflihah](#), Department of Pharmacology, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

[Herri S Sastramihardja](#), Department of Pharmacology, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

[Hidayat Wijayanegara](#), Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

[Ieva B Akbar](#), Department of Physiology, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

[Johanes Cornelius Mose](#), Department of Obstetric and Gynecology, Faculty of Medicine, Universitas Padjadjaran, Bandung, Indonesia

[Lelly Yuniarti](#), Department of Biochemistry, Nutrition, and Biomolecular, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

[Leri Septiani](#), STIKes Dharma Husada/RSIA Grha Bunda, Bandung, Indonesia

[Lia Faridah](#), Department of Biomedical Science, Faculty of Medicine, Universitas Padjadjaran, Bandung, Indonesia

[M Ahmad Djojosugito](#), Department of Orthopedics and Traumatology, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

[Mas Rizky Anggun Adipurna Syamsunarno](#), Department of Biochemistry and Molecular Biology, Faculty of Medicine, Universitas Padjadjaran, Jatinangor, Indonesia

[Maya Tejasari](#), Department of Histology, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

[Mia Kusmiati](#), Department of Medical Education, Bioethics and Humanity, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

[Nuzirwan Acang](#), Department of Internal Medicine, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

[Riyadi Adrizain](#), Department of Child Health, Faculty of Medicine, Universitas Padjadjaran, Bandung, Indonesia

[Rizky Suganda Prawiradilaga](#), Department of Biochemistry, Nutrition and Biomolecular, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

[Titik Respati](#), Department of Public Health, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

[Umar Fahmi Achmadi](#), Department of Environmental Health, Faculty of Public Health, Universitas Indonesia, Depok, Indonesia

[Wahyu Widowati](#), Department of Biomolecular, Faculty of Medicine, Universitas Kristen Maranatha, Bandung, Indonesia

[Wawang Setiawan Sukarya](#), Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Muhammadiyah Prof. Dr. HAMKA, Jakarta Selatan, Indonesia

[Wida Purbaningsih](#), Department of Histology, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

[Yani Triyani](#), Department of Clinical Pathology, Faculty of Medicine, Universitas Islam Bandung, Bandung, Indonesia

[Yulia Sofiatin](#), Division of Epidemiology and Biostatistics, Department of Public Health, Universitas Padjadjaran, Bandung, Indonesia

Pada Min, 14 Jul 2024 pukul 17.57 eveline-margo fk <[eveline-margo@trisakti.ac.id](mailto:eveline-margo@trisakti.ac.id)> menulis:

Dear  
GMHC reviewer,  
Thank you for your information.  
I already makes my manuscripts revision regarding from reviewer's inputs and also already input in GMHC link journal.  
Please kindly to review again.  
Thank you very much.

Best regards,  
Eveline Margo

On Wed, Jul 10, 2024 at 1:37 PM gmhc. unisba <[gmhc.unisba@gmail.com](mailto:gmhc.unisba@gmail.com)> wrote:

Dear  
**Mr./Mrs./Ms. Eveline Margo**

We have decided on your submission to Global Medical & Health Communication (GMHC) entitled:  
**Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages.**

We have decided that your article needs some revision.  
The revised file has to be submitted on **17 July 2024**.

## [GMHC] Revisions Required for Manuscript No.#12299 (Third)

External

Inbox x



**gmhc. unisba**

to me ▾

Fri, Jul 19, 2:01PM

Dear  
**Mr./Mrs./Ms. Eveline Margo**

We have decided on your submission to Global Medical & Health Communication (GMHC) entitled:  
Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages.

We decided that your article needs revision, especially regarding the **English**. Please fix it.  
We offer translation services at an additional cost.

Regards,  
Editorial Board of Global Medical & Health Communication (GMHC)  
Faculty of Medicine, Universitas Islam Bandung  
Mobile: 085220013654  
E-mail: [gmhc.unisba@gmail.com](mailto:gmhc.unisba@gmail.com)  
Website: <https://ejournal.unisba.ac.id/index.php/gmhc>

## Review manuscript 12299

External

Inbox x



**eveline-margo fk** <[eveline-margo@trisakti.ac.id](mailto:eveline-margo@trisakti.ac.id)>  
to unisba ▾

Mon, Jul 22, 6:56 PM



Dear  
Editorial Board of Global Medical & Health Communication,  
  
Thank you for the suggestions for improving my manuscript.

## [GMHC] Accept Submission for Manuscript No.#12299

External

Inbox x



gmhc.unisba

to me ▾

Jul 25, 2024, 2:28 PM



Dear

**Mrs. Eveline Margo**

Your article submission to Global Medical and Health Communication, "[Smoking Habit and Coffee Consumption related to Gastritis Incidence Rate at Productive Ages](#)."

We decide to: **ACCEPT SUBMISSION**.

Your article will be publishing on GMHC Vol. 12 No. 2, 2024.

In addition, please immediately send the following requirements to [gmhc.unisba@gmail.com](mailto:gmhc.unisba@gmail.com) (latest **28 July 2024**):

1. Receipt of Author Article Publication Fee (**IDR 1,500,000.00**)  
to  
Bank Syariah Indonesia (BSI)  
Account number: 5003001120  
Account holder's name: Fakultas Kedokteran Unisba

## [GMHC] Proofreading Request

External

Inbox x



gmhc.unisba

to me ▾

Fri, Aug 2, 2:46 PM (12 days ago)



Dear

**Mrs. Eveline Margo**

Your submission "[Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages](#)" to Global Medical and Health Communication now needs to be proofread by following these steps.

1. Click on the Submission URL below.
2. Log into the journal and view **PROOFING INSTRUCTIONS**
3. Click on **VIEW PROOF** in Layout and proof the galley in the one or more formats used.
4. Enter corrections (typographical and format) in Proofreading Corrections.
5. Save and email corrections to Layout Editor and Proofreader.
6. Send the **COMPLETE** email to the editor.

Submission URL: <https://ejournal.unisba.ac.id/index.php/gmhc/author/submissionEditing/12299>

Username: evemargo



eveline-margo fk <eveline-margo@trisakti.ac.id>

to unisba ▾

Tue, Aug 13, 3:09 PM (21 hours ago)



Kepada Yth,  
Editor GMHC

Dengan hormat,  
mohon maaf sepertinya terjadi pemasukkan data terakhir.

 14 Aug 2024 01:27:30 An issue has been published.	<a href="#">Go To URL</a> <a href="#">Delete</a>
 14 Aug 2024 01:25:54 An issue has been published.	<a href="#">Go To URL</a> <a href="#">Delete</a>
 14 Aug 2024 01:23:02 A galley has been modified for "Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages".	<a href="#">Go To URL</a> <a href="#">Delete</a>
 14 Aug 2024 00:43:05 A galley has been modified for "Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages".	<a href="#">Go To URL</a> <a href="#">Delete</a>
 02 Aug 2024 15:33:47 "Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages's" metadata has been modified.	<a href="#">Go To URL</a> <a href="#">Delete</a>
 31 Jul 2024 13:04:08 "Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages's" supplementary file(s) has been modified.	<a href="#">Go To URL</a> <a href="#">Delete</a>
 15 Jul 2024 15:35:02 "Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages's" metadata has been modified.	<a href="#">Go To URL</a> <a href="#">Delete</a>
 10 Jul 2024 14:24:45 "Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages's" supplementary file(s) has been modified.	<a href="#">Go To URL</a> <a href="#">Delete</a>
 10 Jul 2024 14:22:23 "Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages's" supplementary file(s) has been modified.	<a href="#">Go To URL</a> <a href="#">Delete</a>
 10 Jul 2024 14:22:00 "Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages's" supplementary file(s) has been modified.	<a href="#">Go To URL</a> <a href="#">Delete</a>
 10 Jul 2024 14:21:56 "Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages's" supplementary file(s) has been modified.	<a href="#">Go To URL</a> <a href="#">Delete</a>
 10 Jul 2024 14:20:27 "Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages's" supplementary file(s) has been modified.	<a href="#">Go To URL</a> <a href="#">Delete</a>
 01 May 2024 00:41:51 An issue has been published.	<a href="#">Go To URL</a> <a href="#">Delete</a>

## RESEARCH ARTICLE

## Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages

Elsi Septira Wibowo,<sup>1</sup> Eveline Margo<sup>2</sup>

<sup>1</sup>Faculty of Medicine, Universitas Trisakti, East Jakarta, Indonesia,

<sup>2</sup>Department of Physiology, Faculty of Medicine, Universitas Trisakti, East Jakarta, Indonesia

### Abstract

Gastritis is a disease caused by an inflammation of the mucous and submucosal lining of the stomach. Several studies show that there is an influence of smoking habit and coffee consumption on gastritis incidence rate. The study was to determine the correlation between smoking habit and coffee consumption toward gastritis at productive ages. The research is an analytical observational study with a cross-sectional design conducted in Makale Health Center, South Sulawesi, in February 2020. Required data was gathered by conducting guided interviews using a questionnaire that includes age range, sex, and education level. Brinkman Index was used to measure smoking habits, coffee consumption, and gastritis questionnaire. Data was analyzed using a chi-square test with a significance level of  $p=0.05$ . From a total of 115 respondents, there are 69.6% included in the age range of 26–35 years old, 58.3% have mild smoking habits, 56.5% have severe coffee consumption, and 85.2% are suffering from gastritis. There is a relation between smoking habit ( $p=0.029$ ) and coffee consumption ( $p=0.003$ ) with gastritis ( $p<0.05$ ). The conclusion is that there is a relationship between smoking habits, coffee consumption, and gastritis incidence.

**Keywords:** Coffee, gastritis, productive age, smoking

### Introduction

Gastritis is a common disease that often occurs in everyone and is a health problem with a high prevalence. Lifestyle is an essential factor in the occurrence of gastritis.<sup>1,2</sup> Gastritis is an inflammation of the mucous and submucosal lining of the stomach, which can be caused by several factors, such as smoking, coffee drinking, and drug consumption, such as aspirin or NSAID.<sup>3</sup>

The incidence of world gastritis is around 1.8–2.1 million of the population every year. Based on research conducted by the World Health Organization, gastritis incidents occurred highly in the USA (47%), India (43%), Canada (35%), and China (31%). Based on data from the Ministry of Health of the Republic of Indonesia (2019), 40.8% of gastritis incidents occur in Indonesia.<sup>1,4</sup> Gastritis patient prevalence is often found in the age range of 20 to 44 years old. Research conducted by the University of Medicine and Pharmacy Tirgu-Mures mentioned that gastritis prevalence in people with coffee consumption and smoking habits often occurred in those who

are 30 to 49 years old.<sup>5</sup>

Smoking is one of many modern lifestyles that are not only found in adults but also in children and young adults. According to the 5<sup>th</sup> Indonesian Family Life Survey (IFLS-5) data, the percentage of smoking in Indonesia was 58%, with 95% dominated by males with an average age of 19 years old. Most (48%) choose kretek filtered as the preferred cigarette.<sup>6</sup>

Smoking habits can cause gastric disturbance because nicotine in cigarettes will pucker and disrupt blood vessels in the stomach, which will further cause irritation that triggers excessive increase of gastric acid production.<sup>7</sup>

Another factor that influences gastritis occurrence is consuming coffee/caffeine. Long-term and high-frequency caffeine consumption (>3 glasses a day) will accelerate gastric acid production that would irritate the mucosa. When mucosa in the stomach is irritated, hydrochloride acid (HCl) diffusion towards the mucosa will disrupt mucosa and eliminate its ability to protect the stomach from HCl and pepsin auto digestion.<sup>3,5,8,9</sup> Research conducted by Ilham

Copyright ©2024 by authors. This is an open access article under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License (<https://creativecommons.org/licenses/by-nc-sa/4.0>).

Received: 12 July 2023; Revised: 24 July 2024; Accepted: 25 July 2024; Published: 14 August 2024

**Correspondence:** Eveline Margo, dr., M.Biomed. Department of Physiology, Faculty of Medicine, Universitas Trisakti, Jln. Kyai Tapa No.1, Grogol, West Jakarta 11440, Special Capital Region of Jakarta, Indonesia. E-mail: [eveline-margo@trisakti.ac.id](mailto:eveline-margo@trisakti.ac.id)

et al.<sup>10</sup> stated that the caffeine compound in coffee can accelerate gastric acid production stimulation and cause gastritis. However, a different result was published by Novitasary et al.,<sup>11</sup> which said that there is no relationship between coffee consumption and gastritis, which is highly dependent on the type of coffee, where in this research, the subjects consume more instant coffee with a lower caffeine compound than ground coffee beans. Research by Cheng et al.<sup>12</sup> said that there is a relationship between smoking habit and gastritis, but contrary research by Novitasary et al.,<sup>11</sup> Hairuddin et al.,<sup>13</sup> and Khandelwal et al.,<sup>14</sup> which said that there is no relationship between smoking habit and gastritis occurrence because there are factors with stronger influence on gastritis such as consumption habit and stress.

Based on the description above, the authors are interested in researching the relationship between smoking habits and coffee consumption and gastritis incidence rates at productive ages, which can further identify the contributing factors to preventive and curative sources of gastritis.

## Methods

This research is a type of observational analytical study using a cross-sectional design. This study searched for a relationship between smoking habits and coffee consumption with gastritis incidence rate in Makale Health Center, South Sulawesi, conducted in February 2020. Subjects were men and women aged 26–45 years at the study site. Subjects in this study were taken through simple random sampling on subjects by the inclusion criteria, namely men or women aged 26–45 years, have smoking habits and coffee consumption, are not taking NSAIDs and aspirin, and are willing to sign informed consent. Determination of sample size is done by using an infinite-finite population formula with a significance level of 95% from 1.96, and with gastritis prevalence in Indonesia of 40.8%<sup>1</sup> where the measurement accuracy is 0.05 and 115 subjects are willing to participate in this study.

Data collection was collected through questionnaire filling that gathers information regarding identity (name, age, sex, and education level), Brinkman index questions to measure the number of cigarettes consumed in a day multiplied by length of smoking habit in a year with valuation category of light between 0–199,

medium between 200–599, and heavy >600.<sup>15</sup> The questionnaire also measures the length and frequency of coffee consumption practiced by the subjects with categories of light with a frequency of  $\leq 3$  glasses/day and heavy with a frequency of  $> 3$  glasses/day.<sup>16</sup> A gastritis questionnaire was also included to measure the perceived gastritis symptom complaints with a valuation of  $\geq 12$  and  $< 12$  for non-gastritis complaints.<sup>2</sup>

The Research Ethics Committee of the Faculty of Medicine, Universitas Trisakti, Jakarta, has already approved this research, with approval number 11/KER-FK/1/2019.

## Results

Univariate analysis was used to determine the frequency distribution of respondents' characteristics in terms of age, sex, education level, smoking habits, coffee consumption, and gastritis events of the research subjects.

Table 1 suggests that most subjects in the Makale Health Center were aged between 26 and 35 years (69.6%), with a slightly higher proportion of males (56.5%) than females. Eighty subjects (69.6%) had a high level of education, 67 subjects (58.3%) could be considered to have a mild smoking habit, and 65 subjects (56.5%) are known to consume coffee to a significant extent.

**Table 1 Characteristic Data**

Variables	n=115 (%)
Age (years)	
26–35	80 (69.6)
36–45	35 (30.4)
Gender	
Male	65 (56.5)
Female	50 (43.5)
Level of education (year)	
Low ( $< 9$ )	35 (30.4)
High ( $\geq 9$ )	80 (69.6)
Smoking habit	
Mild ( $< 200$ )	67 (58.3)
Moderate (200–599)	45 (39.1)
Severe ( $> 600$ )	3 (2.6)
Coffee consumption	
Mild ( $\leq 3$ )	50 (43.5)
Severe ( $> 3$ )	65 (56.5)
Gastritis occurrences	
Gastritis	98 (85.2)
Non-gastritis	17 (14.8)

In the assessment of the incidence of gastritis, it was found that 98 subjects (85.2%) had gastritis.

Bivariate analysis with a chi-square test was used to determine the relation between socio-demographic factors (age, sex, and level of education), smoking habits, and coffee consumption on the incidence of gastritis at the age of 26–45.

Table 2 shows that gastritis occurrence is found more in people in the age group of 26–35 years old with 68 subjects (59.1%), and from chi-square statistical test result, it is revealed that there is no significant relation between age group and gastritis incidence rate in subjects of 26–45 years of age ( $p=0.921$ ). As many as 55 subjects (47.8%) are male with gastritis, which is slightly higher than female subjects. This shows that there is no significant difference between the two sexes. Based on the chi-square statistical test result, a value of  $p=0.836$  was obtained. Based on that, we can conclude no significant relation exists between sex and gastritis incidence rate in subjects at productive ages. In groups with higher education levels, there are 66 subjects (57.4%) that suffer from gastritis, which is more than the one with lower education. In this group, the chi-square statistical test result shows a value of  $p=0.214$ , which means no relation exists between education level and gastritis incidence rate in

subjects at productive age.

On a bivariate result of the smoker group, authors combine moderate and severe smokers because there are only three subjects who are heavy smokers. The result obtained shows that the light smokers group suffers more gastritis incidence with 53 subjects (46.1%). On the chi-square test result, a value of  $p=0.029$  was obtained, which means a relationship exists between smoking habits and the incidence rate of gastritis in subjects at a productive age. As for the coffee consumption variable, gastritis incidence was found more in heavy coffee consumers, with 61 subjects (53%). The chi-square test result shows a value of  $p=0.003$ , which means that there is a relationship between coffee consumption and gastritis incidence. The results indicated that the subjects consuming more coffee can increase the incidence of gastritis at a productive age.

Discussion

In this research, demographic characteristics (age, sex, and education level) are unrelated to the gastritis incidence rate in subjects aged 26–45. Gastritis tends to happen at a younger age due to uncontrolled eating patterns caused by daily activity.<sup>1,17</sup> The results by Feyiza and Woldeamanuel<sup>3</sup> showed younger aged 18–28

Table 2 Bivariate Relation between Demographic Characteristics, Smoking Habits and Coffee Consumption with Gastritis Incidence Rate at Productive Ages

Variables	Gastritis Occurrence				p <sup>c</sup>
	Gastritis		Non-gastritis		
	n=98	%	n=17	%	
Age (years)					
26–35	68	59.1	12	10.4	0.921
36–45	30	26.1	5	4.3	
Gender					
Male	55	47.8	10	8.7	0.836
Female	43	37.4	7	6.1	
Level of education (year)					
Low (<9)	32	27.8	3	2.6	0.214
High (≥9)	66	57.4	14	12.2	
Smoking habit					
Mild	53	46.1	14	12.2	0.029*
Moderate–severe	45	39.1	3	2.6	
Coffee consumption					
Mild	37	32.2	13	11.3	0.003*
Severe	61	53	4	3.5	

Note: <sup>e</sup>chi-square statistical test, \*significance  $p<0.05$

years (57.5%) tend to suffer gastritis because they don't know about good health behaviors, but according to Khandelwal V et al.,<sup>14</sup> those aged 14–25 years and 36–45 years has more tend to be gastritis. However, gastritis can also occur in older subjects because of several factors such as age increase, degenerative process, *Helicobacter pylori* infection, and NSAID drug usage.<sup>18</sup> The research by Putri et al.<sup>17</sup> concluded that gastritis is found more in females because females tend to lower their weight by managing food consumption frequency, size, and type, which would lead to gastritis. This statement was also made by Yunanda et al.<sup>19</sup> Women also tended to have gastritis due to lack of rest,<sup>3</sup> but some studies said the opposite.<sup>14</sup> The research by Liu et al.<sup>20</sup> revealed that gastritis risk factors depend on each country's economics, education, and health problems. Sipponen and Maaroos<sup>21</sup> also say that the prevalence of gastritis is increasing in line with the increasing age in developed countries. Still, other statements say that 50% of the prevalence of gastritis occurs in young age groups and even in childhood in developing countries. The research by Wulandari et al.<sup>22</sup> said that education level does not have any relationship with gastritis because both education levels can suffer from gastritis. A low education doesn't mean a low knowledge level because it can be obtained from non-formal education, such as information and personal experience, as their future references.<sup>23</sup> Still, Umasugi et al.'s<sup>4</sup> research said that a lack of education could cause a lack of behavior regarding gastritis prevention.

This research shows that there is a relation between smoking habits and gastritis incidence rate at productive ages. The mechanism of smoking influence on the mucosa is by stimulating HCl and pepsinogen, reducing pancreatic bicarbonate flow, removing the protective mucus layer, disturbing epithelial repair, microcirculation disruption, reducing gaster's motility and decreasing prostaglandin E<sub>2</sub>.<sup>5,24</sup> Smoking can lower prostaglandin synthesis in gastric mucosa and lower barrier function on stomach.<sup>5</sup> Smoking has various factors, such as reactive oxygen species, peroxynitrite, peroxynitrite, free radicals, and other reactive compounds that can cause oxidative stress, impacting endothelial function. This inflammation reaction will continue to occur and further lead to gastritis.<sup>5,12</sup> Smokers who have smoked for more than two years could increase gastric acid secretion and lower gastric pH.<sup>24</sup> This

research had similar findings by Cheng et al.,<sup>12</sup> where a smoking lifestyle could lead to gastritis. Still, some had different conclusions, which said there's no relationship between smoking habit and gastritis because there are other stronger influencing factors that might lead to gastritis occurrence, which are eating patterns and stress.<sup>11,14,25</sup>

This research showed a relation between coffee consumption and gastritis incidence rate at productive ages. Coffees are available in several forms. The chemical composition of coffee depends on green beans and roasted and brewed coffee. The green coffee beans depend on roasting, which can cause a Maillard reaction. This reaction can produce coffee acrylamide and reduce free chlorogenic acids and antioxidant forms. Meanwhile, roasted coffee consists of many components, and some effects are antioxidant, anti-inflammatory, antifibrotic, and antiproliferative. Coffee brewing also influenced the coffee composition. It depends on coffee solubilization in hot or cold water, separation of the water extract from the coffee ground, and water absorption. A recent study showed that hot and cold brew coffees have essential differences in total antioxidant capacity. But, whatever your coffee's methods, its components will affect your body.<sup>8</sup> The caffeine compound in coffee stimulates the secretion of HCl and gastrin through G cells, which are present in the gaster and duodenum but don't expedite gastric emptying.<sup>9,10</sup> Increased gastric acid secretion can increase the possibility of dyspepsia, gastritis, ulcers, and gastroesophageal reflux disease.<sup>9</sup> Caffeine can stimulate the central neural system and increase stomach activity to produce gastrin hormones in the stomach and pepsin, which are acidic, leading to irritation and stomach mucosa erosion.<sup>8,26</sup> Caffeine can accelerate acid production in gaster and produce excess gas, which can cause bloating.<sup>27</sup> This result is similar to the one Mahmoud et al.<sup>18</sup> conducted, which shows a significant association between gastritis and drinking coffee. The opposite study by Novitasary et al.<sup>11</sup> showed that the respondents rarely or not excessively drink coffee. Our body can absorb caffeine wholly and rapidly (95% within 45 minutes), and plasma levels increase within 30–60 minutes.<sup>28</sup> The International Agency for Research on Cancer also said there is evidence that coffee consumption helps reduce incidents of certain cancers, like colon, prostate, endometrium, melanoma, and liver.<sup>8</sup> Based on

International Food Information Council data, the safe consumption of coffee within the moderate range is up to 400 mg of caffeine per day or 3–5 cups per day for healthy adults.<sup>28</sup> Regarding this research, which reveals that most of the subjects consumed >3 glasses of pure coffee per day (black coffee), although every individual experiences different effects, it would risk them to contract gastritis. This result is the same with Santoso,<sup>27</sup> who said drinking coffee for an extended period and continuously with more than three glasses per day can irritate mucosa gaster, leading to gastritis and recurrence of gastritis.

The limitation of this research is that endoscopy examination is not used for gastritis diagnosis due to resource limitations. Therefore, the authors are unable to exclude gastritis contributors due to *Helicobacter pylori* infection or other irritant exposure.

## Conclusion

Smoking behavior and coffee consumption are related to the incidence rate of gastritis at productive ages in Makale Health Center, South Sulawesi.

## Conflict of Interest

The authors declare that there is no conflict of interest.

## Acknowledgment

The authors would like to thank the staff and all participants from Makale Health Center, South Sulawesi respondents, for their hospitality and cooperation in the research.

## References

1. Andreas, Tambunan LN, Baringbing EP. Hubungan pola makan dengan kejadian gastritis di Puskesmas Marina Permai Kota Palangka Raya. JSM. 2022;8(3):159–65.
2. Jusuf H, Adityaningrum A, Yunus R. Determinan kejadian gastritis pada mahasiswa. JHSJ. 2022;4(2):108–18.
3. Feyisa ZT, Woldeamanuel BT. Prevalence and associated risk factors of gastritis among patients visiting Saint Paul Hospital Millennium Medical College, Addis Ababa, Ethiopia. PLoS One. 2021;16(2):e0246619.
4. Umasugi MT, Souliassa FF, Susanti I, Latuperissa GR. The effect of health education on gastritis prevention behavior among high school students. J Ners. 2020;15(2):476–79.
5. Jîtcă G, Cîmpean T, Tero-Vescan A, Vari CE, Filip C, Ósz BE. Retrospective study on the frequency and causative mechanisms of chronic gastritis in an integrative medicine center in Tîrgu-Mureş. Acta Medica Transilvanica. 2018;23(3):67–9.
6. Sujatmiko B, Anggaeni TK, Wismandanu O, Indraswari N. Smoking behavior in Indonesian society based on the 5<sup>th</sup> Indonesian Family Life Survey data. In: Proceedings the 6th International Conference on Public Health; 2019 October 23–24; Surakarta, Indonesia. Surakarta: International Conference on Public Health; 2019. p. 110.
7. Naisali MN, Putri RSM, Nurmaningsari T. Hubungan perilaku merokok dengan kejadian gastritis pada mahasiswa teknik sipil Universitas Tribhuwana Tunggaladewi Malang. Nurs News. 2017;2(1):304–17.
8. Iriondo-DeHond A, Uranga JA, del Castillo MD, Abalo R. Effects of Coffee and its components on the gastrointestinal tract and the brain-gut axis. Nutrients. 2020;13(1):88.
9. Nehlig A. Effects of coffee on the gastrointestinal tract: a narrative review and literature update. Nutrients. 2022;14(2):399.
10. Ilham MI, Haniarti, Usman. Hubungan pola konsumsi kopi terhadap kejadian gastritis pada mahasiswa Muhammadiyah Parepare. J Ilm Manusia Kesehat. 2019;2(3):433–46.
11. Novitasary A, Sabilu Y, Ismail CS. Faktor determinan gastritis klinis pada mahasiswa di Fakultas Kesehatan Masyarakat Universitas Halu Oleo tahun 2016. Jimkesmas. 2017;2(6):2864.
12. Cheng Y, Keita M, Si XB. Investigation on risk factors of chronic gastritis in the population of Conakry in Guinea. Indian J Pharm Sci. 2021;Spec 1:1–5.
13. Hairuddin K, Yulis DM, Sahalessy Y, Tomaso J, Rumaseb E, Said A, et al. Factors associated with the incidence of gastritis in patients at Faisal Islamic Hospital Makassar. IJHS. 2023;1(2):156–61.
14. Khandelwal V, Deshmukh S, Aurangabadi K. Epidemiology and associated risk factors of gastritis in patients at District General Hospital, Amravati. IJHSR. 2024;14(2):93–

100.

15. Arumsari D, Martini S, Artanti KD, Widati S. The description of smoking degree based on Brinkman index in patients with lung cancer. *JBE*. 2019;7(3):249–56.
16. Sani W, Tina L, Jufri NN. Analisis faktor kejadian penyakit gastritis pada petani nilam di wilayah kerja Puskesmas Tiworo Selatan Kab. Muna Barat Desa Kasimpa Jaya tahun 2016. *Jimkesmas*. 2019;4(2):1-12.
17. Putri RSM, Agustin H, Wulansari. Hubungan pola makan dengan timbulnya gastritis pada pasien di Universitas Muhammadiyah Malang Medical Center (UMC). *J Keperawatan* 2010;1(2):156–64.
18. Mahmoud SS, Gasmi FM, Solan YO, Al-Harbi FA, Al-Harbi SA, Hummadi TA, et al. Prevalence and predictors of gastritis among patients attending health care facilities in Jazan, KSA. *Int J Preventive Public Health Sci*. 2016;2(1):1–7.
19. Yunanda FT, Wahyurianto Y, Retna T, Triana W. Gambaran faktor penyebab terjadinya gastritis di Desa Tlogowaru wilayah kerja Puskesmas Temandang Kabupaten Tuban. *JMI*. 2023;2(8):1742–57.
20. Liu Y, Zhang J, Guo Y, Tian S, Wu Y, Liu C, et al. Global burden and risk factors of gastritis and duodenitis: an observational trend study from 1990 to 2019. *Sci Rep*. 2024;14(1):2697.
21. Sipponen P, Maaros HI. Chronic gastritis. *Scand J Gastroenterol*. 2015;50(6):657–67.
22. Wulandari RH, Kalsum U, Izhar MD. Determinan yang berhubungan dengan kejadian gastritis di wilayah kerja di Puskesmas Tanjung Pinang Kota Jambi. *JKMI*. 2022[Unpublished]. Available at <https://repository.unja.ac.id/39278/1/JURNAL%20RIZKY%20HARTATI%20WULANDARI.pdf>.
23. Rujiantie F, Richard SD, Sulistyarini T. Pengetahuan pasien tentang faktor penyebab gastritis. *J STIKES* 2018;11(1):1–10.
24. Berkowitz L, Schultz BM, Salazar GA, Pardo-Roa C, Sebastian VP, Álvarez-Lobos MM, et al. Impact of cigarette smoking on the gastrointestinal tract inflammation: opposing effects in crohn's disease and ulcerative colitis. *Front Immunol*. 2018;9:74.
25. Firdous J, Muhamad N, Latif NA, Syazwani D, Hidayah N, Nurulnasuha, et al. A descriptive study on lifestyle factors influencing gastritis among University Students of UniKL RCMP in Malaysia. *Indian Journal of Natural Sciences* 2016;6(35):10753–6.
26. Arikah, Muniroh L. Riwayat makanan yang meningkatkan asam lambung sebagai faktor risiko gastritis. *Gizi Indones*. 2015;38(1):9–20.
27. Santoso P. Coffee consumption with the incidence of gastritis: literature review. *JANH*. 2023;5(2):225–32.
28. International Food Information Council. Health Professional Webinar for Registered Dietitians. Myths and facts about caffeine: an expert evaluation of the current science [Internet]. Washington: International Food Information Council; 2021 [cited 2023 May 15]. Available from: <https://ifc.org/wp-content/uploads/2021/04/Caffeine-Webinar.pdf>.

# Smoking Habit and Coffee Consumption with Gastritis.docx

*by* Eveline Margo FK

---

**Submission date:** 16-Jul-2024 09:20AM (UTC+0700)

**Submission ID:** 2417548326

**File name:** Smoking\_Habit\_and\_Coffee\_Consumption\_with\_Gastritis.docx (46.67K)

**Word count:** 3374

**Character count:** 20497

## Smoking Habit and Coffee Consumption with Gastritis Incidence Rate at Productive Ages

### Abstract

Gastritis is a disease caused by an inflammation of mucous and sub mucosal lining of the stomach. Several studies show that there are influence of smoking habit and coffee consumption toward gastritis incidence rate. The study was to determine the correlation between smoking habit and coffee consumption toward gastritis at productive ages. The research is an analytical observational study with a cross-sectional design, and was conducted in Puskesmas Makale, South Sulawesi. Required data was gathered by conducting guided interview by using a questionnaire that includes age range, sex, and education level. Brinkman Index were used to measure smoking habit, coffee consumption and gastritis questionnaire. Data was analyzed by using chi-square test with significance level at  $p=0,05$ . From total of 115 respondents, there are 69.6% included in the age range of 26-35 years old, 58.3% have mild smoking habit, 56.5% have severe coffee consumption and 85.2% of them are suffering gastritis. There are correlation between smoking habit and coffee consumption with gastritis with  $p=0.029(p<0.05)$ . The conclusion is there are relation between smoking habit and coffee consumption with gastritis incidence.

**Keywords:** coffee, gastritis, productive age, smoking

### INTRODUCTION

Gastritis is a common disease that often occurs in everyone and is a health problem with a fairly high prevalence. Lifestyle is an important factors for the occurance of gastritis.<sup>(1,2)</sup> Gastritis is an inflammation of mucous and sub <sup>11</sup> mucosal lining of the

stomach which can be caused by several factors, such as smoking habit, coffee drinking habit, and drug consumption for example aspirin or NSAID. <sup>(3)</sup>

<sup>4</sup> The incidence of world gastritis is around 1.8-2.1 million of the population every year. Based on research conducted by World Health Organization (WHO), gastritis incidents occurred high in USA (47%), India (43%), Canada (35%), China (31%). <sup>4</sup> Based on data from the Ministry of Health of the Republic of Indonesia (2019), there are 40.8% gastritis incidents occurred in Indonesia.<sup>(1,4)</sup> Gastritis patient prevalence often found on the age range of 20 to 44 years old. A research conducted by George University of Medicine and Pharmacy Tirgu-Mures mentioned that gastritis prevalence on people with coffee consumption and smoking habit often occurred on they who are 30 to 49 years old.<sup>(5)</sup>

Smoking is one of many modern lifestyles which nowadays not only found on adults, but also on children and young adults. According Indonesian Family Life Survey (IFLS5) data, the percentage of smoking in Indonesia was 58% with 95% dominated by male with average age was 19 years old. Most of them (48%) was kretek-per filtered as the common cigarette.<sup>(6)</sup>

Smoking habit can cause gastric disturbance because nicotine in cigarette will pucker and disrupt blood vessels in the stomach, which will further cause irritation that triggers excessive increase of gastric acid production.<sup>(7)</sup>

Other causing factor that influence gastritis occurrence is the habit to consume coffee/caffeine. Long-term and high frequency caffeine consumption (> 3 glasses a day) will accelerate gastric acid production that would irritate mucosa. When mucosa in the stomach is irritated, hydrochloride acid (HCl) diffusion towards the mucosa will disrupt mucosa and eliminate its ability to protect stomach from HCl and pepsin auto digestion.<sup>(3,5,8,9)</sup> A research conducted by Ilham MI et al. stated that there caffeine compound in coffee can accelerate gastric acid production stimulation and causes

gastritis.<sup>(10)</sup> However, different result was published by Novitasary A et al. which said that no relationship between coffee consumption and gastritis which it is highly depended on the type of the coffee, where in this research the subjects consume more instant coffee with lower caffeine compound than ground coffee beans.<sup>(11)</sup> A research by Cheng Y et al. said that there is relationship between smoking habit and gastritis,<sup>(12)</sup> but contrary researches by Novitasary A et al., Hairuddin K et al. and Khandelwal V et al., which said that there is no relationship between smoking habit and gastritis occurrence because there are factors with stronger influence on gastritis such as consumption habit and stress.<sup>(11,13,14)</sup>

Based on the description above, authors are interested to research on relation between smoking habit and coffee consumption with gastritis incidence rate at productive ages which can further identify the contributing factors to preventive and curative source for gastritis incidence rate.

## METHOD

This research is a type of observational analytical study using cross-sectional design. This study searched for a relation between smoking habits and coffee consumption with gastritis incidence rate in the Puskesmas Makale, South Sulawesi, which was conducted in February 2020. Subjects were men and women aged 26-45 years at the study site. Subjects in this study was taken through simple random sampling on subjects that accordance with the inclusion criteria, namely men or women aged 26-45 years, have smoking habits and/or coffee consumption, are not taking NSAIDs and/or aspirin and are willing to sign informed consent. Determination of sample size is done by using an <sup>1</sup>infinite-finite population formula with a significance level of 95% from 1.96, and with gastritis prevalence in Indonesia of 40.8%<sup>(1)</sup> where

the measurement accuracy is 0.05 and 115 subjects are willing to participate for this study.

Data collection was collected through questionnaire filling that gathers information regarding identity (name, age, sex, and education level), Brinkman index questions to measure the amount of cigarette consumed in a day multiplied by length of smoking habit in year with valuation category of light between 0-199, medium between 200-599, and heavy >600.<sup>(15)</sup> The questionnaire also measure the length and frequency of coffee consumption practiced by the subjects with categories of light with a frequency of  $\leq 3$  glasses/day, and heavy with a frequency of  $> 3$  glasses/day.<sup>(16)</sup> Gastritis questionnaire was also included to measure the perceived gastritis symptom complaints with valuation of  $\geq 12$  and  $< 12$  for non-gastritis complaints.<sup>(2)</sup>

#### ETHICAL CLEARENCE

This research is already approved by Research Ethics Commission of the Faculty of Medicine, Trisakti University, Jakarta, with approval number of 11/KER-FK/1/2019.

#### RESULTS

Univariate analysis was used to determine the frequency distribution of respondents' characteristics in the form of age, sex, education level, smoking habits, coffee consumption, and gastritis events of the research subjects.

Based on table 1, from of 115 total subjects in the Puskesmas Makale, 80 subjects (69.6%) were dominated by ages 26-35 years and 65 subjects (56.5%) were males, 80 subjects (69.6%) have a high education level, 67 subjects (58.3%) are in the mild group for smoking habit, and 65 subjects (56.5%) are severe consumers of coffee. In the assessment of the incidence of gastritis, it was known that 98 subjects (85.2%) had gastritis.

Table 1. Characteristic Data		
Frequency Distribution	Frequency (n)	Percentage (%)
<b>Age (years)</b>		
26-35	80	69.6
36-45	35	30.4
<b>Sex</b>		
Male	65	56.5
Female	50	43.5
<b>Level of education (years)</b>		
Low (< 9)	35	30.4
High (≥9)	80	69.6
<b>Smoking habit</b>		
Mild (< 200)	67	58.3
Moderate (200-599)	48	41.7
Severe (> 600)	3	2.6
<b>Coffee consumption</b>		
Mild (≤3)	50	43.5
Severe (>3)	65	56.5
<b>Gastritis occurrences</b>		
Gastritis	98	85.2
Non-Gastritis	17	14.8

Bivariate analysis with chi-square test was used to determine the relation between socio-demographic factors (age, sex, and level of education), smoking habits and coffee consumption on the incidence of gastritis at the age of 26-45 years.

**Table 2. Bivariate Relationship between Demographic Characteristics, Smoking Habits and Coffee Consumption with Gastritis Incidence Rate at productive ages**

Variable	Gastritis Occurrence				<i>p value</i>
	Gastritis		Non-Gastritis		
	(n)	(%)	(n)	(%)	
<b>Age (years)</b>					
26-35	68	59.1	12	10.4	0.921 <sup>€</sup>
36-45	30	26.1	5	4.3	
<b>Gender</b>					
Male	55	47.8	10	8.7	0.836 <sup>€</sup>
Female	43	37.4	7	6.1	
<b>Level of education (years)</b>					
Low (< 9 )	32	27.8	3	2.6	0.214 <sup>€</sup>
High (≥ 9)	66	57.4	14	12.2	
<b>Smoking habit</b>					
Mild	53	46.1	14	12.2	<b>0.029<sup>€</sup></b>
Moderate-Severe	45	39.1	3	2.6	
<b>Coffee consumption</b>					
Mild	37	32.2	13	11.3	<b>0.003<sup>€</sup></b>
Severe	61	53	4	3.5	

n = amount; % = percentage; € = Chi-Square Statistical Test; p value <0.05

Table 2 shows that gastritis occurrence are found more on people in the age group of 26-35 years old with 68 subjects (59.1%) and from chi-square statistical test result, it is revealed that there is no significant relation between age group and gastritis incidence rate in subjects of 26-45 years of age (p=0.921). As much as 55 subjects (47.8%) are male with gastritis, which is slightly higher than female subjects. This shows that there is no significant difference between two sexes. Based on chi-square statistical test result, a value of p=0.836 was obtained. Based on that, we can conclude that there is no significant relation between sex and gastritis incidence rate on subjects at productive ages. On group with higher education level, there are 66 subjects (57.4%) that suffer from gastritis, which is more than the one with lower education. In this group chi-square statistical test result shows a value of p=0.214 which means that there is no relation between education level and gastritis incidence rate on subjects at productive age.

On bivariate result of smoker group, authors combine moderate and severe smoker because there are only 3 subjects who are severe smokers. The result obtained shows that light smokers group suffers more gastritis incidence with 53 subjects (46.1%). On chi-square test result, a value of  $p=0.029$  was obtained which means that there is a significant relation between smoking habit and gastritis on subjects at productive age. As for coffee consumption variable, gastritis incidence found more in heavy coffee consumer with 61 subjects (53%). Chi-square test result shows a value of  $p=0.03$  which means that there is a significant relation between coffee consumption and gastritis incidence. The result showed that the subjects consume more coffee can increase gastritis incidence rate at productive age.

## DISCUSSION

In this research, demographic characteristic (age, sex, and education level) are not related with gastritis incidence rate on subjects at 26-45 years of age. Gastritis tends to happen at younger age due to uncontrolled eating pattern caused by daily activity,<sup>(1,17)</sup> and the results by Feyiza ZT et al., showed younger aged at 18-28 years (57.5%) has tend to suffered gastritis because don't know about good health behaviors,<sup>(18)</sup> but according to Khandelwal V et al., aged 14-25 years and 36-45 years has more tend to be gastritis.<sup>(14)</sup> However, gastritis can also occur on older subjects because of a number of factors such as age increase, degenerative process, H. Pylori infection, and NSAID drug usage.<sup>(19)</sup> Putri RSM et al. research concluded that gastritis are found more on female, because female tend to lower their weight by managing food consumption frequency, size, and type which would lead to gastritis.<sup>(17)</sup> This statement was also made by Yunanda FT et al.<sup>(20)</sup> Women also had tendency to be gastritis, due to lack of rest,<sup>(18)</sup> but some said the opposite.<sup>(14)</sup> The research by Liu Y et al. revealed that risk factors of gastritis depends of each country's economics, education and health problems.<sup>(21)</sup> This statement is also said by Sipponen P et al., which the prevalence of

gastritis is increasing in line with increasing age in the developed countries, but there are other statement said that 50% prevalence gastritis occurs in young age groups and even in childhood in the developing countries.<sup>(22)</sup> The research by Wulandari RH et al. said that education level is does not have any relationship with gastritis because both education level can suffer gastritis.<sup>(23)</sup> A low education level doesn't mean a low knowledge level, because it can be obtained from non-formal education, such as information and personal experience as their future references,<sup>(24)</sup> but Umasugi MT et al. research said that lack of education level can caused lack of behavior regrading of gastritis prevention.<sup>(4)</sup>

This research shows that there is a relation between smoking habits with gastritis incidence rate at productive ages. The mechanism of smoking influence towards mucosa is by more stimulate HCl and pepsinogen, reduce pancreatic bicarbonate flow, remove proctective mucus layer, disturbs epithelial repair, micro circulation disruption, reduces gaster's motility and decreases prostaglandin E2.<sup>(5,25)</sup> Smoking can lower prostaglandin synthesize in gastric mucosa and lower barrier function on stomach.<sup>(5)</sup> Smoking habit has various factors such as **reactive oxygen species (ROS), peroxynitrite, peroxynitrate, free radicals and other reactive compounds that** can make **oxidative stress** so will impact the endothelial function. This inflammation reaction will continue to occur and further lead to gastritis. <sup>(5,12)</sup> Smoker **who have smoked for more than 2 years could increase** gastric acid **secretion** and lower gastric pH.<sup>(25)</sup> This research had similar finding by Cheng Y et al. where smoking lifestyle could lead to gastritis,<sup>(12)</sup> but some had different conclusion which said there're no relationship between smoking habit and gastritis because there are other stronger influencing factors that might lead to gastritis occurrence which are eating pattern and stress.<sup>(11,14,26)</sup>

This research showed that there is relation between coffee consumption with gastritis incidence rate at productive ages. Coffees are available in several form. <sup>3</sup> The chemical composition of coffee depends on green beans, roasted and brewed coffee. The green coffee beans are depended on roasting, which can occurs Maillard reaction. This reaction can produces acrylamide form in coffee, reduces free chlorogenic acids and antioxidant forms. Meanwhile, roasted coffee is consists of many components, and some effects are as antioxidant, anti-inflammatory, antifibrotic and antiproliverative. Coffee brewing also had influence on the coffee composition. It's depend on coffee solubilization in hot or cold water, <sup>8</sup> separation of the water extract from coffee ground and water absorption. Recent study showed <sup>3</sup> that hot and cold brew coffees have important differences in total antioxidant capacity. But, whatever your coffee's methods, it's components will effects on your body.<sup>(8)</sup> Caffeine compound in coffee stimulates secretion HCl and gastrin through G cells which present in gaster and duodenum, but doesn't expedite gastric emptying.<sup>(9,10)</sup> Increased gastric acid secretion can increases the possible of dyspepsia, gastritis, ulcers and gastro-esophageal reflux disease (GERD).<sup>(9)</sup> Caffeine can stimulate central neural system and increase stomach activity to produce gastrin hormone in the stomach and pepsin with acidic nature and this would lead to irritation and stomach mucosa erosion.<sup>(8,27)</sup> Caffeine can accelerates acid production in gaster, produces excess gas which can causes bloating.<sup>(28)</sup> This result is similar as the one conducted by Mahmoud SS et al. which shows that there is a significant assosiation between gastritis and drinking coffee.<sup>(19)</sup> The opposite study by Novitasary A showed that the respondents rarely or not excessively drank coffee.<sup>(11)</sup> Our body can absorb caffeine rapidly and completely (95% within 45 minutes) and plasma levels increase within 30-60 minutes.<sup>(29)</sup> In addition to <sup>8</sup> International Agency for Research on Cancer (IARC) <sup>3</sup> said there is evidence that coffee consumption help to reduce incident of certain cancer, like colon, prostate, endometrium, melanoma and

liver.<sup>(8)</sup> Based on International Food Information Council (IFIC) data, the safe consumption of coffee within the moderate range is up to 400 mg caffeine per day or 3-5 cups per day for healthy adults.<sup>(29)</sup> Regarding this research which reveals that most of the subjects consumed >3 glasses of pure coffee per day (black coffee), although every individual experience different effects it would risk them to gastritis. This results is the same with Santoso, which said drinking coffee habit for long period and continuously with more than three glasses per day can irritate mucosa gaster, that will lead to gastritis and recurrence of gastritis.<sup>(28)</sup>

The limitation in this research is that gastritis diagnosis is not strengthened by endoscopy examination due to resource limitation, which unable the authors to exclude gastritis contributor due to H. Pylori infection or due to other irritant exposure.

## CONCLUSION

Smoking behavior and coffee consumption are relation with gastritis incidence rate at productive ages in Puskesmas Makale , South Sulawesi.

## DECLARATION

6

*The authors declare that there is no conflict of interest*

## ACKNOWLEDGMENT

The authors would like to express their gratitude to the Puskesmas Makale, South Sulawesi for their hospitality and cooperation in the research.

## REFERENCES

1. Andreas, Tambunan LN, Baringbing EP. Hubungan Pola Makan dengan Kejadian Gastritis di Puskesmas Marina Permai Kota Palangka Raya. JSM 2022; 8(3):159-65. DOI: <https://doi.org/10.33084/jsm.vxiix.xxx>
2. Jusuf H, Adityaningrum A, Yunus R. Determinan Kejadian Gastritis Pada Mahasiswa. JHSJ 2022; 4(2):108-18. DOI: <https://doi.org/10.37311/jhsj.v4i2.15171>
3. Feyisa ZT, Woldeamanuel BT. Prevalence and associated risk factors of gastritis among patients visiting Saint Paul Hospital Millennium Medical College, Addis Ababa, Ethiopia. Plos one 2021;16(2):1-16. <https://doi.org/10.1371/journal.pone.0246619>
4. Umasugi MT, Souliissa FF, Susanti I, et.al. The Effect of Health Education on Gastritis Prevention Behavior Among High School Students. Jurnal Ners. 2020;15(2):476-79. DOI: [10.20473/JN.V15I1SP.20515](https://doi.org/10.20473/JN.V15I1SP.20515)
5. Jitca G, Cimpean T, Tero-Vescan A, Vari CE, Filip C, Osz BE. Retrospective Study on The Frequency and Causative Mechanisms of Chronic Gastritis in An Integrative Medicine Center in Tirgu-Mures. AMT 2018;23(3):67-9. Available at : <https://www.researchgate.net/publication/327916560>
6. Sujatmiko B, Anggaeni TK, Wismandanu O, et.al. Smoking Behavior in Indonesian Society Based on The 5<sup>th</sup> Indonesian Family Life Survey Data. ICPH 2019. <https://doi.org/10.26911/the6thicph.02.17>
7. Naisali MN, Putri RSM, Nurmaningsari T. The Relationship between Smoking Behavior and the Occurrence of Gastritis in Civil Engineering Students of the University of Tribhuwana Tungadewi Malang. Nursing News 2017;2(1):305-17. <https://publikasi.unitri.ac.id/index.php/fikes/article/download/172/206>
8. Iriondo-DeHond A, Uranga JA, del Castillo MD, Abalo R. Effects of Coffee and Its Components on the Gastrointestinal Tract and the Brain-Gut Axis. Nutrients 2021;13(88):1-33. <https://doi.org/10.3390/nu13010088>
9. Nehlig A. Effects of Coffee on the Gastro-Intestinal Tract: A Narrative Review and Literature Update. Nutrients 2022; 14(399):1-31. <https://doi.org/10.3390/nu14020399>
10. Ilham MI, Haniarti, Usman. The Relationship of Coffee Consumption Pattern to the Occurrence of Gastritis in Muhammadiyah Parepare Students. Jurnal Ilmiah Manusia dan Kesehatan 2019;2(3):433-46. <https://doi.org/10.31850/makes.v2i3.189>
11. Novitasary A, Sabilu Y, Ismail CS. Determinants of Clinical Gastritis in 2016 Halu Oleo University's Faculty of Public Health. Jimkesmas 2017;2(6):1-11 <https://www.neliti.com/publications/183949/faktor-determinan-gastritis-klinis-pada-mahasiswa-di-fakultas-kesehatan-masyarakat>
12. Cheng Y, Keita M, Si XB. Investigation on Risk Factors of Chronic Gastritis in the Population of Conakry in Guinea. IJPS 2021; 1:1-5. DOI: 10.36468/pharmaceutical-sciences.spl.161
13. Hairuddin K, Yulis DM, Sahalessy Y, Tomaso J, Rumaseb E, Said A, Lestari SA. Factors Associated with the Incidence of Gastritis in Patients at Faisal Islamic Hospital Makassar. IJHS 2023;1(2):156-61. <https://doi.org/10.59585/ijhs.v1i2.78>
14. Khandelwal V, Deshmukh S, Aurangabadi K. Epidemiology and Associated Risk Factors of Gastritis in Patients at District General Hospital, Amravati. IJHSR 2024;14(2):93-100. <https://doi.org/10.52403/ijhsr.20240212>
15. Arumsari D, Martini S, Artanti KD, et. al. The Description of Smoking Degree Based on Brinkman Index in Patients with Lung Cancer. JBE 2019; 7(3): 249-56. <https://e-journal.unair.ac.id/JBE/article/view/12184>

16. Sani W, Tina L, Jufri NN. Analisis Faktor Kejadian Penyakit Gastritis pada Petani Nilam di Wilayah Kerja Puskesmas Tiworo Selatan Kab. Muna Barat Desa Kasimpa Jaya Tahun 2016. *JIMKESMAS* 2016;1(4):1-12. Available at: <https://media.neliti.com/media/publications/184565-ID-analisis-faktor-kejadian-penyakit-gastri.pdf>
17. Putri RSM, Agustin H, Wulansari. Hubungan Pola Makan dengan Timbulnya Gastritis Pada Pasien di Universitas Muhammadiyah Malang *Medical Center* (UMC). *Jurnal Keperawatan* 2010;1(2):156-64. Available at: <https://ejournal.umm.ac.id/index.php/keperawatan/article/view/406/3366>
18. Feyisa ZT, Woldeamanuel BT. Prevalence and Associated Risk Factors of Gastritis Among Patients Visiting Saint Paul Hospital Millennium Medical College, Addis Ababa, Ethiopia. *PloS One* 2021;16(2):1-16. <https://doi.org/10.1371/journal.pone.0246619>
19. Mahmoud SS, Gasmi FM, Solan YO, et al. Prevalence and Predictors of Gastritis among Patients Attending Health Care Facilities in Jazan, KSA. *Int J of Preventive and Pub health Sci* 2016;2(1):1-7. <http://dx.doi.org/10.17354/ijpphs/2016/18>
20. Yunanda FT, Wahyurianto Y, Retna T, Triana W. Gambaran Faktor Penyebab Terjadinya Gastritis di Desa Tlogowaru Wilayah Kerja Puskesmas Temandang Kabupaten Tuban. *Global Health Science* 2023;8(1):17-26. <http://dx.doi.org/10.33846/ghs8104>
21. Liu Y, Zhang J, Guo Y, Tian S, Wu Y, Liu C, et.al. Global Burden and Risk Factors of Gastritis and Duodenitis: an Observational Trend Study from 1990 to 2019. *Scientific Reports* 2024;14(2697):1-12. <https://doi.org/10.1038/s41598-024-52936-1>
22. Sipponen P, Maaroos HI. Chronic Gastritis. *Scand J Gastroenterol.* 2015;50(6):657- 67. <https://doi.org/10.3109/00365521.2015.1019918>
23. Wulandari RH, Kalsum U, Izhar MD. Determinan yang Berhubungan dengan Kejadian Gastritis di Wilayah Kerja di Puskesmas Tanjung Pinang kota Jambi. *JKMI.* Available at; <https://repository.unja.ac.id/39278/1/JURNAL%20RIZKY%20HARTATI%20WULANDARI.pdf>
24. Rujiantie F, Richard SD, Sulistyarini T. Pengetahuan Pasien tentang Faktor Penyebab Gastritis. *Jurnal STIKES* 2018;11(1):1-10. Available at: <https://jurnal.stikesbaptis.ac.id/index.php/STIKES/article/view/520/433>
25. Berkowitz L, Schultz BM, Salazar GA, Pardo-Roa C, Sebastian VP, et.al. Impact of Cigarette Smoking on the Gastrointestinal Tract Inflammation: Opposing Effects in Crohn's Disease and Ulcerative Colitis. *Front Immunol* 2018;9(74):1-10. <https://doi.org/10.3389/fimmu.2018.00074>
26. Firdous J, Muhamad N, Latif NA, et al. A Descriptive Study on LifeStyle Factors Influencing Gastritis among University Students of UniKL RCMP in Malaysia. *Indian Journal of Natural Sciences* 2016;6(35):10753-6. [https://www.researchgate.net/publication/316472217\\_A\\_descriptive\\_study\\_on\\_lifestyle\\_factors\\_influencing\\_gastritis\\_among\\_university\\_students\\_of\\_UniKL RCMP\\_in\\_Malaysia](https://www.researchgate.net/publication/316472217_A_descriptive_study_on_lifestyle_factors_influencing_gastritis_among_university_students_of_UniKL RCMP_in_Malaysia)
27. Arikah, Muniroh L. Riwayat Makanan yang Meningkatkan Asam Lambung sebagai Faktor Risiko Gastritis. *Gizi indon* 2015;38(1):9-20. <http://dx.doi.org/10.36457/gizindo.v38i1.163>
28. Santoso P. Coffee Consumption with The Incidence of Gastritis: Literature Review. *JANH* 2023;5(2):225-32. <https://doi.org/10.55018/janh.v5i2.153>

29. Health Profesional Webinar for Registered Dietitians. Myths and Facts about Caffeine: An Expert Evaluation of the Current Science. International Food Information Council (IFIC) 2021. Available at: chrome-extension://efaidnbmnnnibpcajpcglclefindmkaj/https://ific.org/wp-content/uploads/2021/04/Caffeine-Webinar.pdf

# Smoking Habit and Coffee Consumption with Gastritis.docx

## ORIGINALITY REPORT

12%

SIMILARITY INDEX

11%

INTERNET SOURCES

6%

PUBLICATIONS

1%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://jbiomedkes.org">jbiomedkes.org</a> Internet Source	2%
2	<a href="http://newinera.com">newinera.com</a> Internet Source	2%
3	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	1%
4	<a href="http://repo.stikesicme-jbg.ac.id">repo.stikesicme-jbg.ac.id</a> Internet Source	1%
5	<a href="http://www.frontiersin.org">www.frontiersin.org</a> Internet Source	1%
6	<a href="http://submission.als-journal.com">submission.als-journal.com</a> Internet Source	1%
7	<a href="http://journal2.um.ac.id">journal2.um.ac.id</a> Internet Source	1%
8	Amaia Iriondo-DeHond, José Antonio Uranga, Maria Dolores del Castillo, Raquel Abalo. "Effects of Coffee and Its Components on the Gastrointestinal Tract and the Brain-Gut Axis", Nutrients, 2020 Publication	1%

9	healthdocbox.com Internet Source	<1 %
10	journals.umkt.ac.id Internet Source	<1 %
11	Submitted to Adtalem Global Education, Inc. Student Paper	<1 %
12	jurnal.uinsu.ac.id Internet Source	<1 %
13	www.ncbi.nlm.nih.gov Internet Source	<1 %
14	bmcinfectdis.biomedcentral.com Internet Source	<1 %

Exclude quotes      On

Exclude bibliography      On

Exclude matches

< 10 words

Kepada Yth.  
Dewan Redaksi Global Medical and Health Communication  
Fakultas Kedokteran Universitas Islam Bandung  
Jln. Hariangbanga No. 2 Bandung  
Telepon/faksimile: (022) 4321213  
<https://ejournal.unisba.ac.id/index.php/gmhc>

### PERNYATAAN HAK CIPTA (Copyright Statement)

Naskah yang berjudul:

Smoking Habit and Coffee Consumption with gastritis Incidence  
Rate at Productive Age

Nama penulis: 1. Elvi Septira Untawo  
2. Euline Margo  
3. \_\_\_\_\_  
.. \_\_\_\_\_

akan diterbitkan dalam jurnal Global Medical and Health Communication.

Penulis menyatakan bahwa

[Silakan beri tanda (√)],

✓ Kutipan utuh data sekunder (bentuk kata, angka, gambar, tabel) yang merupakan barang hak cipta (*copyright*), disalin (*reproduce*), digambar ulang (*redrawn*), ditabelkan (*reuse*) dalam versi sendiri, sudah seizin pemegang hak cipta (penulis, penerbit, organisasi).

✓ Kutipan sebagian data sekunder (bentuk kata dan angka) yang disalin (*reproduce*), digambar ulang (*redrawn*), ditabelkan (*reuse*) untuk pembandingan dengan data primer atau pelengkap tabel/gambar sendiri, sudah menuliskan referensi sesuai dengan format pengutipan data.

✓ Naskah ini asli dan penulis mengalihkan hak cipta (*transfer of copyright*) naskah ini kepada jurnal Global Medical & Health Communication.

Jakarta, 24 April 2024  
Tempat, tgl, bulan, tahun  
Disetujui oleh penulis utama

Mete

Nam



Elvi Septira Untawo

### Surat Pernyataan Etik Ethical Statement

Judul Artikel: ..... Etiologi, Mekanisme, dan Konsumsi Kopi dengan Kejadian Gasvitis pada  
Article Title: ..... Urea Produktif .....  
..... Sedasi, Habits, and Coffee Consumption with Gasvitis Incidence Rate .....  
..... at Productive Age .....

Nama Seluruh Penulis: ..... Eli Septira Wibawa .....  
Names of All Authors: ..... Gueline Mago .....  
.....

Dengan ini kami menyatakan bahwa:  
We hereby confirm that:

1. Artikel yang kami kirimkan adalah asli yang ditulis oleh nama-nama penulis yang tercantum di atas dan belum pernah dipublikasi pada media manapun;  
*The article we have submitted to the journal for review is original, has been written by the stated authors and has not been published elsewhere;*
2. Artikel tidak sedang dalam proses atau pertimbangan publikasi oleh jurnal lain dan tidak akan dikirimkan ke jurnal lain selama proses penelaahan oleh jurnal ini berlangsung;  
*The article is not currently being considered for publication by any other journal and will not be submitted for such review while under review by this journal;*
3. Artikel tidak mengandung pernyataan-pernyataan yang berbahaya, menentang hukum, dan mengancam individu atau hak-hak individu dan kelompok lain;  
*The article contains no libellous or other unlawful statements and does not contain any materials that violate any personal or proprietary rights of any other person or entity;*
4. Kami telah memperoleh izin tertulis dari pemilik hak cipta setiap pernyataan atau dokumen yang diperoleh dari produk ber-hak cipta, serta telah menyebutkan sumber referensi yang digunakan dalam artikel ini.  
*We have obtained written permission from copyright owners for any excerpts from copyrighted works that are included and have credited the sources in my article.*



Jakarta, 21 April 2024  
Tanda Tangan / Author signature(s) Tanggal / Date

Eli Septira Wibawa  
Nama / Name(s)