



CONTENT

CASE REPORTS

Hyper-Reactive Malarial Splenomegaly Syndrome (HMS): Management Challenges in a Rural Community in Southwest Nigeria – Case Report

ORIGINAL ARTICLES

Early Versus Late Administration of Intravenous Oxytocin Infusion in the Prevention of Blood Loss During Elective Cesarean Section – A Randomized-Controlled Trial

Assessment of Attitude and Knowledge Towards the End-Of-Life Care amongst Nurses Working at Kigali University Teaching Hospital, Rwanda

Prevalence of Anemia and Associated Socio-Economic Determinants amongst Malnourished Children Aged 6-59 Months, Centre Hospitalier Universitaire de Kigali (CHUK) – A Retrospective Observational Study

REVIEW ARTICLES

Acute Kidney Injury and Renal Replacement Therapy (RRT) in ICU: The Nephrology or Intensive Care Nurse Role

The Need for Interdisciplinary Collaboration between Dental Professionals, Nurses and General Practitioners to Control NCDs

Editorial Team

CHEF EDITOR

Leon Mutesa, MD, PhD, Professor, Center for Human Genetics, University of Rwanda

DEPUTY CHIEF EDITORS

Peter Cartledge, MD, BSc, MRCOB, MRCPCH, POME, MSc, Assistant Professor, Yale University, Rwanda Human Resources for Health (HRH) Program, University Teaching Hospital of Kigali, Rwanda

Jean Paul Rwabishama, MD, PhD, Professor at the University of Rwanda, Kigali, Rwanda, Paris-Est Creteil University, Paris, France

Joseph Mucumbizi, MD, MPH, Honorary Associate Professor, Department of Pediatrics, King Faisal Hospital, Kigali, Rwanda

MEDICAL EDITOR & EDITORIAL ASSISTANT

Christian Nsanzabaganwa, MD, Rwanda Military Hospital, Rwanda

Fidele Byiringiro, MD, MCS (ECSA), MMed Department of Surgery, Rwanda Military Hospital, Rwanda

DESKTOP PUBLISHER

Christian Nsanzabaganwa, MD, Rwanda Military Hospital, Rwanda

COMMUNICATIONS MANAGER

Oscar Mulderwa, MD, Department of pediatrics, University of Rwanda

ASSOCIATE EDITORS

Cameron Page MD, Internal Medicine, Brooklyn, New York, USA

Chantal Ingabire, PhD, Social & Community Health, College of Medicine and Health Sciences, University of Rwanda, Kigali

EDITORIAL BOARD MEMBERS

Jesse Raiten MD, Anesthesiology, Critical Care, Perioperative Medicine, University of Pennsylvania, Anesthesiology and Critical Care Department, Philadelphia, USA

Paulin Banguti, MD, Anesthesiology, Critical Care, Cardiac Anesthesiologist, College of Medicine and Health Sciences, University of Rwanda, Kigali

Brian Swan DDS, MPH, Dentistry, Cambridge Health Alliance, Cambridge, MA, USA

Elena Stauf DDS, MSc, PhD, Dentistry, Oral Medicine & Oral Pathology, Harvard School of Dental Medicine, HRH, Rwanda

Ladan Bashir MA, DMD, Dentistry, Washington DC, USA

Amelia Pousson MD, MPH, Emergency medicine, CHUK, Kigali, Rwanda

Giles Cattermole BM BCh FRCSM DTMBM, Emergency medicine & Medical ethics, King's College Hospital NHS Trust, London UK

EDITORIAL BOARD MEMBERS

Jesse Raiten MD, Anesthesiology, Critical Care, Perioperative Medicine, University of Pennsylvania, Anesthesiology and Critical Care Department, Philadelphia, USA

Paulin Banguti, MD, Anesthesiology, Critical Care, Cardiac Anesthesiologist, College of Medicine and Health Sciences, University of Rwanda, Kigali

Brian Swan DDS, MPH, Dentistry, Cambridge Health Alliance, Cambridge, MA, USA

Elena Stauf DDS, MSc, PhD, Dentistry, Oral Medicine & Oral Pathology, Harvard School of Dental Medicine, HRH, Rwanda

Ladan Bashir MA, DMD, Dentistry, Washington DC, USA

Amelia Pousson MD, MPH, Emergency medicine, CHUK, Kigali, Rwanda

Giles Cattermole BM BCh FRCSM DTMBM, Emergency medicine & Medical ethics, King's College Hospital NHS Trust, London UK

Katie Cartledge BSc, MRCOB, DRCCO, DFRPH, RCOR, Family medicine & Medical education, International dispensary & UK, Kigali Rwanda

Navin Kumar Devaraj MD MMed, Department of Family Medicine, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, 43400 Serdang, Selangor, MALAYSIA

Claude Mambu Muvunyi, MD, PhD, Microbiology & Infectious Diseases, College of Medicine and Health Sciences, University of Rwanda, Kigali

Aline Uwimana, MD, MPH, Tropical & Infectious Diseases, Rwanda Biomedical Center

Krzysztof Bujarski MD, Internal medicine & Neurology, Dartmouth-Hitchcock Medical Center, Lebanon, New Hampshire, USA

Norbert Lameire MD, Nephrology, Internal Medicine, Retired professor of medicine University of Cent, Belgium

Joseph H Friedman MD, Neurology, Providence, Boston, USA

Florence Masaisa, MD, PhD, Hematology & Internal Medicine, College of Medicine and Health Sciences University of Rwanda, Kigali

Dik J van Leeuwen MD, PhD, FASLD, Gastroenterology and Hepatology, The Geisel School of Medicine at Dartmouth College, Hanover NH, USA

Tim Walker MBBS(Hons), FRACP, MPHMT, Gastroenterology, Internal medicine & Tropical medicine, Department of Medicine, Calvary Mater Hospital, Newcastle, Australia

Linda Baxter CNM, MSN, Clinical Educator Nurse-Midwifery, Crest Barrington MA, USA

Cedline Chironde BSc, MSc, PhD, General Nursing, nephrology & Public Health, Human Resources for Health, Rwanda

Maria Kisher APRN, CNP, FNP-BC, FAANP, Clinical Educator & nursing, Essentia Health, Fargo, North Dakota, USA

Renee Pyburn PMHNP-BC, RN, APRN, MSN, PMHNP at Samaritan Behavioral Health, Watertown, NY USA

Sheila Shaibu BEd, MSc, PhD, General Nursing

Stephen Rulisa, MD, PhD, Reproductive Health, College of Medicine and Health Sciences, University of Rwanda, Kigali

Lucy Baxter BSc, MRCOB, FRCOphth, Ophthalmology, Moorfields Eye Hospital, London, UK

Douglas Blackall MD, MPH, Pathology, St Louis University, Missouri, USA

Justin Wane, MD, MMed, Pathology, King Faisal Hospital, Rwanda

Craig J Conrad MD, MPH, FAAP, Pediatric Hospitalist, Ochsner Health System, New Orleans, LA, USA

Jeanvier Hityezu MD, MMed, Pediatrics and critical care, Department of Pediatrics, School of Medicine and Pharmacy, University Teaching Hospital of Kigali, Kigali, Rwanda

Natalie McCall MD, MPH, FAAP, Pediatric emergency medicine, Yale University, Human Resources for Health, Rwanda

Tanya Rogo MD, MPH/MTM, Pediatric infectious diseases, Icahn School of Medicine at Mount Sinai, New York, NY, USA

Stefan Jansen, MSc, PhD, Psychology & Mental Health, College of Medicine and Health Sciences, University of Rwanda, Kigali

Vedatane Nshindwa, MD, MSc, Biostatistics & Public Health, School Public Health, University of Rwanda, Kigali

Jean Claude Byiringiro, MD, MMed, FCS(ECSA), Surgery (Orthopedics), Surgery (Education), College of Medicine and Health Sciences, University of Rwanda, Kigali

Michael Sinclair MD, Cardiothoracic surgery, CHUB, Butare, Rwanda

David Skavathi MD MPH, Surgery (general), Maine Medical Center, Portland Maine USA

Georges Nkalyuta, MD, MMed, FCS(ECSA), General Surgery, College of Medicine and Health Sciences University of Rwanda, Kigali

Published: 2024-07-12

Articles

[Impact of educational interventions on knowledge, attitude, practice toward pharmacovigilance and adverse drug reaction reporting among healthcare professionals at the University Teaching Hospital, Rwanda](#)

O. Uwambayimana Cashumba, E. Muranyaneza, S. Twahirwa, A. Nzamukocha, V. Musingemana, D. Ryamukuru
9-19

DOI: [10.32464/rmj.v8i2.7](#)

[Download PDF](#)

[Barbiting Early Palliative Care for Suspected Lung Cancer Patient: a Case Series from Resource-Limited Setting in Indonesia](#)

Muhammad I. D. Rokasari, W. Prasetya, I. Ryatna, Yana K. Putri Fasal
5-8

DOI: [10.32464/rmj.v8i2.1](#)

[Download PDF](#)

[Factors Influencing the choice of place of delivery among mothers of children less than two years in Oyo State, Nigeria](#)

Nyemile S. Awonor, Abiodun O. Oloru, O. Oloru, A. Bisi, Isaac O. Aibunadigbo, Aissa Desmond A. Bello, Jerry O. Chukwura
20-27

DOI: [10.32464/rmj.v8i2.3](#)

[Download PDF](#)

[Prevalence and forms of violence against healthcare workers in the emergency department of a teaching hospital in Nigeria](#)

A. R. Bala, M. W. Agososo, D. O. Agbonbosa
28-35

DOI: [10.32464/rmj.v8i2.4](#)

[Download PDF](#)

[Enhancing sexual and reproductive health services uptake in Sub-Saharan Africa: the role of community Pharmacists in promoting self-care interventions: a systematic review](#)

S. Ndayishimiye, A. Odoakur, M. F. Mukanyangasi
36-46

DOI: [10.32464/rmj.v8i2.2](#)

[Download PDF](#)

[Supplementary vitamin D3 and vitamin D receptor polymorphisms affect blood vitamin D levels in type 2 diabetes mellitus in Indonesia](#)

Yenni R. Witasongka, E. Herwana, J. V. Kalumpu, B. P. Litan
47-57

DOI: [10.32464/rmj.v8i2.5](#)

[Download PDF](#)

[Assessing the Impact of Noninvasive Local Vibration Voice Therapy on Voice Disorders: A Comprehensive Review](#)

Roberto Tedeschi
58-63

DOI: [10.32464/rmj.v8i2.6](#)

[Download PDF](#)

Supplementary vitamin D3 and vitamin D receptor polymorphisms affect blood vitamin D levels in type-2 diabetes mellitus in Indonesia

Authors: Yenny^{1,*}; R. Wratsangka¹; E. Herwana¹; J. V. Kalumpiu¹, P. B. Liman¹

Affiliation: ¹Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

ABSTRACT

INTRODUCTION: There are no data on vitamin D receptor (VDR) gene single nucleotide polymorphism (SNP) influence on blood 25-hydroxy-cholecalciferol [25(OH)D] levels after supplementary vitamin D in Indonesian type 2 diabetes mellitus (T2DM) patients. This study evaluated the effects of the supplementary vitamin D3 and VDR gene SNPs rs1555410 and rs2228570 on blood 25(OH)D levels in T2DM cases.

METHODS: A randomized, double-blind placebo-controlled trial (RDPCT) was conducted at one research setting using 85 T2DM subjects divided into vitamin D group (VDG) and control group (CG) and receiving 5,000 IU/day vitamin D3 (cholecalciferol) or placebo once daily for 84 days. Levels of 25(OH)D were determined baseline and after supplementary vitamin D3 administration for 84 days. Circulatory 25(OH)D was assayed using ELISA. VDR polymorphisms were detected using sequencing.

RESULTS: Post-supplementary blood 25(OH)D rose appreciably from baseline in VDG for VDR rs1544410 genotypes G/G ($p=0.001$) and G/A ($p=0.010$), and in VDR rs2228570 genotypes T/T ($p=0.012$), T/C ($p<0.001$), and C/C ($p=0.001$). Post-supplementary VDG still contained 30.3% of subjects not reaching blood 25(OH)D ≥ 30 ng/mL.

In attaining blood 25(OH)D ≥ 30 ng/mL post-supplementation, VDR rs2228570 genotype T/C differed significantly from T/T (52.4% v. 100%; $p=0.027$), but there were no appreciable differences between genotypes C/C and T/T (78.6% v. 100%; $p=0.273$), as well as between VDR rs1544410 genotypes G/G and G/A (67.5% v. 100%; $p=0.542$).

CONCLUSION: Only 52.4% of subjects with VDR rs2228570 genotype T/C achieved sufficiently high blood 25(OH)D levels. VDR rs2228570 polymorphisms apparently influence T2DM response to supplementary vitamin D.

Keywords: Diabetes Mellitus, Vitamin D, Single Nucleotide Polymorphism, Indonesia

INTRODUCTION

Indonesia, with 10.7 million type 2 diabetes mellitus (T2DM) patients, occupies the 7th global

rank, with predicted T2DM prevalence rising to 16.6 million in 2045 [1]. T2DM vitamin D deficiency prevalence is higher than in the general global population, with prevalences of 63.2- 83.2% [2,3].

***Corresponding author:** Yenny, Department of Pharmacology and Clinical Pharmacy, Faculty of Medicine, Universitas Trisakti, Jakarta post code 11440, Indonesia; Email: yennyfarmako@trisakti.ac.id; Orchid id: 0000-0001-9390-5527; **Potential Conflicts of Interest (Col):** All authors: no potential conflicts of interest disclosed; **Funding:** All authors: Research funding from Universitas Trisakti No. 0303/PUF/FK/2021-2022; **Academic Integrity:** All authors confirm that they have made substantial academic contributions to this manuscript as defined by the ICMJE; **Ethics of human subject participation:** The study was approved by the local Institutional Review Board. Informed consent was sought and gained where applicable; **Originality:** All authors: this manuscript is original has not been published elsewhere; **Review:** This manuscript was peer-reviewed by three reviewers in a double-blind review process; **Type-editor:** Peter (USA).

Received: 21st February 2024; **Initial decision given:** 14th April 2024; **Revised manuscript received:** 14th April 2024; **Accepted:** 27th May 2024.

Copyright: © The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND) ([click here](https://creativecommons.org/licenses/by-nc-nd/4.0/)) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Biomedical Centre (RBC)/Rwanda Health Communication Center, P. O. Box 4586, Kigali. ISSN: 2079-097X (print); 2410-8626 (online)

Citation for this article: Yenny; R. Wratsangka; E. Herwana et al. Supplementary vitamin D3 and vitamin D receptor polymorphisms affect blood vitamin D levels in type-2 diabetes mellitus in Indonesia. Rwanda Medical Journal, Vol. 81, no. 2, p. 36-46, 2024. <https://dx.doi.org/10.4314/rmj.v81i2.5>

The latter are higher than in Europe, with a vitamin D deficiency prevalence of 40.4% (blood 25(OH)D) <20 ng/mL [4], while vitamin D deficiency prevalence in South Asia is 68% [5].

Vitamin D supplementation raises serum 25(OH)D concentration, influences health outcomes, and achieves maximum mortality rate reductions in developed countries from the commonest fatal diseases, such as cardiovascular disease and T2DM [6]. Gross vitamin D deficiency at 25(OH)D <12ng/mL (or <30nmol/L) is a health hazard [7] and should be corrected through vitamin D supplementation [8].

The utility of providing additional vitamin D to T2DM patients for glucose hemostasis and reducing insulin resistance is still debated, presumably because of the diversity of study populations for serum vitamin D status, supplementation dose and duration, methodology, gender, BMI, and ethnicity [9,10].

Vitamin D regulates signal transmission through the vitamin D receptor (VDR) [11]. The vitamin D-responsive element (VDRE) gene on chromosome 12q13.1 comprises nine exons and eight introns [11]. VDR activates and controls gene transcription via the target gene promoter VDRE. The VDR gene has over 470 polymorphisms [11], the most common being FokI (rs2228570 C to T) and BsmI (rs1544410 A to G)[11]. VDR rs1544410 at intron 8 regulates mRNA stability, thereby affecting gene expression [12]. VDR rs2228570 lies in the exon 2 start codon and changes the initiation sites [13]. VDR gene SNPs may influence VDR mRNA and protein stability and activity, resulting in a suboptimal response to supplementary vitamin D [14,15].

Multiple factors may affect post-supplementary blood 25(OH)D, such as sunshine exposure, aging, body mass index, calcium intake, supplementary vitamin D, and genetics [4,16]. It is currently uncertain whether VDR SNPs affect blood vitamin D levels after supplementary vitamin D in T2DM patients [17], due to difficulties in detecting gene involvement in the development of T2DM, because of possible minute differences in the gene and its interaction with genetic or non-genetic factors [18]. VDR SNP influence on increases in blood 25(OH)D levels that depict vitamin D condition, can be evaluated only with a randomized, double-blind placebo-controlled trial (RDBPCT).

Currently, few randomized double-blind placebo-controlled trials (RDBPCT) exist for evaluating

VDR rs1544410 and rs2228570 relationships with vitamin D therapeutic response. Waterhouse et al. [14]. Studied Australian elderly aged 60 – 80 years receiving vitamin D3 supplementation (30.000 vs. 60.000 IU/month) for 12 months and found rs2228570 was not associated with increases in 25(OH)D levels after supplementary vitamin D. A 20-week Chinese RDBPCT [15] on vitamin D deficiency cases receiving vitamin D3 at 2000 IU/day or placebo, demonstrated that rs2228570 showed stronger influences on 25(OH)D levels ($p < 0.04$). Post-treatment, VDR rs2228570-G, and its alleles had higher 25(OH)D levels ($p = 0.009$).

The RDBPCT of Cavalcante et al. [19]. on elderly females aged 68 ± 6 years with vitamin D insufficiency and receiving 200.000 IU vitamin D3 supplementation for 4 weeks, showed higher blood 25(OH)D concentrations in persons with BsmI genotypes BB/Bb ($p < 0.001$) who responded better to supplementary vitamin D than those with bb genotype.

A prospective case-control study involving 125 T2DM patients and 125 controls, revealed that low blood 25(OH)D and gene rs2228570 correlated with T2DM risk [20]. The inconsistent study results on VDR SNP relationships were caused by variations in pre-supplementary blood vitamin D levels, study designs, small sample sizes, vitamin D dose, duration of supplementary vitamin D, dietary intake, and ethnicity.

There are currently no reports on the causal relationship of VDR SNPs rs1544410 and rs2228570 with supplementary vitamin D among Indonesian T2DM patients, for which an RDBPCT is necessary. The outcome may be useful for vitamin D3 therapeutic dose personalization in T2DM patients to reduce morbidity and mortality rates. Primary research outcomes would be responses to supplementary vitamin D by comparing post-supplementary blood 25(OH)D values. Secondary outcomes would be the impact of rs1544410 and rs2228570 polymorphisms on blood 25(OH)D changes and attainment of blood 25(OH)D of ≥ 30 ng/mL

METHODS

Research design

This RDBPCT was conducted from June to August 2022 at Puskesmas Mampang in South Jakarta, with subjects signing informed consent. Study protocol approval was by the Research Ethics Committee,

Faculty of Medicine, Universitas Trisakti, under No. 001/KER/FK/1/2022.

Patients and intervention

The study subjects were Mampang area residents with T2DM. Inclusion criteria: males and females ≥ 18 years old, T2DM, HbA1c 7-8.5%, oral antidiabetic drug monotherapy, agreeing to follow-up controls. T2DM was diagnosed using American Diabetes Association criteria [21], namely fasting blood glucose ≥ 126 mg/dL, or 2-hour postprandial blood glucose ≥ 200 mg/dL and HbA1c $\geq 6.5\%$. Exclusion criteria: previously and currently on insulin therapy, suspect kidney disease (estimated glomerular filtration rate < 30 mL/min/1.73m²), abnormal liver function (SGPT 3 times normal upper limit), pregnant or lactating, allergy, hypercalcemia (plasma calcium > 2.65 mmol/L), or receiving daily supplementary vitamin D in preceding 84 days. Dropout criteria: blood 25(OH) D > 100 ng/mL, hypercalcemia, cholecalciferol hypersensitivity.

The study enrolled 115 subjects allocated by simple randomization to vitamin D (intervention) and control (placebo) groups at 1:1 ratio, which was conducted by personnel blinded to the intervention. VDG received daily vitamin D3 tablets containing 5,000 IU cholecalciferol, whereas controls received once-daily placebo tablets (120mg calcium), all tablets taken for 84 days. The vitamin D and placebo tablets contained in dark-colored glass bottles coded A and B were identical in visual, olfactory, and gustatory qualities. Participants and the statistician were all blinded to the origin of the tablets, whether from VDG or CG. Compliance with the intervention was determined by weekly counting the remaining tablets. VDG had 6 dropouts because of protocol non-compliance, returning to the home village, and diarrhea, while CG had 7 dropouts because of not agreeing to participate, protocol non-compliance, and nausea and vomiting. Subjects completing the study were 85 in number, consisting of 43 in VDG and 42 in CG (Figure 1).

The researchers were blinded to subject allocation in all study phases (recruitment, enrolment, data collection, and group assignment). For improving verification and compliance, all empty medication bottles were returned monthly to the cadres for evaluation of subject compliance to supplementation at the completion of the study. Subjects' complaints (potential adverse events)

were noted for recording. The principal investigator evaluated the complaints and their connection with the supplements. The development of all reported symptoms was monitored until the study was completed. The allocation codes stored by an independent third party were opened after the study was completed.

Measurements

On day zero (admission date), before administration of vitamin D tablets, subjects meeting recruitment criteria were interviewed to collect subject data on age, gender, and duration of diabetes mellitus, followed by blood collection at 08.00 and 9.00 a.m. local time.

Biochemical measurements

From each participant, 5 mL of venous whole blood was collected, with 3 mL for blood 25(OH) D determination and 2 mL in EDTA tubes for VDR SNP genotyping.

Blood 25(OH)D level was determined by chemiluminescent microparticle immunoassay (ARCHITECT 25-OH Vitamin D assay, Abbott), with measuring interval 8.0- 160.0 ng/mL (20.0- 400.0 nmol/L), limit of detection (LoD) ≤ 10.0 ng/mL, limit of quantitation (LoQ) ≤ 20 ng/mL, and imprecision $\leq 10\%$ within total CV.

Blood 25(OH)D data were presented as median \pm SD, and categorized according to Endocrine Society 2011 clinical practice guideline, with < 20 ng/mL defined as vitamin D deficiency, 20 – 30 ng/mL as vitamin D insufficiency, 30 – 100 ng/mL as vitamin D sufficiency [8].

Vitamin D receptor single nucleotide polymorphisms

Genomic DNA was extracted using QIAamp DNA Blood Mini Kit (QIAGEN) from 200 μ L EDTA blood and its purity and level were determined using a NanoDrop 2000 spectrophotometer (ThermoScientific). VDR SNPs were detected by PCR, followed by sequencing. PCR used MyTaq HS Red Mix Kit (Bioline) to amplify the target sequence using specific primers, namely for rs1544410: forward primer 5'- GGG AGT ATG AAG GAC AAA GAC C-3' and reverse primer 5'- CCC GCA AGA AAC CTC AAA TAA C-3' and for rs2228570: forward primer 5'-GCA CTG ACT CTG GCT CTG-3' and reverse primer 5'-TGG ACA TTG TAA GGA AGG AGA TG-3'. PCR amplification used 2 μ L DNA, 8.5 μ L nuclease-free water, 2x 12.5 μ L My Taq HS Red Mix,

and 1.0 μ l each of forward and reverse primers. Conditions: initial activation at 95oC for 5 minutes, denaturation at 95oC for 15 seconds, annealing at 58oC for 30 seconds, extension at 72oC for 30 seconds for 40 cycles, and final extension at 72oC for 3 minutes. PCR products on 2% agarose gel were visualized by electrophoresis. Purified PCR products were sequenced using BigDye Terminator Kit (Thermo Fisher Scientific) and ABI 3500 sequencer (Applied Biosystems). Sequence analysis was performed with BioEdit software to confirm mutations, which were compared to NCBI BLAST

database with accession number NG_008731.1.

Sample size

In each group, 40 subjects were required to detect a treatment difference at 5% two-sided significance level (0.05) and 90% study power. $\mu_1 - \mu_2$ = predicted between-group mean difference, estimated at 4; α_2 = expected population variance from the preliminary study, estimated at 1.02. To anticipate dropouts, the sample size became 95 for adequate power to detect outcome measure differences.




Figure 1: Flow of participants throughout trial

Statistical analysis

Continuous data of normal distribution were shown as mean and standard deviation (SD), continuous data of skewed distribution as median (minimum-maximum), and categorical data as percentages. Data were checked for normal distribution using the Kolmogorov-Smirnov test. Unpaired Student's t-test, Mann-Whitney U test, Chi-square test, and Fisher exact test were used for baseline comparisons between VDG and CG, based on the variable type and data distribution. Mann-Whitney test was used to compare blood 25(OH)D in VDG and CG at baseline and post-supplementation. Baseline and post-supplementation vitamin D status differences were evaluated using Fisher's exact test and Chi-square test. VDG baseline and post-supplementation blood 25(OH)D differences between rs1544410 genotypes were evaluated by the Mann-Whitney test. VDG baseline and post-supplementation blood 25(OH)D levels between rs2228570 genotypes were compared using the Kruskal-Wallis test, Wilcoxon signed rank test, and paired t-test at $p < 0.05$. Statistical analysis used SPSS for Windows version 23.

RESULTS

Baseline subject characteristics

We had 85 T2DM participants, with 68 (80%) females, mean age 55.8 ± 0.6 years, median T2DM duration 12 (1 – 36) months, and median blood 23(OH)D 11.6 (2.4 – 30.3) ng/mL. The most frequent rs1555410 genotype was G/G, comprising 79 subjects (92.9%), followed by G/A with 6 subjects (7.1%), without any T/T. Gene rs2228570 had genotypes T/C, C/C, and T/T with 38 (44.7%), 32 (37.6%), and 15 (17.6%) subjects, respectively. Vitamin D status was mostly deficient in 74 subjects (84.1%), insufficiency in 9 subjects (10.6%), and sufficiency in 2 subjects (2.4%). After group randomization, no significant differences were observed in age, gender, T2DM duration, and VDR genotype between VDG and CG (Table 1).

Baseline and post-supplementation blood 25(OH)D and vitamin D status

After vitamin D3 supplementation for 84 days, there was a much greater increase in VDG blood 25(OH)D than in CG (46(9.4-79.4) v. 14.4(6.9-38.3); ($p < 0.001$) (Table 2).

Table 1: Subject characteristics at the start of the study in VDG and CG

Characteristic	Treatment group		P-value
	VDG (n=43)	CG (n=42)	
Age (years)	56 (35 – 80)	56 (35 – 69)	0.396 ^a
Gender			
Male	8 (47.1)	9 (52.9)	0.745 ^b
Female	35 (51.5)	33 (48.5)	
Duration of DM (months)	12 (1 – 36)	12 (1 – 36)	0.967 ^a
VDR genotype (n,%)			
rs1544410			
G/G	40 (50.6)	39 (49.4)	1.000 ^c
G/A	3 (50)	3 (50)	
A/A	0 (0)	0 (0)	
rs 2228570			
T/T	8 (53.3)	7 (46.7)	0.614 ^b
T/C	21 (55.3)	17 (44.7)	
C/C	14 (43.8)	18 (56.2)	

Values presented as median (min-max) or n(%). Statistical analysis: ^aMann-Whitney test; ^bChi-square test; ^cFisher's exact test; $p < 0.05$ = statistically significant. VDG = vitamin D group; CG = control group

Table 2: Blood 25 (OH)D level and vitamin D status in VDG and CG at baseline versus 84 days after supplementation with vitamin D3.

Group	Start of study		p value	After supplementation		p-value
	VDG (n=43)	CG (n=42)		VDG (n=43)	CG (n=42)	
Blood 25(OH)D level (ng/mL)						
	10.5 (4.7 – 30.3)	13.05 (2.4 – 26.9)	0.264 ^a	46 (9.4 – 79.4)	14.4 (6.9 – 38.3)	0.001 ^{a*}
Vitamin D status						
Deficiency	40 (50.6)	34 (45.9)		3 (8.6)	32 (91.4)	
Insufficiency	2 (22.2)	7 (77.8)	1.000 ^c	10 (62.5)	6 (37.5)	0.001 ^{b*}
Sufficiency	1 (50)	1 (50)		30 (88.2)	4 (11.8)	

Groups: VDG= vitamin D3 5.000IU/day, CG = placebo
 Vitamin D status (blood 25(OH)D level): deficiency (< 20 ng/mL); insufficiency (20 – 30ng/mL); sufficiency (30 – 100 ng/mL). Statistical analysis: ^aMann Whitney test; ^bChi-square test; ^cFisher's exact test; *p<0.05 =statistically significant
 VDG = vitamin D group; CG = control group

Comparison of blood 25(OH)D levels by VDR genotype after supplementary vitamin D3 administration in VDG
 To find the effects of VDR genotype on post-supplementation blood 25(OH)D, 25(OH)D blood levels were compared between genotypes in VDG (Table 3). Blood 25(OH)D levels increased significantly above baseline after supplementary vitamin D3 in rs1544410 genotypes G/G [10.5 (4.7

– 30.5) v. 46.5 (9.4 – 79.4) ng/mL; p=0.001] and G/A [10.8 ± 1.3 v. 45.7 ± 7.2 ng/mL; p =0.010]. No great differences were found in 25(OH)D between rs1544410 genotypes G/G and G/A [46.5 (9.4 – 79.4) ng/mL v. 45.7 ± 7.2 ng/mL; p=0.924].
 As compared to baseline, post-supplementation rs2228570 blood 25(OH)D levels increased significantly for genotypes T/T [11.4 (6.2 – 17.9) v.

Table 3: Blood 25(OH)D levels in VDG by VDR genotype after supplementary vitamin D3 administration

VDG	Blood 25(OH)D level (ng/mL)		p-value
	Start of study (n=43)	84 days (n=42)	
VDR genotypes			
rs1544410			
G/G (n= 40)	10.5 (4.7 – 30.5)	46.5 (9.4 – 79.4)	0.001 ^{c*}
G/A (n= 3)	10.8 ± 1.3	45.7 ± 7.2	0.010 ^{d*}
P value	0.924 ^a	0.924 ^a	
rs2228570			
T/T (n=8)	11.4 (6.2 – 17.9)	61.2 (32.1 – 79.4)	0.012 ^{c*}
T/C (n=21)	10.5 (7.6 – 30.3)	34.2 (9.4 – 67.4)	<0.001 ^{c*}
C/C (n=14)	11.0 ± 4.4	43.7 ± 12.4	0.001 ^{d*}
P value	0.373 ^b	0.024 ^{b*}	

Statistical analysis: ^aMann-Whitney test; ^bKruskal-Wallis test; ^cWilcoxon signed rank test; ^dpaired t-test; *p value <0.05 = statistically significant
 VDG = vitamin D group; CG = control group

Table 4: Attainment of blood 25(OH)D levels in VDG by VDR genotype after supplementary vitamin D administration

VDR genotype	Blood 25(OH)D level		p-value
	< 30 ng/mL (n,%)	≥30 ng/mL (n,%)	
VDG			
rs 1544410			
G/G (n=40)	13 (32.5)	27 (67.5)	0.542
G/A (n=3)	0 (0)	3 (100)	
rs 2228570			
T/T (n=8)	0 (0)	8 (100)	0.027*
T/C (n= 21)	10 (47.6)	11 (52.4)	
C/C (n=14)	3 (21.4)	11 (78.6)	0.273

Statistical analysis: logistic regression test; *p value <0.05 = statistically significant
VDG = vitamin D group

61.2 (32.1 – 79.4); p=0.012], T/C [10.5 (7.6 – 30.3) v. 34.2 (9.4 – 67.4); p<0.001], and C/C [11.0 ± 4.4 v. 43.7 ± 12.4; p=0.001] (Table 3).

There were also significant differences in post-supplementation blood 25(OH)D between rs2228570 genotypes T/T, T/C, and C/C themselves [61.2 (32.1 – 79.4) ng/mL v. 34.2 (9.4 – 67.4) ng/mL v. 43.7 ± 12.4 ng/mL; p=0.024] (Table 3).

To determine the genotypes causing the post-supplementation differences in rs2228570, a post-hoc analysis was conducted, showing significant differences between genotypes T/T and T/C (p=0.015) and between T/T and C/C (p=0.017), but not between T/C and C/C (p=0.391).

Regarding blood 25(OH)D responses to supplementary vitamin D of rs1544410 genotypes G/G and G/A, among the 43 VDG subjects, only 30 (69.7%) attained blood 25 (OH)D levels ≥30 ng/mL. The same is true for the three rs2228570 genotypes T/T, T/C, and C/C (Table 4).

No prominent differences were found between rs1544410 G/G and G/A in attaining blood 25(OH) D ≥30 ng/mL (67.5% v. 100%; p=0.542). In rs2228570, significant differences occurred in blood 25(OH)D level attainment between T/C and T/T (52.4% v. 100%; p=0.027).

DISCUSSION

Vitamin D deficiency was found in 84.1% of subjects, with a median blood 23(OH)D level of

11.6 (2.4 – 30.3) ng/mL, which is much greater than in Europe, where 25(OH)D levels below 20 ng/mL and 12 ng/mL are observed in 40.4% and 13.0%, respectively, of the population [4]. Conversely, adult vitamin D deficiency prevalence in 5 South Asian countries was 68% [5]. Our results approximate those of earlier studies demonstrating that T2DM vitamin D deficiency prevalence is around 63.2 [83.2% [2,3], and that vitamin D deficiency is also found in tropical countries, such as Indonesia, with abundant sunshine for cutaneous vitamin D synthesis.

Vitamin D as a prohormone is available as vitamin D2 (ergocalciferol) in foods and vitamin D3 (cholecalciferol) in UV-exposed human skin. Generally, vitamin D deficiency is caused by low dietary intake and reduced cutaneous synthesis from inadequate sunlight exposure due to geographic location, skin color, age, indoor lifestyle, and cultural or religious practices[22,23]. Our greater vitamin D deficiency prevalence shows that foods and sunlight alone cannot maintain optimal vitamin D status, necessitating vitamin D supplementation.

Signal transmission in humans occurs through vitamin D binding with the vitamin D receptor (VDR) [11] expressed by insulin-sensitive tissues. Apparently, vitamin D may have a direct influence on insulin sensitivity and insulin receptor expression, thereby enhancing insulin-stimulated glucose transport. Vitamin D may also have an indirect influence [17] by reducing insulin resistance-inducing inflammatory responses [24]. A meta-analysis showed that vitamin D

supplements can raise blood 25(OH)D and reduce insulin resistance in T2DM [25]. However, systematic reviews of T2DM RCTs failed to find evidence for the efficacy of vitamin D supplements in glucose hemostasis and in decreasing insulin resistance [26,27]. According to another meta-analysis, vitamin D dosage, status, and length of supplementation affect the therapeutic response[9]. High-dose vitamin D supplementation produces greater effects in obese vitamin D-deficient Asians[9]. Irrespective of vitamin D supplementation's impact on T2DM patients, vitamin D deficiency should be corrected because vitamin D serves an essential function in calcium hemostasis and bone metabolism [22].

Our study showed that rs1555410 had genotype G/G in 79 (92.9%) subjects and no genotype A/A, whereas Chinese T2DM patients [17] had mostly rs1544410 genotype G/A (93.75%), but similarly no A/A. Our rs2228570 genotype proportions comprised T/C in 44.7%, C/C in 37.6%, and T/T in 17.6% of subjects, whereas the Chinese study [17] comprised rs2228570 genotype C/C in 59.8%, T/T in 23.2%, and C/T in 16.9% of subjects, showing that Asians vary in rs1544410 and rs2228570 gene proportions. Similarly, the research results of Sari et al. [28]. on healthy North Sumatran women differed from ours, because all subjects had heterozygous A>G for BsmI (rs1544410), and T>C for TaqI (rs2228570), showing that Indonesians also have differing VDR genotype proportions, supporting the supplementary vitamin D dosage personalization concept.

In our study, supplementary vitamin D at 5000 IU/day caused a 3.2-fold higher rise in blood 25(OH)D in VDG compared to CG [46 (9.4 – 79.4) ng/mL v. 14.4(6.9 – 38.3) ng/mL; $p=0.001$] (Table 2). Blood 25(OH)D may rise around 1 ng/mL (2.5 nmol/L) per 100 IU daily vitamin D3 supplement given for 56 - 84 days [29], although the supplementary dose may not be linearly correlated with blood 25(OH)D [30].

After 84 days of vitamin D supplementation at 5000 IU/day, among our 43 subjects, 13 (30.2%) subjects still did not attain blood 25(OH)D levels of ≥ 30 ng/mL (Table 4.) This agrees with the findings of Yao et al. [15], from a 140-day RDBPCT on 448 Chinese with vitamin D deficiency receiving 2000 IU/day vitamin D3 or placebo, where the vitamin D increased blood 25(OH)D, but could not overcome vitamin D deficiency in 25% of subjects. Al-Daghari et al. [31]. Studying T2DM patients on 2000 IU/day

supplementary vitamin D for 12 months showed that 42% of subjects still could not reach target blood 25(OH)D. Hu et al. [17]. in their study on T2DM subjects receiving supplementary vitamin D at 800 IU/day for 12 months, also found that 44.6% of subjects did not attain vitamin D sufficiency.

The influence of VDR SNP on supplementary vitamin D3 results remains unclear. We found a significant increase in post-supplementary blood 25(OH)D levels above baseline values in rs1544410 genotypes G/G ($p=0.001$) and G/A ($p=0.010$) which agrees with Cavalcante et al. [19]. showing that supplementary vitamin D significantly increased blood 25(OH)D in BB/Bb ($p=0.009$), but not in bb subjects.

In our VDG subjects with rs2228570, large differences were found in post-supplementary 25(OH)D blood concentrations of genotypes T/T, T/C, and C/C ($p=0.024$) (Table 3.). Post-hoc analysis showed differences between T/T and T/C ($p=0.015$) and between T/T and C/C ($p=0.017$), but not between T/C and C/C ($p=0.391$). In our study, rs2228570 apparently influenced supplementary vitamin D response in T2DM subjects. In T/C subjects, only 52.4% attained 25(OH)D ≥ 30 ng/mL, lower than in the other genotypes (Table 4).

One meta-analysis showed that TaqI and FokI polymorphisms may modulate supplementary vitamin D response for better results [32]. Our study confirms that the doses should be adapted (“personalized”) in subjects with rs2228570 genotype T/C for optimal benefits of supplementary vitamin D.

Our study results differ from those of Al-Daghari et al. [31]. on T2DM subjects with genotype-related differences in post-supplementary blood 25(OH)D, in that genotype T/T subjects evidenced better therapeutic responses than the other genotypes. Our results also differ from those of Hu et al. [17]. showing in T2DM subjects that rs2228570 genotypes T/C and T/T had no remarkable differences in blood 25(OH)D ($p=0.964$). In addition, Hu’s study and ours showed differences in subject characteristics regarding age, baseline blood 25(OH)D, and supplementary vitamin D dose. Hu’s subjects were aged 66.3 ± 9.1 years, whereas ours were 55.8 ± 0.6 years old. Our baseline blood 25(OH)D of $11.6 (2.4 – 30.3)$ ng/mL exceeded that of Hu’s 22.7 ± 1.9 ng/mL, presumably because Hu et al. used a lower vitamin D3 supplementation dose (800 IU) [17]. Another study found that lower baseline blood 25(OH)D levels were associated

with significantly higher blood 25(OH)D responses [15].

Opinions regarding optimal blood 25(OH)D levels in humans are inconsistent, with no uniform definition of vitamin D deficiency and insufficiency in different guidelines. The IOM recommends a minimum blood 25(OH)D concentration of 20 ng/mL (50 nmol/L), in connection with bone health [33]. However, the Endocrine Society recommends 25(OH)D levels exceeding 30 ng/mL (or 75 nmol/L) for preventing infections and obtaining other non-calcemic vitamin D benefits [8]. We showed that supplementary vitamin D3 at 5000 IU/day for 84 days still could not prevent 30% of subjects from attaining vitamin D sufficiency with blood 25(OH)D \geq 30 ng/mL. The Endocrine Society clinical practice guideline [8] recommends supplementary vitamin D3 for increasing vitamin D levels and determining blood 25(OH)D concentrations because 25(OH)D is the most frequent circulatory vitamin D, with a half-life of 14 – 21 days, and extremely useful for monitoring vitamin D status in persons at high risk of vitamin D deficiency.

Our study confirms the need for supplementary vitamin D dose personalization and blood 25(OH)D measurement in high-risk patients in relation to VDR SNPs, apart from the contradictory relationship of vitamin D deficiency in glucose hemostasis and insulin resistance reduction [20,26,27,34]. There is a need for dosage adjustment (“personalization”) in subjects with rs2228570 genotype T/C to obtain better gains from supplementary vitamin D. Our study results may provide inputs on management policies of T2DM patients susceptible to vitamin D deficiency, particularly in Indonesia.

In some populations, the interplay of genes and lifestyle may obscure the genetic component; therefore, studies on gene interactions with diet and physical activity are mandatory to confirm the relationship. Other longer-term RCTs with larger sample sizes are also necessary to better utilize the results of vitamin D supplements in patients with type 1 and type 2 diabetes.

We used an RDBPCT design that is best for measuring cause-and-effect relationships. We used an identical therapeutic procedure and supplementary vitamin D3 dosage to minimize subject variation.

One limitation of this study was that our subjects were Indonesian T2DM patients; therefore, our results may not apply to other nations. We also

did not account for physical activity, diet, sunlight exposure, BMI, and parathyroid hormone as confounders.

CONCLUSION

After vitamin D supplementation, blood 25(OH)D levels rose perceptibly, but a third of subjects still failed to attain blood 25(OH)D levels of \geq 30 ng/mL. VDR rs2228570 genotype T/C had only 52.4% of its subjects attaining a sufficiently large 25(OH)D level, but perceptibly lower than in genotypes T/T and C/C. VDR rs2228570 polymorphisms apparently influence T2DM response to supplementary vitamin D. There is a need for personalization of vitamin D dosage and blood 25(OH)D measurement in high-risk patients due to VDR SNPs.

REFERENCES

1. IDF Diabetes Atlas; 9th ed.; International Diabetes Federation, 2019; ISBN 978-2-930229-87-4. Available from: https://diabetesatlas.org/upload/resources/material/20200302_133351_IDFATLAS9e-final-web.pdf
2. Khudayar, M.; Nadeem, A.; Lodi, M.N.; Rehman, K.; Jawaid, S.I.; Mehboob, A.; Aleem, A.S.; Mirza, R.E.F.; Ahmed, M.; Abbas, K. The Association Between Deficiency of Vitamin D and Diabetes Mellitus Type 2 (DMT2). *Cureus* 2022, 14, e22221, doi:10.7759/cureus.22221.
3. Ac, A. Serum Vitamin D Levels in Persons with Type 2 Diabetes Mellitus in Lagos, Nigeria., doi:10.23937/2377-3634/1410133.
4. Cashman, K.D.; Dowling, K.G.; Škrabáková, Z.; Gonzalez-Gross, M.; Valtueña, J.; De Henauw, S.; Moreno, L.; Damsgaard, C.T.; Michaelsen, K.F.; Mølgaard, C.; et al. Vitamin D Deficiency in Europe: Pandemic? *Am. J. Clin. Nutr.* 2016, 103, 1033–1044, doi:10.3945/ajcn.115.120873.
5. Siddiquee, M.H.; Bhattacharjee, B.; Siddiqi, U.R.; MeshbahrRahman, M. High Prevalence of Vitamin D Deficiency among the South Asian Adults: A Systematic Review and Meta-Analysis. *BMC Public Health* 2021, 21, 1823, doi:10.1186/s12889-021-11888-1.
6. Nutrients | Free Full-Text | A Narrative Review of the Evidence for Variations in Serum 25-Hydroxyvitamin D Concentration Thresholds for Optimal Health Available online: <https://www.mdpi.com/2072-6643/14/3/639> (accessed on 29 February 2024).

7. Vitamin D Deficiency 2.0: An Update on the Current Status Worldwide | *European Journal of Clinical Nutrition* Available online: <https://www.nature.com/articles/s41430-020-0558-y> (accessed on 29 February 2024).
8. Holick, M.F.; Binkley, N.C.; Bischoff-Ferrari, H.A.; Gordon, C.M.; Hanley, D.A.; Heaney, R.P.; Murad, M.H.; Weaver, C.M.; Endocrine Society Evaluation, Treatment, and Prevention of Vitamin D Deficiency: An Endocrine Society Clinical Practice Guideline. *J. Clin. Endocrinol. Metab.* 2011, 96, 1911–1930, doi:10.1210/jc.2011-0385.
9. Farahmand, M.A.; Daneshzad, E.; Fung, T.T.; Zahidi, F.; Muhammadi, M.; Bellissimo, N.; Azadbakht, L. What Is the Impact of Vitamin D Supplementation on Glycemic Control in People with Type-2 Diabetes: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *BMC Endocr. Disord.* 2023, 23, 15, doi:10.1186/s12902-022-01209-x.
10. Musazadeh, V.; Kavyani, Z.; Mirhosseini, N.; Dehghan, P.; Vajdi, M. Effect of Vitamin D Supplementation on Type 2 Diabetes Biomarkers: An Umbrella of Interventional Meta-Analyses. *Diabetol. Metab. Syndr.* 2023, 15, 76, doi:10.1186/s13098-023-01010-3.
11. Molecular Epidemiology of Vitamin D Receptor Gene Variants | *Epidemiologic Reviews* | Oxford Academic Available online: <https://academic.oup.com/epirev/article/22/2/203/456955> (accessed on 29 February 2024).
12. Decker, C.J.; Parker, R. Diversity of Cytoplasmic Functions for the 3' Untranslated Region of Eukaryotic Transcripts. *Curr. Opin. Cell Biol.* 1995, 7, 386–392, doi:10.1016/0955-0674(95)80094-8.
13. Uitterlinden, A.G.; Fang, Y.; Van Meurs, J.B.J.; Pols, H.A.P.; Van Leeuwen, J.P.T.M. Genetics and Biology of Vitamin D Receptor Polymorphisms. *Gene* 2004, 338, 143–156, doi:10.1016/j.gene.2004.05.014.
14. Environmental, Personal, and Genetic Determinants of Response to Vitamin D Supplementation in Older Adults - PubMed Available online: <https://pubmed.ncbi.nlm.nih.gov/24694335/> (accessed on 29 February 2024).
15. Yao, P.; Sun, L.; Lu, L.; Ding, H.; Chen, X.; Tang, L.; Xu, X.; Liu, G.; Hu, Y.; Ma, Y.; et al. Effects of Genetic and Nongenetic Factors on Total and Bioavailable 25(OH)D Responses to Vitamin D Supplementation. *J. Clin. Endocrinol. Metab.* 2017, 102, 100–110, doi:10.1210/jc.2016-2930.
16. Mazahery, H.; von Hurst, P.R. Factors Affecting 25-Hydroxyvitamin D Concentration in Response to Vitamin D Supplementation. *Nutrients* 2015, 7, 5111–5142, doi:10.3390/nu7075111.
17. Hu, Z.; Tao, S.; Liu, H.; Pan, G.; Li, B.; Zhang, Z. The Association between Polymorphisms of Vitamin D Metabolic-Related Genes and Vitamin D3 Supplementation in Type 2 Diabetic Patients. *J. Diabetes Res.* 2019, 2019, e8289741, doi:10.1155/2019/8289741.
18. Xia, Z.; Hu, Y.; Han, Z.; Gao, Y.; Bai, J.; He, Y.; Zhao, H.; Zhang, H. Association of Vitamin D Receptor Gene Polymorphisms with Diabetic Dyslipidemia in the Elderly Male Population in North China. *Clin. Interv. Aging* 2017, 12, 1673–1679, doi:10.2147/CIA.S145700.
19. Cavalcante, I.G. de M.; Silva, A.S.; Costa, M.J.C.; Persuhn, D.C.; Issa, C.I.; Freire, T.L. de L.; Gonçalves, M. da C.R. Effect of Vitamin D3 Supplementation and Influence of Bsm1 Polymorphism of the VDR Gene of the Inflammatory Profile and Oxidative Stress in Elderly Women with Vitamin D Insufficiency: Vitamin D3 Megadose Reduces Inflammatory Markers. *Exp. Gerontol.* 2015, 66, 10–16, doi:10.1016/j.exger.2015.03.011.
20. AlFaqih, M.A. Association of Vitamin D Levels and Polymorphisms in Vitamin D Receptor with Type 2 Diabetes Mellitus Available online: <https://www.spandidos-publications.com/10.3892/br.2022.1585> (accessed on 29 February 2024).
21. American Diabetes Association Standards of Medical Care in Diabetes-2022 Abridged for Primary Care Providers. *Clin. Diabetes Publ. Am. Diabetes Assoc.* 2022, 40, 10–38, doi:10.2337/cd22-as01.
22. Holick, M.F. The Vitamin D Deficiency Pandemic: Approaches for Diagnosis, Treatment and Prevention. *Rev. Endocr. Metab. Disord.* 2017, 18, 153–165, doi:10.1007/s11154-017-9424-1.
23. Nimitphong, H.; Holick, M.F. Vitamin D Status and Sun Exposure in Southeast Asia. *Dermatoendocrinol.* 2013, 5, 34–37, doi:10.4161/derm.24054.
24. Sung, C.-C.; Liao, M.-T.; Lu, W.; Wu, C.-C. Role of Vitamin D in Insulin Resistance. *J. Biomed. Biotechnol.* 2012, 2012, 634195, doi:10.1155/2012/634195.
25. Li, X.; Liu, Y.; Zheng, Y.; Wang, P.; Zhang, Y. The Effect of Vitamin D Supplementation on Glycemic Control in Type 2 Diabetes Patients: A Systematic Review and Meta-Analysis. *Nutrients* 2018, 10, 375, doi:10.3390/nu10030375.
26. Seida, J.C.; Mitri, J.; Colmers, I.N.; Majumdar,

- S.R.; Davidson, M.B.; Edwards, A.L.; Hanley, D.A.; Pittas, A.G.; Tjosvold, L.; Johnson, J.A. Clinical Review: Effect of Vitamin D3 Supplementation on Improving Glucose Homeostasis and Preventing Diabetes: A Systematic Review and Meta-Analysis. *J. Clin. Endocrinol. Metab.* 2014, 99, 3551–3560, doi:10.1210/jc.2014-2136.
27. George, P.S.; Pearson, E.R.; Witham, M.D. Effect of Vitamin D Supplementation on Glycaemic Control and Insulin Resistance: A Systematic Review and Meta-Analysis. *Diabet. Med.* 2012, 29, e142–e150, doi:10.1111/j.1464-5491.2012.03672.x.
28. Is Micro Evolution in Tropical Country Women Resulting Low 25(OH) Available online: <https://www.longdom.org/open-access/is-micro-evolution-in-tropical-country-women-resulting-low-25ohd-level-a-cross-sectional-study-in-indonesia-33467.html> (accessed on 29 February 2024).
29. Heaney, R.P. Vitamin D in Health and Disease. *Clin. J. Am. Soc. Nephrol. CJASN* 2008, 3, 1535–1541, doi:10.2215/CJN.01160308.
30. Institute of Medicine (US) Committee to Review Dietary Reference Intakes for Vitamin D and Calcium Dietary Reference Intakes for Calcium and Vitamin D; Ross, A.C., Taylor, C.L., Yaktine, A.L., Del Valle, H.B., Eds.; The National Academies Collection: Reports funded by National Institutes of Health; National Academies Press (US): Washington (DC), 2011;
31. Al-Daghri, N.M.; Al-Attas, O.S.; Alkharfy, K.M.; Khan, N.; Mohammed, A.K.; Vinodson, B.; Ansari, M.G.A.; Alenad, A.; Alokail, M.S. Association of VDR-Gene Variants with Factors Related to the Metabolic Syndrome, Type 2 Diabetes and Vitamin D Deficiency. *Gene* 2014, 542, 129–133, doi:10.1016/j.gene.2014.03.044.
32. Usategui-Martín, R.; De Luis-Román, D.-A.; Fernández-Gómez, J.M.; Ruiz-Mambrilla, M.; Pérez-Castrillón, J.-L. Vitamin D Receptor (VDR) Gene Polymorphisms Modify the Response to Vitamin D Supplementation: A Systematic Review and Meta-Analysis. *Nutrients* 2022, 14, 360, doi:10.3390/nu14020360.
33. Ross, A.C.; Manson, J.E.; Abrams, S.A.; Aloia, J.F.; Brannon, P.M.; Clinton, S.K.; Durazo-Arvizu, R.A.; Gallagher, J.C.; Gallo, R.L.; Jones, G.; et al. The 2011 Report on Dietary Reference Intakes for Calcium and Vitamin D from the Institute of Medicine: What Clinicians Need to Know. *J. Clin. Endocrinol. Metab.* 2011, 96, 53–58, doi:10.1210/jc.2010-2704.
34. Alvina, A.; Immanuel, S.; Harbuwono, D.S.; Suyatna, F.D.; Harahap, A.; Prihartono, J.; Pusparini, P. Effect of Three and Six Months of Vitamin D Supplementation on Glycemic Control and Insulin Resistance in Type 2 Diabetes Mellitus: Randomized Placebo-Controlled Trial. *Indones. Biomed. J.* 2023, 15, 287–295, doi:10.18585/inabj.v15i3.2370.

supp vit D

by joice kalumpiu

Submission date: 11-Oct-2024 10:28AM (UTC+0700)

Submission ID: 2481822254

File name: ajol-file-journals_595_articles_273830_66a107c7a8655.pdf (403.85K)

Word count: 6985

Character count: 35341

Supplementary vitamin D3 and vitamin D receptor polymorphisms affect blood vitamin D levels in type-2 diabetes mellitus in Indonesia

Authors: Yenny^{1,*}; R. Wratsangka¹; E. Herwana¹; J. V. Kalumpiu¹; P. B. Liman¹

Affiliation: ¹Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

ABSTRACT

INTRODUCTION: There are no data on vitamin D receptor (VDR) gene single nucleotide polymorphism (SNP) influence on blood 25-hydroxy-cholecalciferol [25(OH)D] levels after supplementary vitamin D in Indonesian type 2 diabetes mellitus (T2DM) patients. This study evaluated the effects of the supplementary vitamin D3 and VDR gene SNPs rs1555410 and rs2228570 on blood 25(OH)D levels in T2DM cases.

METHODS: A randomized, double-blind, placebo-controlled trial (RDPCT) was conducted at one research setting using T2DM subjects divided into vitamin D group (VDG) and control group (CG) and receiving 5,000 IU/day vitamin D3 (cholecalciferol) or placebo once daily for 84 days. Levels of 25(OH)D were determined baseline and after supplementary vitamin D3 administration for 84 days. Circulatory 25(OH)D was assayed using ELISA. VDR polymorphisms were detected using sequencing.

RESULTS: Post-supplementary blood 25(OH)D rose appreciably from baseline in VDG for VDR rs1555410 genotypes G/G ($p=0.001$) and G/A ($p=0.010$), and in VDR rs2228570 genotypes T/T ($p=0.012$), T/C ($p<0.001$), and C/C ($p=0.001$). Post-supplementary VDG still contained 30.3% of subjects not reaching blood 25(OH)D ≥ 30 ng/mL.

In attaining blood 25(OH)D ≥ 30 ng/mL post-supplementation, VDR rs2228570 genotype T/C differed significantly from T/T (52.4% v. 100%; $p=0.027$), but there were no appreciable differences between genotypes C/C and T/T (78.6% v. 100%; $p=0.273$), as well as between VDR rs1555410 genotypes G/G and G/A (67.5% v. 100%; $p=0.542$).

CONCLUSION: Only 52.4% of subjects with VDR rs2228570 genotype T/C achieved sufficiently high blood 25(OH)D levels. VDR rs2228570 polymorphisms apparently influence T2DM response to supplementary vitamin D.

Keywords: Diabetes Mellitus, Vitamin D, Single Nucleotide Polymorphism, Indonesia

INTRODUCTION

Indonesia, with 10.7 million type 2 diabetes mellitus (T2DM) patients, occupies the 7th global

rank, with predicted T2DM prevalence rising to 16.6 million in 2045 [1]. T2DM vitamin D deficiency prevalence is higher than in the general global population, with prevalences of 63.2- 83.2% [2,3].

***Corresponding author:** Yenny, Department of Pharmacology and Clinical Pharmacy, Faculty of Medicine, Universitas Trisakti, Jakarta post code 11440, Indonesia; Email: yennyfarmako@trisakti.ac.id; Orcid id: 0000-0001-9390-5527; **Potential Conflicts of Interest:** All authors: no potential conflicts of interest disclosed; **Funding:** All authors: Research funding from Universitas Trisakti No. 0303/PUF/FK/2021-2022; **Academic integrity:** All authors confirm that they have made substantial academic contributions to this manuscript as defined by the ICMJE; **Ethics of human subject participation:** The study was approved by the local Institutional Review Board. **Informed consent** was sought and gained where applicable; **Originality:** All authors: this manuscript is original has not been published elsewhere; **Review:** This manuscript was peer-reviewed by three reviewers in a double-blind review process; **Type-editor:** Peter (USA).

Received: 10 February 2024; **Initial decision given:** 14th April 2024; **Revised manuscript received:** 14th April 2024; **Accepted:** 27th May 2024.

Copyright: © The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND) (click here) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Biomedical Centre (RBC)/Rwanda Health Communication Center, P. O. Box 4586, Kigali. ISSN: 2079-097X (print); 2410-8626 (online)

Citation for this article: Yenny; R. Wratsangka; E. Herwana et al. Supplementary vitamin D3 and vitamin D receptor polymorphisms affect blood vitamin D levels in type-2 diabetes mellitus in Indonesia. Rwanda Medical Journal, Vol. 81, no. 2, p. 36-46, 2024. <https://dx.doi.org/10.4314/rmj.v81i2.5>

The latter are higher than in Europe, with a vitamin D deficiency prevalence of 40.4% (blood 25(OH)D <20 ng/mL) [4], while vitamin D deficiency prevalence in South Asia is 68% [5].

Vitamin D supplementation raises serum 25(OH)D concentration, influences health outcomes, and achieves maximum mortality rate reductions in developed countries from the commonest fatal diseases, such as cardiovascular disease and T2DM [6]. Gross vitamin D deficiency at 25(OH)D <12 ng/mL (or <30 nmol/L) is a health hazard [7] and should be corrected through vitamin D supplementation [8].

The utility of providing additional vitamin D to T2DM patients for glucose hemostasis and reducing insulin resistance is still debated, presumably because of the diversity of study populations for serum vitamin D status, supplementation dose and duration, methodology, gender, BMI, and ethnicity [9,10].

Vitamin ²D regulates signal transmission through the vitamin D receptor (VDR) [11]. The vitamin D-responsive element (VDRE) gene on chromosome 12q13.1 comprises nine exons and eight introns [11]. VDR activates and controls gene transcription via the target gene promoter VDRE. The VDR gene has over 470 polymorphisms [11], the most common being FokI (rs2228570 C to T) and BsmI (rs1544410 A to G) [11]. VDR rs1544410 at intron 8 regulates mRNA stability, thereby affecting gene expression [12]. VDR rs2228570 lies in the exon 2 start codon and changes the initiation sites [13]. VDR gene SNPs may influence VDR mRNA and protein stability and activity, resulting in a suboptimal response to supplementary vitamin D [14,15].

Multiple factors may affect post-supplementary blood 25(OH)D, such as sunshine exposure, aging, body mass index, calcium intake, supplementary vitamin D, and genetics [4,16]. It is currently uncertain whether VDR SNPs affect blood vitamin D levels after supplementary vitamin D in T2DM patients [17], due to difficulties in detecting gene involvement in the development of T2DM, because of possible minute differences in the gene and its interaction with genetic or non-genetic factors [18]. VDR SNP influence on increases in blood 25(OH)D levels that depict ⁴⁰min D condition, can be evaluated only with a randomized, double-blind placebo-controlled trial (RDBPCT).

Currently, few randomized double-blind placebo-controlled trials (RDBPCT) exist for evaluating

VDR rs1544410 and rs2228570 relationships with vitamin D therapeutic response. Waterhouse et al. [14]. Studied Australian elderly aged 60 – 80 years receiving vitamin D3 supplementation (30.000 vs. 60.000 IU/month) ¹³ for 12 months and found rs2228570 was not associated with increases in 25(OH)D levels after supplementary vitamin D. A 20-week Chinese RDBPCT [15] on vitamin D deficiency cases receiving vitamin D3 at 2000 IU/day or placebo, demonstrated that rs2228570 showed stronger influences on 25(OH)D levels (p<0.04). Post-treatment, ²² VDR rs2228570-G, and its alleles had higher 25(OH)D levels (p = 0.009).

The RDBPCT ¹⁷ of Cavalcante et al. [19]. on elderly females aged 68 ± 6 years with vitamin D insufficiency and receiving 200.000 IU ²⁹ vitamin D3 supplementation for 4 weeks, showed higher blood 25(OH)D concentrations in persons with BsmI genotypes BB/Bb (p<0.001) who responded better to supplementary vitamin D than those with bb genotype.

A prospective case-control study involving 125 T2DM patients and 125 controls, revealed that low blood 25(OH)D and gene rs2228570 correlated with T2DM risk [20]. The inconsistent study results on VDR SNP relationships were caused by variations in pre-supplementary blood vitamin D levels, study designs, small sample sizes, vitamin D dose, duration of supplementary vitamin D, dietary intake, and ethnicity.

There are currently no reports on the causal relationship of VDR SNPs rs1544410 and rs2228570 with supplementary vitamin D among Indonesian T2DM patients, for which an RDBPCT is necessary. The outcome may be useful for vitamin D3 therapeutic dose personalization in T2DM patients to reduce morbidity and mortality rates. Primary research outcomes would be responses to supplementary vitamin D by comparing post-supplementary blood 25(OH)D values. Secondary outcomes would be the impact of rs1544410 and rs2228570 polymorphisms on blood 25(OH)D changes and attainment of blood 25(OH)D of ≥30 ng/mL

METHODS

Research design

This RDBPCT was conducted from June to August 2022 at Puskesmas Mampang in South ⁸ Jakarta, with subjects signing informed consent. Study protocol approval was by the Research Ethics Committee,

Faculty of Medicine, Universitas Trisakti, under No. 001/KER/FK/1/2022.

Patients and intervention

The study subjects were Mampang area residents with T2DM. Inclusion criteria: males and females ≥ 18 years old, T2DM, HbA1c 7-8.5%, oral antidiabetic drug monotherapy, agreeing to follow-up controls. T2DM was diagnosed using American Diabetes Association criteria [21], namely fasting blood glucose ≥ 126 mg/dL, or 2-hour postprandial blood glucose ≥ 200 mg/dL and HbA1c $\geq 6.5\%$. Exclusion criteria: previously and currently on insulin therapy, suspect kidney disease (estimated glomerular filtration rate < 30 mL/min/1.73m²), abnormal liver function (SGPT 3 times normal upper limit), pregnant or lactating, allergy, hypercalcemia (plasma calcium > 2.65 mmol/L), or receiving daily supplementary vitamin D in preceding 84 days. Dropout criteria: blood 25(OH) D > 100 ng/mL, hypercalcemia, cholecalciferol hypersensitivity.

The study enrolled 115 subjects allocated by simple randomization to vitamin D (intervention) and control (placebo) groups at 1:1 ratio, which was conducted by personnel blinded to the intervention. VDG received daily vitamin D3 tablets containing 5,000 IU cholecalciferol, whereas controls received once-daily placebo tablets (120mg calcium), all tablets taken for 84 days. The vitamin D and placebo tablets contained in dark-colored glass bottles coded A and B were identical in visual, olfactory, and gustatory qualities. Participants and the statistician were all blinded to the origin of the tablets, whether from VDG or CG. Compliance with the intervention was determined by weekly counting the remaining tablets. VDG had 6 dropouts because of protocol non-compliance, returning to the home village, and diarrhea, while CG had 7 dropouts because of not agreeing to participate, protocol non-compliance, and nausea and vomiting. Subjects completing the study were 85 in number, consisting of 43 in VDG and 42 in CG (Figure 1).

The researchers were blinded to subject allocation in all study phases (recruitment, enrolment, data collection, and group assignment). For improving verification and compliance, all empty medication bottles were returned monthly to the cadres for evaluation of subject compliance to supplementation at the completion of the study. Subjects' complaints (potential adverse events)

were noted for recording. The principal investigator evaluated the complaints and their connection with the supplements. The development of all reported symptoms was monitored until the study was completed. The allocation codes stored by an independent third party were opened after the study was completed.

Measurements

On day zero (admission date), before administration of vitamin D tablets, subjects meeting recruitment criteria were interviewed to collect subject data on age, gender, and duration of diabetes mellitus, followed by blood collection at 08.00 and 9.00 a.m. local time.

Biochemical measurements

From each participant, 5 mL of venous whole blood was collected, with 3 mL for blood 25(OH) D determination and 2 mL in EDTA tubes for VDR SNP genotyping.

Blood 25(OH)D level was determined by chemiluminescent microparticle immunoassay (ARCHITECT 25-OH Vitamin D assay, Abbott), with measuring interval 8.0- 160.0 ng/mL (20.0- 400.0 nmol/L), limit of detection (LoD) ≤ 10.0 ng/mL, limit of quantitation (LoQ) ≤ 20 ng/mL, and imprecision $\leq 10\%$ within total CV.

Blood 25(OH)D data were presented as median \pm SD, and categorized according to Endocrine Society 2011 clinical practice guideline, with < 20 ng/mL defined as vitamin D deficiency, 20 – 30 ng/mL as vitamin D insufficiency, 30 – 100 ng/mL as vitamin D sufficiency [8].

Vitamin D receptor single nucleotide polymorphisms

Genomic DNA was extracted using QIAamp DNA Blood Mini Kit (QIAGEN) from 200 μ L EDTA blood and its purity and level were determined using a NanoDrop 2000 spectrophotometer (ThermoScientific). VDR SNPs were detected by PCR, followed by sequencing. PCR used MyTaq HS Red Mix Kit (Bioline) to amplify the target sequence using specific primers, namely for rs1544410: forward primer 5'- GGG AGT ATG AAG GAC AAA GAC C-3' and reverse primer 5'- CCC GCA AGA AAC CTC AAA TAA C-3' and for rs2228570: forward primer 5'-GCA CTG ACT CTG GCT CTG-3' and reverse primer 5'-TGG ACA TTG TAA GGA AGG AGA TG-3'. PCR amplification used 2 μ L DNA, 8.5 μ L nuclease-free water, 2x 12.5 μ L My Taq HS Red Mix,

and 1.0 μ l each of forward and reverse primers. Conditions: initial activation at 95oC for 5 minutes, denaturation at 95oC for 15 seconds, annealing at 58oC for 30 seconds, extension at 72oC for 30 seconds for 40 cycles, and final extension at 72oC for 3 minutes. PCR products on 2% agarose gel were visualized by electrophoresis. Purified PCR products were sequenced using BigDye Terminator Kit (Thermo Fisher Scientific) and ABI 3500 sequencer (Applied Biosystems). Sequence analysis was performed with BioEdit software to confirm mutations, which were compared to NCBI BLAST

database with accession number NG_008731.1.

Sample size

In each group, 40 subjects were required to detect a treatment difference at 5% two-sided significance level (0.05) and 90% study power. $\mu_1 - \mu_2$ = predicted between-group mean difference, estimated at 4; σ^2 = expected population variance from the preliminary study, estimated at 1.02. To anticipate dropouts, the sample size became 95 for adequate power to detect outcome measure differences.




Figure 1: Flow of participants throughout trial

Statistical analysis

Continuous data of normal distribution were shown as mean and standard deviation (SD), continuous data of skewed distribution as median (minimum-maximum), and categorical data as percentages. Data were checked for normal distribution using the Kolmogorov-Smirnov test. Unpaired Student's t-test, Mann-Whitney U test, Chi-square test, and Fisher exact test were used for baseline comparisons between VDG and CG based on the variable type and data distribution. Mann-Whitney test was used to compare blood 25(OH)D in VDG and CG at baseline and post-supplementation. Baseline and post-supplementation vitamin D status differences were evaluated using Fisher's exact test and Chi-square test. VDG baseline and post-supplementation blood 25(OH)D differences between rs1544410 genotypes were evaluated by the Mann-Whitney test. VDG baseline and post-supplementation blood 25(OH)D levels between rs2228570 genotypes were compared using the Kruskal-Wallis test, Wilcoxon signed rank test, and paired t-test at $p < 0.05$. Statistical analysis used SPSS for Windows version 23.

RESULTS

Baseline subject characteristics

We had 85 T2DM participants, with 68 (80%) females, mean age 55.8 ± 0.6 years, median T2DM duration 12 (1 – 36) months, and median blood 23(OH)D 11.6 (2.4 – 30.3) ng/mL. The most frequent rs1555410 genotype was G/G, comprising 79 subjects (92.9%), followed by G/A with 6 subjects (7.1%), without any T/T. Gene rs2228570 had genotypes T/C, C/C, and T/T with 38 (44.7%), 32 (37.6%), and 15 (17.6%) subjects, respectively. Vitamin D status was mostly deficient in 74 subjects (84.1%), insufficiency in 9 subjects (10.6%), and sufficiency 2 subjects (2.4%). After group randomization, no significant differences were observed in age, gender, T2DM duration, and VDR genotype between VDG and CG (Table 1).

Baseline and post-supplementation blood 25(OH)D and vitamin D status

After vitamin D3 supplementation for 84 days, there was a much greater increase in VDG blood 25(OH)D than in CG (46(9.4-79.4) v. 14.4(6.9-38.3); ($p < 0.001$) (Table 2).

Table 1: Subject characteristics at the start of the study in VDG and CG

Characteristic	Treatment group		P-value
	VDG (n=43)	CG (n=42)	
Age (years)	56 (35 – 80)	56 (35 – 69)	0.396 ^a
Gender			
Male	8 (47.1)	9 (52.9)	0.745 ^b
Female	35 (51.5)	33 (48.5)	
Duration of DM (months)	12 (1 – 36)	12 (1 – 36)	0.967 ^a
VDR genotype (n,%)			
rs1544410			
G/G	40 (50.6)	39 (49.4)	1.000 ^c
G/A	3 (50)	3 (50)	
A/A	0 (0)	0 (0)	
rs 2228570			
T/T	8 (53.3)	7 (46.7)	0.614 ^b
T/C	21 (55.3)	17 (44.7)	
C/C	14 (43.8)	18 (56.2)	

Values presented as median (min-max) or n(%). Statistical analysis: ^aMann-Whitney test; ^bChi-square test; ^cFisher's exact test; $p < 0.05$ = statistically significant. VDG = vitamin D group; CG = control group

Table 2: Blood 25 (OH)D level and vitamin D status in VDG and CG at baseline versus 84 days after supplementation with vitamin D3.

Group	Start of study		p value	After supplementation		p-value
	VDG (n=43)	CG (n=42)		VDG (n=43)	CG (n=42)	
Blood 25(OH)D level (ng/mL)						
	10.5	13.05	0.264 ^a	46	14.4	0.001 ^{a*}
	(4.7 – 30.3)	(2.4 – 26.9)		(9.4 – 79.4)	(6.9 – 38.3)	
Vitamin D status						
Deficiency	40 (50.6)	34 (45.9)		3 (8.6)	32 (91.4)	
Insufficiency	2 (22.2)	7 (77.8)	1.000 ^c	10 (62.5)	6 (37.5)	0.001 ^{b*}
Sufficiency	1 (50)	1 (50)		30 (88.2)	4 (11.8)	

Groups: VDG= vitamin 25,000IU/day, CG = placebo

Vitamin D status (blood 25(OH)D level): deficiency (< 20 ng/mL); insufficiency (20 – 30ng/mL); sufficiency (30 – 100 ng/mL). Statistical analysis: aMann-Whitney test; bChi-square test; cFisher's exact test; *p<0.05 =statistically significant

VDG = vitamin D group; CG = control group

Comparison of blood 25(OH)D levels by VDR genotype after supplementary vitamin D3 administration in VDG

To find the effect of VDR genotype on post-supplementation blood 25(OH)D, 25(OH)D blood levels were compared between genotypes in VDG (Table 3). Blood 25(OH)D levels increased significantly above baseline after supplementary vitamin D3 in rs1544410 genotypes G/G [10.5 (4.7

– 30.5) v. 46.5 (9.4 – 79.4) ng/mL; p=0.001] and G/A [10.8 ± 1.3 v. 45.7 ± 7.2 ng/mL; p =0.010]. No great differences were found in 25(OH)D between rs1544410 genotypes G/G and G/A [46.5 (9.4 – 79.4) ng/mL v. 45.7 ± 7.2 ng/mL; p=0.924].

As compared to baseline, post-supplementation rs2228570 blood 25(OH)D levels increased significantly for genotypes T/T [11.4 (6.2 – 17.9) v.

Table 3: Blood 25(OH)D levels in VDG by VDR genotype after supplementary vitamin D3 administration

VDG	Blood 25(OH)D level (ng/mL)		p-value
	Start of study (n=43)	84 days (n=42)	
VDR genotypes			
rs1544410			
G/G (n= 40)	10.5 (4.7 – 30.5)	46.5 (9.4 – 79.4)	0.001 ^{c*}
G/A (n= 3)	10.8 ± 1.3	45.7 ± 7.2	0.010 ^{d*}
P value	0.924 ^a	0.924 ^a	
rs2228570			
T/T (n=8)	11.4 (6.2 – 17.9)	61.2 (32.1 – 79.4)	0.012 ^{c*}
T/C (n=21)	10.5 (7.6 – 30.3)	34.2 (9.4 – 67.4)	<0.001 ^{c*}
C/C (n=14)	11.0 ± 4.4	43.7 ± 12.4	0.001 ^{d*}
P value	0.373 ^b	0.024 ^{b*}	

Statistical analysis: aMann-Whitney test; bKruskal-Wallis test; cWilcoxon signed rank test; dpaired t-test; *p value <0.05 = statistically significant

VDG = vitamin D group; CG = control group

Table 4: Attainment of blood 25(OH)D levels in VDG by VDR genotype after supplementary vitamin D administration

Administration		15	
	Blood 25(OH)D level		p-value
VDR genotype	< 30 ng/mL (n,%)	≥30 ng/mL (n,%)	
VDG			
rs 1544410			
G/G (n=40)	13 (32.5)	27 (67.5)	0.542
G/A (n=3)	0 (0)	3 (100)	
rs 2228570			
T/T (n=8)	0 (0)	8 (100)	0.027*
T/C (n= 21)	10 (47.6)	11 (52.4)	
C/C (n=14)	3 (21.4)	11 (78.6)	0.273

Statistical analysis: logistic regression test; *p value <0.05 = statistically significant

VDG = vitamin D group

61.2 (32.1 – 79.4); $p=0.012$], T/C [10.5 (7.6 – 30.3) v. 34.2 (9.4 – 67.4); $p<0.001$], and C/C [11.0 ± 4.4 v. 43.7 ± 12.4; $p=0.001$] (Table 3).

There were also significant differences in post-supplementation blood 25(OH)D between rs2228570 genotypes T/T, T/C, and C/C themselves [61.2 (32.1 – 79.4) ng/mL v. 34.2 (9.4 – 67.4) ng/mL v. 43.7 ± 12.4 ng/mL; $p=0.024$] (Table 3).

To determine the genotypes causing the post-supplementation differences in rs2228570, a post-hoc analysis was conducted, showing significant differences between genotypes T/T and T/C ($p=0.015$) and between T/T and C/C ($p=0.017$), but not between T/C and C/C ($p=0.391$).

Regarding blood 25(OH)D responses to supplementary vitamin D of rs1544410 genotypes G/G and G/A, among the 42 VDG subjects, only 30 (69.7%) attained blood 25 (OH)D levels ≥30 ng/mL. The same is true for the three rs2228570 genotypes T/T, T/C, and C/C (Table 4).

No prominent differences were found between rs1544410 G/G and G/A in attaining blood 25(OH)D ≥30 ng/mL (67.5% v. 100%; $p=0.542$). In rs2228570, significant differences occurred in blood 25(OH)D level attainment between T/C and T/T (52.4% v. 100%; $p=0.027$).

DISCUSSION

Vitamin D deficiency was found in 84.1% of subjects, with a median blood 23(OH)D level of

11.6 (2.4 – 30.3) ng/mL, which is much greater than in Europe, where 25(OH)D levels below 20 ng/mL and 12 ng/mL are observed in 40.4% and 13.0%, respectively, of the population [4]. Conversely, adult vitamin D deficiency prevalence in 5 South Asian countries was 68% [5]. Our results approximate those of earlier studies demonstrating that T2DM vitamin D deficiency prevalence is around 63.2 – 83.2% [2,3], and that vitamin D deficiency is also found in tropical countries, such as Indonesia, with abundant sunshine for cutaneous vitamin D synthesis.

Vitamin D as a prohormone is available as vitamin D2 (ergocalciferol) in foods and vitamin D3 (cholecalciferol) in UV-exposed human skin. Generally, vitamin D deficiency is caused by low dietary intake and reduced cutaneous synthesis from inadequate sunlight exposure due to geographic location, skin color, age, indoor lifestyle, and cultural or religious practices[22,23]. Our greater vitamin D deficiency prevalence shows that foods and sunlight alone cannot maintain optimal vitamin D status, necessitating vitamin D supplementation.

Genal transmission in humans occurs through vitamin D binding with the vitamin D receptor (VDR) [11] expressed by insulin-sensitive tissues. Apparently, vitamin D may have a direct influence on insulin sensitivity and insulin receptor expression, thereby enhancing insulin-stimulated glucose transport. Vitamin D may also have an indirect influence [17] by reducing insulin resistance-inducing inflammatory responses [24]. A meta-analysis showed that vitamin D

supplements can raise blood 25(OH)D and reduce insulin resistance in T2DM [25]. However, systematic reviews of T2DM RCTs failed to find evidence for the efficacy of vitamin D supplements in glucose hemostasis and in decreasing insulin resistance [26,27]. According to another meta-analysis, vitamin D dosage, status, and length of supplementation affect the therapeutic response [9]. High-dose vitamin D supplementation produces greater effects in obese vitamin D-deficient Asians [9]. Irrespective of vitamin D supplementation's impact on T2DM patients, vitamin D deficiency should be corrected because vitamin D serves an essential function in calcium hemostasis and bone metabolism [22].

Our study showed that rs1555410 had genotype G/G in 79 (92.9%) subjects and no genotype A/A, whereas Chinese T2DM patients [17] had mostly rs1544410 genotype G/A (93.75%), but similarly no A/A. Our rs2228570 genotype proportions comprised T/C in 44.7%, C/C in 37.6%, and T/T in 17.6% of subjects, whereas the Chinese study [17] comprised rs2228570 genotype C/C in 59.8%, T/T in 23.2%, and C/T in 16.9% of subjects, showing that Asians vary in rs1544410 and rs2228570 gene proportions. Similarly, the research results of Sari et al. [28], on healthy North Sumatran women differed from ours, because all subjects had heterozygous A>G for BsmI (rs1544410), and T>C for TaqI (rs2228570), showing that Indonesians also have differing VDR genotype proportions, supporting the supplementary vitamin D dosage personalization concept.

In our study, supplementary vitamin D at 5000 IU/day caused a 3.2-fold higher rise in blood 25(OH)D in VDG compared to CG [46 (9.4 – 79.4) ng/mL v. 9.4 (6.9 – 38.3) ng/mL; $p=0.001$] (Table 2). Blood 25(OH)D may rise around 1 ng/mL (2.5 nmol/L) per 100 IU daily vitamin D3 supplement given for 56 – 84 days [29], although the supplementary dose may not be linearly correlated with blood 25(OH)D [30].

After 84 days of vitamin D supplementation at 5000 IU/day, among our 43 subjects, 13 (30.2%) subjects still did not attain blood 25(OH)D levels of ≥ 30 ng/mL (Table 4.) This agrees with the findings of Yao et al. [15], from a 140-day RDBPCT on 448 Chinese with vitamin D deficiency receiving 2000 IU/day vitamin D3 or placebo, where the vitamin D increased blood 25(OH)D, but could not overcome vitamin D deficiency in 25% of subjects. Al-Daghari et al. [31]. Studying T2DM patients on 2000 IU/day

supplementary vitamin D for 12 months showed that 42% of subjects still could not reach target blood 25(OH)D. Hu et al. [17]. in their study on T2DM subjects receiving supplementary vitamin D at 800 IU/day for 12 months, also found that 44.6% of subjects did not attain vitamin D sufficiency.

The influence of VDR SNP on supplementary vitamin D3 results remains unclear. We found a significant increase in post-supplementary blood 25(OH)D levels above baseline values in rs1544410 genotypes G/G ($p=0.001$) and G/A ($p=0.010$) which agrees with Cavalcante et al. [19]. showing that supplementary vitamin D significantly increased blood 25(OH)D in BB/Bb ($p=0.009$), but not in bb subjects.

our VDG subjects with rs2228570, large differences were found in post-supplementary 25(OH)D blood concentrations of genotypes T/T, T/C, and C/C ($p=0.024$) (Table 3.). Post-hoc analysis showed differences between T/T and T/C ($p=0.015$) and between T/T and C/C ($p=0.017$), but not between T/C and C/C ($p=0.391$). In our study, rs2228570 apparently influenced supplementary vitamin D response in T2DM subjects. In T/C subjects, only 52.4% attained 25(OH)D ≥ 30 ng/mL, lower than in the other genotypes (Table 4).

One meta-analysis showed that TaqI and FokI polymorphisms may modulate supplementary vitamin D response for better results [32]. Our study confirms that the doses should be adapted ("personalized") in subjects with rs2228570 genotype T/C for optimal benefits of supplementary vitamin D.

Our study results differ from those of Al-Daghari et al. [31]. on T2DM subjects with genotype-related differences in post-supplementary blood 25(OH)D, in that genotype T/T subjects evidenced better therapeutic responses than the other genotypes. Our results also differ from those of Hu et al. [17]. showing in T2DM subjects that rs2228570 genotypes T/C and T/T had no remarkable differences in blood 25(OH)D ($p=0.964$). In addition, Hu's study and ours showed differences in subject characteristics regarding age, baseline blood 25(OH)D, and supplementary vitamin D dose. Hu's subjects were aged 66.3 ± 9.1 years, whereas ours were 55.8 ± 0.6 years old. Our baseline blood 25(OH)D of $11.6 (2.4 – 30.3)$ ng/mL exceeded that of Hu's 22.7 ± 1.9 ng/mL, presumably because Hu et al. used a lower vitamin D3 supplementation dose (800 IU) [17]. Another study found that lower baseline blood 25(OH)D levels were associated

with significantly higher blood 25(OH)D responses [15].

Opinions regarding optimal blood 25(OH)D levels in humans are inconsistent, with no uniform definition of vitamin D deficiency and insufficiency in different guidelines. The IOM recommends a minimum blood 25(OH)D concentration of 20 ng/mL (50 nmol/L), in connection with bone health [7,3]. However, the Endocrine Society recommends 25(OH)D levels exceeding 30 ng/mL (or 75 nmol/L) for preventing infections and obtaining other non-calcemic vitamin D benefits [8]. We showed that supplementary vitamin D3 at 5000 IU/day for 84 days still could not prevent 30% of subjects from attaining vitamin D sufficiency with blood 25(OH)D \geq 30 ng/mL. The Endocrine Society clinical practice guideline [8] recommends supplementary vitamin D3 for increasing vitamin D levels and determining blood 25(OH)D concentrations because 25(OH)D is the most frequent circulatory vitamin D, with a half-life of 19–21 days, and extremely useful for monitoring vitamin D status in persons at high risk of vitamin D deficiency.

Our study confirms the need for supplementary vitamin D dose personalization and blood 25(OH)D measurement in high-risk patients in relation to VDR SNPs, apart from the contradictory relationship of vitamin D deficiency in glucose hemostasis and insulin resistance reduction [20,26,27,34]. There is a need for dosage adjustment (“personalization”) in subjects with rs2228570 genotype T/C to obtain better gains from supplementary vitamin D. Our study results may provide inputs on management policies of T2DM patients susceptible to vitamin D deficiency, particularly in Indonesia. In some populations, the interplay of genes and lifestyle may obscure the genetic component; therefore, studies on gene interactions with diet and physical activity are mandatory to confirm the relationship. Other longer-term RCTs with larger samples are also necessary to better utilize the results of vitamin D supplements in patients with type 1 and type 2 diabetes.

We used an RDBPCT design that is best for measuring cause-and-effect relationships. We used an identical therapeutic procedure and supplementary vitamin D3 dosage to minimize subject variation.

One limitation of this study was that our subjects were Indonesian T2DM patients; therefore, our results may not apply to other nations. We also

did not account for physical activity, diet, sunlight exposure, BMI, and parathyroid hormone as confounders.

CONCLUSION

After vitamin D supplementation, blood 25(OH)D levels rose perceptibly, but a third of subjects still failed to attain blood 25(OH)D levels of \geq 30 ng/mL. VDR rs2228570 genotype T/C had only 52.4% of its subjects attaining a sufficiently large 25(OH)D level, but perceptibly lower than in genotypes T/T and C/C. VDR rs2228570 polymorphisms apparently influence T2DM response to supplementary vitamin D. There is a need for personalization of vitamin D dosage and blood 25(OH)D measurement in high-risk patients due to VDR SNPs.

REFERENCES

1. IDF Diabetes Atlas; 9th ed.; International Diabetes Federation, 2019; ISBN 978-2-930229-87-4. Available from: https://diabetesatlas.org/upload/resources/material/20200302_133351_IDFATLAS9e-final-web.pdf
2. Khudayar, M.; Nadeem, A.; Lodi, M.N.; Rehman, K.; Jawaid, S.I.; Mehboob, A.; Aleem, A.S.; Mirza, R.E.F.; Ahmed, M.; Abbas, K. The Association Between Deficiency of Vitamin D and Diabetes Mellitus Type 2 (DMT2). *Cureus* 2022, 14, e22221, doi:10.7759/cureus.22221.
3. Ac, A. Serum Vitamin D Levels in Persons with Type 2 Diabetes Mellitus in Lagos, Nigeria., doi:10.23937/2377-3634/1410133.
4. Cashman, K.D.; Dowling, K.G.; Škrabáková, Z.; Gonzalez-Gross, M.; Valtueña, J.; De Henauw, S.; Moreno, L.; Damsgaard, C.T.; Michaelsen, K.F.; Mølgaard, C.; et al. Vitamin D Deficiency in Europe: Pandemic? *Am. J. Clin. Nutr.* 2016, 103, 1033–1044, doi:10.3945/ajcn.115.120873.
5. Siddiquee, M.H.; Bhattacharjee, B.; Siddiqi, U.R.; MeshbahirRahman, M. High Prevalence of Vitamin D Deficiency among the South Asian Adults: A Systematic Review and Meta-Analysis. *BMC Public Health* 2021, 21, 1823, doi:10.1186/s12889-021-11888-1.
6. Nutrients | Free Full-Text | A Narrative Review of the Evidence for Variations in Serum 25-Hydroxyvitamin D Concentration Thresholds for Optimal Health Available online: <https://www.mdpi.com/2072-6643/14/3/639> (accessed on 29 February 2024).

7. Vitamin D Deficiency 2.0: An Update on the Current Status Worldwide | European Journal of Clinical Nutrition Available online: <https://www.nature.com/articles/s41430-020-0558-y> (accessed on 29 February 2024).
8. Holick, M.F.; Binkley, N.C.; Bischoff-Ferrari, H.A.; Gordon, C.M.; Hanley, D.A.; Heaney, R.P.; Murad, M.H.; Weaver, C.M.; Endocrine Society Evaluation, Treatment, and Prevention of Vitamin D Deficiency: An Endocrine Society Clinical Practice Guideline. *J. Clin. Endocrinol. Metab.* 2011, 96, 1911–1930, doi:10.1210/jc.2011-0385.
9. Farahmand, M.A.; Daneshzad, E.; Fung, T.T.; Zahidi, F.; Mohammadi, M.; Bellissimo, N.; Azadbakht, L. What Is the Impact of Vitamin D Supplementation on Glycemic Control in People with Type-2 Diabetes: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *BMC Endocr. Disord.* 2023, 23, 15, doi:10.1186/s12902-022-01209-x.
10. Musazadeh, V.; Kavyani, Z.; Mirhosseini, N.; Dehghan, P.; Vajdi, M. Effect of Vitamin D Supplementation on Type 2 Diabetes Biomarkers: An Umbrella of Interventional Meta-Analyses. *Diabetol. Metab. Syndr.* 2023, 15, 76, doi:10.1186/s13098-023-01010-3.
11. Molecular Epidemiology of Vitamin D Receptor Gene Variants | Epidemiologic Reviews | Oxford Academic Available online: <https://academic.oup.com/epirev/article/22/2/203/456955> (accessed on 29 February 2024).
12. Decker, C.J.; Parker, R. Diversity of Cytoplasmic Functions for the 3' Untranslated Region of Eukaryotic Transcripts. *Curr. Opin. Cell Biol.* 1995, 7, 386–392, doi:10.1016/0955-0674(95)80094-8.
13. Uitterlinden, A.G.; Fang, Y.; Van Meurs, J.B.J.; Pols, H.A.P.; Van Leeuwen, J.P.T.M. Genetics and Biology of Vitamin D Receptor Polymorphisms. *Gene* 2004, 338, 143–156, doi:10.1016/j.gene.2004.05.014.
14. Environmental, Personal, and Genetic Determinants of Response to Vitamin D Supplementation in Older Adults - PubMed Available online: <https://pubmed.ncbi.nlm.nih.gov/24694335/> (accessed on 29 February 2024).
15. Yao, P.; Sun, L.; Lu, L.; Ding, H.; Chen, X.; Tang, L.; Xu, X.; Liu, G.; Hu, Y.; Ma, Y.; et al. Effects of Genetic and Nongenetic Factors on Total and Bioavailable 25(OH)D Responses to Vitamin D Supplementation. *J. Clin. Endocrinol. Metab.* 2017, 102, 100–110, doi:10.1210/jc.2016-2930.
16. Mazahery, H.; von Hurst, P.R. Factors Affecting 25-Hydroxyvitamin D Concentration in Response to Vitamin D Supplementation. *Nutrients* 2015, 7, 5111–5142, doi:10.3390/nu7075111.
17. Hu, Z.; Tao, S.; Liu, H.; Pan, G.; Li, B.; Zhang, Z. The Association between Polymorphisms of Vitamin D Metabolic-Related Genes and Vitamin D3 Supplementation in Type 2 Diabetic Patients. *J. Diabetes Res.* 2019, 2019, e8289741, doi:10.1155/2019/8289741.
18. Xia, Z.; Hu, Y.; Han, Z.; Gao, Y.; Bai, J.; He, Y.; Zhao, H.; Zhang, H. Association of Vitamin D Receptor Gene Polymorphisms with Diabetic Dyslipidemia in the Elderly Male Population in North China. *Clin. Interv. Aging* 2017, 12, 1673–1679, doi:10.2147/CIA.S145700.
19. Cavalcante, I.G. de M.; Silva, A.S.; Costa, M.J.C.; Persuhn, D.C.; Issa, C.I.; Freire, T.L. de L.; Gonçalves, M. da C.R. Effect of Vitamin D3 Supplementation and Influence of Bsm1 Polymorphism of the VDR Gene of the Inflammatory Profile and Oxidative Stress in Elderly Women with Vitamin D Insufficiency: Vitamin D3 Megadose Reduces Inflammatory Markers. *Exp. Gerontol.* 2015, 66, 10–16, doi:10.1016/j.exger.2015.03.011.
20. AlFaqih, M.A. Association of Vitamin D Levels and Polymorphisms in Vitamin D Receptor with Type 2 Diabetes Mellitus Available online: <https://www.spandidos-publications.com/10.3892/br.2022.1585> (accessed on 29 February 2024).
21. American Diabetes Association Standards of Medical Care in Diabetes-2022 Abridged for Primary Care Providers. *Clin. Diabetes Publ. Am. Diabetes Assoc.* 2022, 40, 10–38, doi:10.2337/cd22-as01.
22. Holick, M.F. The Vitamin D Deficiency Pandemic: Approaches for Diagnosis, Treatment and Prevention. *Rev. Endocr. Metab. Disord.* 2017, 18, 153–165, doi:10.1007/s11154-017-9424-1.
23. Nimitphong, H.; Holick, M.F. Vitamin D Status and Sun Exposure in Southeast Asia. *Dermatoendocrinol.* 2013, 5, 34–37, doi:10.4161/derm.24054.
24. Sung, C.-C.; Liao, M.-T.; Lu, W.; Wu, C.-C. Role of Vitamin D in Insulin Resistance. *J. Biomed. Biotechnol.* 2012, 2012, 634195, doi:10.1155/2012/634195.
25. Li, X.; Liu, Y.; Zheng, Y.; Wang, P.; Zhang, Y. The Effect of Vitamin D Supplementation on Glycemic Control in Type 2 Diabetes Patients: A Systematic Review and Meta-Analysis. *Nutrients* 2018, 10, 375, doi:10.3390/nu10030375.
26. Seida, J.C.; Mitri, J.; Colmers, I.N.; Majumdar,

- S.R.; Davidson, M.B.; Edwards, A.L.; Hanley, D.A.; Pittas, A.G.; Tjosvold, L.; Johnson, J.A. Clinical Review: Effect of Vitamin D3 Supplementation on Improving Glucose Homeostasis and Preventing Diabetes: A Systematic Review and Meta-Analysis. *J. Clin. Endocrinol. Metab.* 2014, 99, 3551–3560, doi:10.1210/jc.2014-2136.
27. George, P.S.; Pearson, E.R.; Witham, M.D. Effect of Vitamin D Supplementation on Glycaemic Control and Insulin Resistance: A Systematic Review and Meta-Analysis. *Diabet. Med.* 2012, 29, e142–e150, doi:10.1111/j.1464-5491.2012.03672.x.
28. Is Micro Evolution in Tropical Country Women Resulting Low 25(OH) Available online: <https://www.longdom.org/open-access/is-micro-evolution-in-tropical-country-women-resulting-low-25ohd-level-a-cross-sectional-study-in-indonesia-33467.html> (accessed on 29 February 2024).
29. Heaney, R.P. Vitamin D in Health and Disease. *Clin. J. Am. Soc. Nephrol. CJASN* 2008, 3, 1535–1541, doi:10.2215/CJN.01160308.
30. Institute of Medicine (US) Committee to Review Dietary Reference Intakes for Vitamin D and Calcium Dietary Reference Intakes for Calcium and Vitamin D; Ross, A.C., Taylor, C.L., Yaktine, A.L., Del Valle, H.B., Eds.; The National Academies Collection: Reports funded by National Institutes of Health; National Academies Press (US): Washington (DC), 2011;
31. Al-Daghri, N.M.; Al-Attas, O.S.; Alkharfy, K.M.; Khan, N.; Mohammed, A.K.; Vinodson, B.; Ansari, M.G.A.; Alenad, A.; Alokail, M.S. Association of VDR-Gene Variants with Factors Related to the Metabolic Syndrome, Type 2 Diabetes and Vitamin D Deficiency. *Gene* 2014, 542, 129–133, doi:10.1016/j.gene.2014.03.044.
32. Usategui-Martín, R.; De Luis-Román, D.-A.; Fernández-Gómez, J.M.; Ruiz-Mambrilla, M.; Pérez-Castrillón, J.-L. Vitamin D Receptor (VDR) Gene Polymorphisms Modify the Response to Vitamin D Supplementation: A Systematic Review and Meta-Analysis. *Nutrients* 2022, 14, 360, doi:10.3390/nu14020360.
33. Ross, A.C.; Manson, J.E.; Abrams, S.A.; Aloia, J.F.; Brannon, P.M.; Clinton, S.K.; Durazo-Arvizu, R.A.; Gallagher, J.C.; Gallo, R.L.; Jones, G.; et al. The 2011 Report on Dietary Reference Intakes for Calcium and Vitamin D from the Institute of Medicine: What Clinicians Need to Know. *J. Clin. Endocrinol. Metab.* 2011, 96, 53–58, doi:10.1210/jc.2010-2704.
34. Alvina, A.; Immanuel, S.; Harbuwono, D.S.; Suyatna, F.D.; Harahap, A.; Prihartono, J.; Pusparini, P. Effect of Three and Six Months of Vitamin D Supplementation on Glycemic Control and Insulin Resistance in Type 2 Diabetes Mellitus: Randomized Placebo-Controlled Trial. *Indones. Biomed. J.* 2023, 15, 287–295, doi:10.18585/inabj.v15i3.2370.

15%

SIMILARITY INDEX

%

INTERNET SOURCES

15%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1

Alice D da Silva, Julia S Oliveira, Isabela C de Castro, Wanderléia C Paiva, Júnia M G Gomes, Laura C J P Pimenta. "Association of vitamin D and cognition in people with type 2 diabetes: a systematic review", Nutrition Reviews, 2023

Publication

1%
- 2

"Vitamin D and the Lung", Springer Science and Business Media LLC, 2012

Publication

1%
- 3

Christian Trummer, Verena Schwetz, Martina Kollmann, Monika Wölfler et al. "Effects of vitamin D supplementation on metabolic and endocrine parameters in PCOS: a randomized-controlled trial", European Journal of Nutrition, 2018

Publication

1%
- 4

Stefan Pilz, Andreas Hahn, Christiane Schön, Manfred Wilhelm, Rima Obeid. "Effect of Two Different Multimicronutrient Supplements on Vitamin D Status in Women of Childbearing Age: A Randomized Trial", Nutrients, 2017

Publication

1%

5	Dongyou Liu. "Molecular Detection of Human Viral Pathogens", CRC Press, 2019 Publication	1 %
6	"Handbook of vitamin D in human health", Brill, 2013 Publication	<1 %
7	Adrian F. Gombart. "Vitamin D - Oxidative Stress, Immunity, and Aging", CRC Press, 2019 Publication	<1 %
8	Alvina Alvina, Suzanna Immanuel, Dante Saksono Harbuwono, Frans Dhyhanagiri Suyatna et al. "Effect of Three and Six Months of Vitamin D Supplementation on Glycemic Control and Insulin Resistance in Type 2 Diabetes Mellitus: Randomized Placebo-controlled Trial", The Indonesian Biomedical Journal, 2023 Publication	<1 %
9	Chao, Yi-Sheng, Ludovic Brunel, Peter Faris, and Paul Veugelers. "The Importance of Dose, Frequency and Duration of Vitamin D Supplementation for Plasma 25-Hydroxyvitamin D", Nutrients, 2013. Publication	<1 %
10	Xuan Feng, Zhe Wang, Quentin Howlett-Prieto, Nathan Einhorn, Suad Causevic, Anthony T. Reder. "Vitamin D enhances responses to interferon-β in MS", Neurology	<1 %

11

William B. Grant, Fatme Al Anouti, Barbara J. Boucher, Erdiñç Dursun et al. "A Narrative Review of the Evidence for Variations in Serum 25-Hydroxyvitamin D Concentration Thresholds for Optimal Health", Nutrients, 2022

Publication

<1 %

12

Yenny Yenny, Elly Herwana, Raditya Wratsangka. "Mandatory universal use of cloth mask for prevention of coronavirus disease 2019 transmission", Universa Medicina, 2021

Publication

<1 %

13

Martineau, Adrian, and David Jolliffe. ""Vitamin D and Human Health: from the Gamete to the Grave": Report on a meeting held at Queen Mary University of London, 23rd–25th April 2014", Nutrients, 2014.

Publication

<1 %

14

"European Congress on Osteoporosis & Osteoarthritis (ECCEO11-IOF)", Osteoporosis International, 2011

Publication

<1 %

15

"Handbook of nutrition and diet in therapy of bone diseases", Brill, 2016

Publication

<1 %

16

"Handbook of nutrition in heart health",
Wageningen Academic Publishers, 2017

Publication

<1 %

17

Isa Gabriela de Medeiros Cavalcante,
Alexandre Sérgio Silva, Maria José Carvalho
Costa, Darlene Camati Persuhn et al. "Effect
of vitamin D3 supplementation and
influence of BsmI polymorphism of the VDR
gene of the inflammatory profile and
oxidative stress in elderly women with
vitamin D insufficiency", Experimental
Gerontology, 2015

Publication

<1 %

18

Victoria Rontoyanni, Jaqueline Avila, Sapna
Kaul, Rebeca Wong, Sreenivas Veeranki.
"Association between Obesity and Serum
25(OH)D Concentrations in Older Mexican
Adults", Nutrients, 2017

Publication

<1 %

19

Jaeseok Choi, Raban Dusabimana, Fedine
Urubuto, Faustine Agaba et al. "A
standardised neonatal admission record
(NAR) - increasing quality of neonatal care
in Rwanda-a retrospective observational
study", PAMJ Clinical Medicine, 2020

Publication

<1 %

20

S Sinha, V Munikrishnan, J Montgomery, S
Mitchell. "The Impact of Patient-Controlled
Analgesia on Laparoscopic

<1 %

Cholecystectomy", The Annals of The Royal College of Surgeons of England, 2007

Publication

21

Nasser M. Al-Daghri, Abdul Khader Mohammed, Omar S. Al-Attas, Mohammed Ghouse Ahmed Ansari et al. "Vitamin D Receptor Gene Polymorphisms Modify Cardiometabolic Response to Vitamin D Supplementation in T2DM Patients", Scientific Reports, 2017

Publication

<1 %

22

Tatiana L. Karonova, Ksenia A. Golovatyuk, Igor V. Kudryavtsev, Alena T. Chernikova et al. "Effect of Cholecalciferol Supplementation on the Clinical Features and Inflammatory Markers in Hospitalized COVID-19 Patients: A Randomized, Open-Label, Single-Center Study", Nutrients, 2022

Publication

<1 %

23

Zahra Mirzaei-Azandaryani, Yousef Javadzadeh, Elnaz Shaseb, Mojgan Mirghafourvand. "The effects of vitamin D on sleep quality and pregnancy symptoms in pregnant women: a randomized, tripled-blinded and placebo-controlled clinical trial", Nutrition & Food Science, 2023

Publication

<1 %

24

"Nutritional Influences on Bone Health", Springer Science and Business Media LLC, 2016

Publication

<1 %

-
- 25 Mushtaq A. Khan, Hilal A. Dar, Muneer A. Baba, Altaf H. Shah, Bhagat Singh, Nadeem A. Shiekh. "Impact of Vitamin D Status in Chronic Liver Disease", Journal of Clinical and Experimental Hepatology, 2019
Publication <1 %
-
- 26 Sowell, Krista, Carl Keen, and Janet Uriu-Adams. "Vitamin D and Reproduction: From Gametes to Childhood", Healthcare, 2015.
Publication <1 %
-
- 27 Stefan Pilz. "Vitamin D status and arterial hypertension: a systematic review", Nature Reviews Cardiology, 08/18/2009
Publication <1 %
-
- 28 "Eighth European Congress on Clinical and Economic Aspects of Osteoporosis and Osteoarthritis–ECCEO 8", Osteoporosis International, 2008
Publication <1 %
-
- 29 Pang Yao, Derrick Bennett, Marion Mafham, Xu Lin, Zhengming Chen, Jane Armitage, Robert Clarke. "Vitamin D and Calcium for the Prevention of Fracture", JAMA Network Open, 2019
Publication <1 %
-
- 30 Xinling Wen, Fen Li, Xuewen Yu, Li Wang. "Effects of vitamin D supplementation on metabolic parameters of women with polycystic ovary syndrome: a randomized

31

Manuel Rupprecht, Stefan Wagenpfeil, Jakob Schöpe, Reinhold Vieth, Thomas Vogt, Jörg Reichrath. "Meta-Analysis of European Clinical Trials Characterizing the Healthy-Adult Serum 25-hydroxyvitamin D Response to Vitamin D Supplementation", Nutrients, 2023

Publication

<1 %

32

Hussam Alhawari, Yazun Jarrar, Dina Abulebdah, Sara J. Abaalkhail et al. "Effects of Vitamin D Receptor Genotype on Lipid Profiles and Retinopathy Risk in Type 2 Diabetes Patients: A Pilot Study", Journal of Personalized Medicine, 2022

Publication

<1 %

33

Nicole Weidner, Adronie Verbrugghe. "Current knowledge of vitamin D in dogs", Critical Reviews in Food Science and Nutrition, 2016

Publication

<1 %

34

Nutritional Influences on Bone Health, 2013.

Publication

<1 %

35

Abhai Kumar, Debasis Bagchi. "Antioxidants and Functional Foods for Neurodegenerative Disorders - Uses in Prevention and Therapy", CRC Press, 2021

<1 %

36 Cindy D Davis, John A Milner. "Vitamin D and colon cancer", Expert Review of Gastroenterology & Hepatology, 2014 <1 %

Publication

37 Darina Bassil, Maya Rahme, Maha Hoteit, Ghada El-Hajj Fuleihan. "Hypovitaminosis D in the Middle East and North Africa", Dermato-Endocrinology, 2014 <1 %

Publication

38 Linda Aurbibul, Sirintip Sricharoenchai, Orasri Wittawatmongkol, Virat Sirisanthana et al. "Vitamin D status in perinatally HIV-infected Thai children receiving antiretroviral therapy", Journal of Pediatric Endocrinology and Metabolism, 2016 <1 %

Publication

39 Robin M. Daly. "Prevalence of vitamin D deficiency and its determinants in Australian adults aged 25 years and older: A national, population-based study", Clinical Endocrinology, 12/2011 <1 %

Publication

40 Sakshi Tyagi, Shalini Mani. "Combined Administration of Metformin and Vitamin D: A Futuristic Approach for Management of Hyperglycemia", Cardiovascular & Hematological Agents in Medicinal Chemistry, 2024 <1 %

Publication

41 Amjad Hayat, David O'Brien, Paul O'Rourke, Siobhan McGuckin et al. "CD38 expression level and pattern of expression remains a reliable and robust marker of progressive disease in chronic lymphocytic leukemia", *Leukemia & Lymphoma*, 2009 <1 %

Publication

42 Muthmainah, Hanik Badriyah Hidayati, Budi Yanti. "Improving Health for Better Future Life: Strengthening from Basic Science to Clinical Research", CRC Press, 2023 <1 %

Publication

43 Shen, Liang, Qi-Shuai Zhuang, and Hong-Fang Ji. "Assessment of vitamin D levels in type 1 and type 2 diabetes patients: Results from meta-analysis", *Molecular Nutrition & Food Research*, 2016. <1 %

Publication

44 Sukumar, D., S.A. Shapses, and S.H. Schneider. "Vitamin D supplementation during short-term caloric restriction in healthy overweight/obese older women: Effect on glycemic indices and serum osteocalcin levels", *Molecular and Cellular Endocrinology*, 2015. <1 %

Publication

Exclude bibliography On