

ISSN: 1874-2106

The Open Dentistry Journal



VIEW ARTICLES ON
PUBMED CENTRAL



BENTHAM OPEN



[Submit Abstract Online](#) [Submit Issue Proposal](#) [Become an Editorial Board Member](#)
[Become a Reviewer](#) [Become an Executive Guest Editor](#) [Become a Section Editor](#)
[Become a Member](#)

AIMS AND SCOPE

The Open Dentistry Journal is an Open Access journal, which publishes research articles, reviews/mini-reviews, letters, case reports and guest edited single topic issues in all areas of dentistry and the oral cavity.

The journal encourages submissions related to the following fields of dentistry:

- Dental Implants
- Dental Biomaterials Science
- Endodontology
- Fixed and Removable Prosthodontics
- Management of Dental Disease
- Maxillofacial Surgery
- Periodontology
- Restorative Dentistry
- Operative Dentistry
- Oral Pathology
- Oral Medicine
- Clinical Trials
- Translational Research

The Open Dentistry Journal, a peer-reviewed journal, is an important and reliable source of current information on important recent developments in the field. The emphasis will be on publishing quality papers rapidly and freely available to researchers worldwide.

RECENT ARTICLES

RESEARCH ARTICLE

Periodontal Disease Markers among Patients with Long COVID: A Case-control Study
Marie Louisa, Wita Anggraini, Ricky Anggara Putranto, Olivia Nauli Komala and Nicola De Angelis

The Open Dentistry Journal, 2023; 17: e187421062307030.
Electronic Publication Date: July 24, 2023

[View Abstract](#) [View Fulltext HTML](#) [Download PDF](#) [Download ePub](#)

RESEARCH ARTICLE



Evaluation of the Effect of Two Surface Treatment Methods on Shear Bond Strength between Porcelain Veneer and Two Different Zirconia Blocks
Ali Barzegar, Tahereh Ghaffari, Ramin Negahdari, Solmaz Maleki Dizaj and Reyhaneh Saghari

The Open Dentistry Journal, 2023; 17: e187421062305300.
Electronic Publication Date: July 11, 2023

[View Abstract](#) [View Fulltext HTML](#) [Download PDF](#) [Download ePub](#)

REVIEW ARTICLE



Antimicrobial Irrigation Solutions in Root Canal Treatment: A Glance at the Past, the Present, and the Future
Abubaker Qutieshat, Nutayla Al Harthy, Shima Al Busaidi, Ahmed Al Sadoon, Dima Al Sayahien, Maryam Sedqi, Sumalya Al Rashdi and Samiya Al Ghamman

The Open Dentistry Journal, 2023; 17: e187421062306010.
Electronic Publication Date: July 04, 2023

[View Abstract](#) [View Fulltext HTML](#) [Download PDF](#) [Download ePub](#)

RESEARCH ARTICLE



The Cell Biocompatibility of a new Polycaprolactone-based Endodontics Sealer Compared with AH Plus Sealer
Mostafa Torbati, Anita Basiri Maleki, Mohammadali Torbati, Shahriar Shahi, Solmaz Maleki Dizaj and Simin Sharifi

The Open Dentistry Journal, 2023; 17: e187421062305220.
Electronic Publication Date: July 04, 2023

[View Abstract](#) [View Fulltext HTML](#) [Download PDF](#) [Download ePub](#)

CLINICAL TRIAL STUDY



Effect of Different Desensitizing Protocols on Pulp Inflammatory Responses in Whitened Teeth: A Randomized Clinical Trial
Antonia Patricia Oliveira Barros, Reissa Pinheiro de Mendonça, Crislaine de Melo Alencar, João de Jesus Viana Pinheiro, Cláudia Pires Rothbarth, Sérgio de Melo Alves Júnior and Cecy Martins Silva

The Open Dentistry Journal, 2023; 17: e187421062305190.
Electronic Publication Date: July 04, 2023

[View Abstract](#) [View Fulltext HTML](#) [Download PDF](#) [Download ePub](#)

[View More Articles](#)



EDITORIAL BOARD

EDITOR-IN-CHIEF



Ibtisam Al-Hashimi
Scopus

[Biography and Affiliation](#)

Biography: Prof. Ibtisam Al-Hashimi, BDS, M.S. and
[READ MORE](#)

CO-EDITOR



Luigi Checchi
Scopus

[Biography and Affiliation](#)

Biography: Dr. Luigi Checchi completed Medical an
[READ MORE](#)

FRONTIER SECTION EDITOR



Gerardo Pellegrino
Scopus

[Biography and Affiliation](#)

Biography: Dr. Pellegrino graduated in Dentistry fr
[READ MORE](#)

EDITORIAL BOARD MEMBERS



Emanuel Savio de Souza Andrade
Scopus

[Biography and Affiliation](#)

Biography: Dr. Emanuel Savio is an Associate Profe
[READ MORE](#)



Maria Teresa Vinuesa Aumedes
Scopus

[Biography and Affiliation](#)

Biography: Dr. Teresa Vinuesa Aumedes is a Medici
[READ MORE](#)



Nicola Alberto Valente
Scopus PubMed

[Biography and Affiliation](#)

Biography: Dr. Nicola Alberto Valente DDS, MS, Phl
[READ MORE](#)



Nicola De Angelis
Scopus

[Biography and Affiliation](#)

Biography: Dr. De Angelis received his dental degre
[READ MORE](#)

[SUBMIT MANUSCRIPT](#)

The
Open Dentistry
Journal



ISSN (Online): 1874-2106

VOLUME: 17, 2023

[DOWNLOAD FLYER](#)

IMPACT FACTOR **1.3**



SEARCH ARTICLES

SEARCH TERM

SEARCH CRITERIA:

Article Title / Keywords / Abst

ARTICLE TYPE:

All Article Types

[CLEAR](#)[SEARCH](#)

JOIN OUR TEAM

[Join as a Board Member](#)[Join as an Executive Guest Editor](#)[Join as a Reviewer](#)[Join as a Section Editor](#)

MARKETING OPPORTUNITIES

[Advertising Policy](#)[Kudos](#)[Promotional Services](#)


ADVERTISEMENT

CURRENT DENTISTRY

LIMITED TIME **COMPLEMENTARY**
OPEN ACCESS OFFER

PAPERS SUBMITTED BEFORE
DECEMBER 31, 2023
WILL BE PUBLISHED AS
PROMOTIONAL OPEN ACCESS

BENTHAM SCIENCE
Publishers of Quality Research



The Open Dentistry Journal

Content list available at: <https://opendentistryjournal.com>



RESEARCH ARTICLE

Periodontal Disease Markers among Patients with Long COVID: A Case-control Study

Marie Louisa¹, Wita Anggraini², Ricky Anggara Putranto¹, Olivia Nauli Komala¹ and Nicola De Angelis^{1,3,*}

¹Department of Periodontology, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia

²Department of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia

³Unit of Implant and Prosthetics, Department of Surgical Sciences and Integrated Diagnostics, University of Genoa, Genoa, Italy

Abstract:

Background:

Long-COVID affects approximately 32-87% of COVID-19 patients and causes persistent symptoms that last more than 4 weeks after initial infection. Long COVID is associated with a notable cytokine triad, namely IL-1 β , IL-6, and TNF- α . Periodontal disease also involves proinflammatory cytokines production, including IL-1 β , IL-6, and TNF- α . Consequently, long-COVID, which has an impact on proinflammatory cytokine release, could have an impact on periodontal status.

Objective:

This study aims to see whether long COVID affects periodontal status severity based on proinflammatory cytokines levels involved in both diseases, namely IL-1 β , IL-6, and TNF- α .

Methods:

Patients were divided into periodontitis or gingivitis patients and then were further divided into two groups, previous COVID-19 patients and non-COVID-19 patients (controls). Gingival sulcus fluids were obtained from each patient using paper points inserted in the patients' sulcus, and ELISA tests were carried out to measure IL-1 β , IL-6, and TNF- α levels.

Results:

Levene Test indicated that there were no substantial differences between IL-1 β , IL-6, and TNF- α levels (0.057, 0.135, and 0.341, respectively) in COVID-19 patients with gingivitis in comparison to the control group with gingivitis, with average IL-1 β , IL-6 and TNF- α levels seen higher in the control group compared to COVID-19 patients. There were also no substantial differences between IL-1 β , IL-6, and TNF- α levels (1.00, 0.567, and 0.666, respectively) between COVID-19 patients with periodontitis and the control group with periodontitis. Although higher levels of IL-6 and TNF- α were found higher in COVID-19 patients in comparison to the control group.

Conclusion:

Levels of IL-6 and TNF- α in periodontitis patients with long COVID were higher than controls. But despite that, higher IL-1 β , IL-6, and TNF- α levels were not found in long COVID subjects with gingivitis, as well as IL- β levels in the periodontitis group. Further studies with more subjects are needed to determine the connection between these two diseases further.

Clinical Trial Registration No: The clinical trial registration of this research is 014/S3/KEPK/FKG/10/2021.

Keywords: Long COVID, IL-1 β , IL-6, TNF- α , Long COVID, IL-6, TNF- α , Periodontal disease, Gingivitis.

Article History

Received: April 07, 2023

Revised: June 11, 2023

Accepted: June 21, 2023

1. INTRODUCTION

Coronavirus disease 19 (COVID-19), which has had detrimental consequences over more than the preceding two years, was provoked by Severe Acute Respiratory Distress Syndrome Coronavirus-2 (SARS-CoV-2), causing the demise

of more than 5 million people. Approximately 20% of those who were infected by SARS-CoV-2 would require intensive care unit (ICU) admission, where subsequently, approximately 25% of these patients were deceased, mostly as a consequence of severe inflammation and thromboembolic complications [1]. COVID-19 also has an immense impact on dentistry worldwide, including dentists, dental patients, and also dental practice. Dentists are at risk and prone to COVID-19 infection

* Address correspondence to this author at the Department of Surgical Sciences, Integrated Diagnostics and Periodontology, University of Genoa (IT), Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia;
E-mail: nicolaantonio.deangelis@edu.unige.it

due to exposure to pathogens, long working hours, psychological stress, and fatigue [2]. In addition, COVID-19 has caused mental and mental health problems in both COVID-19 patients and non-COVID patients undergoing self-isolation, social distancing or quarantine, leading to problems such as loneliness, depression and isolation [3].

Periodontal disease is a multifactorial infectious disease, with plaque accumulation as the main cause, predisposed by a lot of factors, such as poor oral hygiene, age, sex, diabetes, HIV, menopause, pregnancy, and smoking, *etc.* Psychological stress affects periodontium by way of neuro-immune endocrinological mechanisms. Psychological stress affects the hypothalamus, which alters the adrenal, producing several substances (catecholamine, chromogranin A, neuropeptides), resulting in a decreased immune response. Stress also affects the autonomic nervous system, where the adrenal glands release prostaglandins and other proteases, subsequently affecting the periodontium. These two factors both contribute to periodontitis and poor effectiveness of periodontal treatment [4]. Patients with COVID-19 also commonly have systemic manifestations, giving rise to complications affecting the respiratory, endocrine, metabolic, cardiovascular, hematological, vascular, gastrointestinal, neurological, renal, musculoskeletal and hepatic systems [5].

First, SARS-CoV-2 enters through the binding of ACE2 protein to alveolar epithelial cells, which subsequently respond to the progression of COVID-19 by initiating the production of chemokines, interferons, interleukins (IL) and tumor necrosis factor (TNF- α), which then provokes the innate and adaptive immune systems [6]. Evidence of late shows that a range of manifestations can still persevere after the resolution of acute infection, which is referred to as long COVID, that exerts an influence on multiple organs [7]. National Health Institute for Health and Care Excellence defines this condition as manifestations that continue or develop after acute COVID-19 infection and cannot be justified by an alternative diagnosis, which entails ongoing symptomatic COVID-19 during 4-12 weeks following initial infection, and post-COVID-19 syndrome surpassing 12 weeks following the initial infection. The National Institutes of Health (NIH) also defines long COVID as a sequelae that persist surpassing 4 weeks following the initial infection [8]. Long COVID is clinically characterized by the presence of fatigue, exercise intolerance, brain fog, shortness of breath, joint pain, fever, sleep, anxiety disorders, gastrointestinal symptoms, and palpitations that could persist for months. Around 21.7-87% of COVID-19 patients undergo long COVID [9].

The immune response is a compelling factor that determines the interaction between pathogens and the host. When this process becomes dysregulated, a cytokine storm can result, resulting in rapid production of cytokines that control immunopathological responses. Altered serum levels of cytokines such as IFN- α , IFN- γ , IL-1 β , IL-6, IL-12, IL-18, IL-33, TNF- α and TGF- β are found correlated to COVID-19 [10]. In spite of the fact that a lot of other risk factors (age, comorbidities) could affect COVID-19 severity, one study found a connection between long COVID with the elevation of notable cytokine triads, IL-1 β , IL-6, and TNF- α , in which three

are all involved in pain perception, anxiety, depression, and inflammation [11].

On the other hand, periodontal disease has a high prevalence of 50% in the adult population all over the world. Even in healthy circumstances, the periodontal tissues have immune cells continuously present in the gingiva, maintaining a balance between the host and the oral biofilm and keeping the immune response active [12]. Periodontal disease is a multifactorial infectious disease that is mediated by immune responses. Higher cytokine levels in periodontal tissue and serum were found in patients with periodontal disease, similar to the cytokine storm occurring in COVID-19 patients [13].

Periodontal infection is also associated with systemic diseases, including bacteremia and systemic release of inflammatory mediators. Cytokines are produced by fibroblasts, epithelial cells, neutrophils, and macrophages in the acute phase of periodontal disease; and produced by lymphocytes in advanced periodontal diseases. During an inflammatory response, the first stage of periodontal disease is characterized by the secretion of proinflammatory cytokines such as IL-1, IL-6, and TNF- α in opposition to periodontal bacteria [14]. An increase in blood flow initiates this disease, amplified vascular permeability, and the ingress of neutrophils, and monocyte macrophages, with ensuing T cells and B cells appearing, producing cytokines (*e.g.*, IL-1 β , IL-6, and TNF- α), degrading epithelial cell tissues, and consequently creating bone resorption [15].

In this regard, it is possible that long COVID, which impacts on proinflammatory cytokine release, could affect periodontal status. This is based on the fact that periodontal disease could occur due to increased host inflammatory responses as a COVID-19 consequence [16]. This study investigates whether long COVID affects proinflammatory cytokines, namely IL-1 β , IL-6 and TNF- α in periodontal disease. Therefore, in this study, we aim to compare inflammatory mediator levels of long COVID patients (patients who have experienced COVID-19 previously) with inflammatory periodontal diseases, compared to non-COVID-19 patients (controls).

2. MATERIALS AND METHODS

The research was a pilot study, in which we divided the subjects into two groups: gingivitis and periodontitis. In the gingivitis group, a sum of twenty samples was used, involving 10 COVID-19 patients and 10 controls. A sum of four samples in the gingivitis group was subtracted due to low levels of protein exhibited after the Bradford protein assay test was done. Therefore, these four samples were unable to be continued with the ELISA test.

Whereas in the periodontitis group, 20 samples were used, including 10 COVID-19 patients and 10 controls. The results of this clinical study suggested that there was a correlation between periodontal disease and long COVID, delineated by altered levels of proinflammatory cytokines. This hypothesis was substantiated by higher levels of IL-6 and TNF- α that were observed higher in previous COVID-19 patients with periodontitis in this study.

2.1. Obtaining Samples

Samples were obtained at the dental and oral hospital of Universitas Trisakti and the research was conducted at the biomedical laboratory of the Faculty of Dentistry, Universitas Trisakti. Subjects were determined based on inclusion criteria (gingivitis patients with a probing depth of ≤ 3 mm and bleeding on probing / BOP score $\geq 10\%$; periodontitis patients with clinical attachment loss / CAL on ≥ 2 non-adjacent teeth or ≥ 3 mm on buccal/oral surfaces on ≥ 2 teeth; patients with a minimum amount of 20 teeth; COVID-19 patients who tested positive with a PCR test and showed symptoms 4 weeks after the onset of the first symptoms, Long-COVID-19 patients who completed an online Long-COVID-Symptoms questionnaire and exclusion criteria (active smokers, type II diabetes mellitus patients, pregnant and lactating patients, patients who underwent periodontal treatment during the pandemic, patients with aggressive periodontitis) were distinguished.

Throughout the course of the study, there has not been a study conducted on periodontal biomarkers observed in previous COVID-19 patients / patients with a COVID-19 history. Hence, this pilot study was commenced.

Informed consent were then obtained from all subjects after an explanation regarding the research was given, in conjunction with periodontal status checking (probing depth, Oral Hygiene Index-simplified/OHI-s, and CAL) to discern gingivitis and periodontitis patients. Subjects were then divided into two groups: previous COVID-19 patients and non-COVID-19 patients (controls), distinguished by a history of COVID-19 (tested positive by PCR test).

Gingival sulcus fluids were obtained to determine IL-1 β , IL-6, and TNF- α levels. One tablet of Phosphate-buffered saline (PBS) was dissolved in 100 ml, then 200 μ l of PBS solution was transferred into an Eppendorf tube, which was then inserted in a cooler box containing ice cubes. Subjects were given cheek retractors, and the area was isolated using cotton rolls. Plaques were then removed, and tooth surfaces were dried. Consequently, samples were collected using paper points on every tooth exhibiting positive BOP. Paper points were inserted in the sulcus to a depth of 1 mm for 30 seconds. Samples with saliva and blood contamination were excluded. The collected gingival sulcus fluids were inserted in Eppendorf tubes containing 200 μ l of PBS solution and stored at -80C degrees.

2.2. Samples Preparation

Samples stored at -80°C degrees were prepared by thawing them at room temperature and then vortexed to obtain even concentrations. Tubes were centrifuged at 2000 g, 4°C for 5 minutes. Concentration of extract obtained from centrifugation was then measured with 9 standards at different concentrations, namely, 2000 μ g/ml, 1500 μ g/ml, 1000 μ g/ml, 750 μ g/ml, 500 μ g/ml, 250 μ g/ml, 125 μ g/ml, 25 μ g/ml, and 0 using Bradford

method. Calculations were done for the samples with the highest total protein concentrations, and consequently, protein concentrations were equated into 250 μ g/ml by adding PBS to gather stock volume samples to the tune of 250 μ l.

2.3. ELISA Test

IL-1 β , IL-6, and TNF- α levels were measured using an ELISA test. Samples were obtained and thawed at room temperature. Twenty ml of wash buffer concentrate was diluted 25 times with deionized or distilled water to yield 500 ml of 1x Wash Buffer. Reagents, standard solutions, and samples are also brought to room temperature. Strips used in the research were inserted in frames (unused frames were stored at 2-8°C). As much as 50 μ l standard was added to the standard wells. In sample wells, 40 μ l samples are added. In addition, in sample wells, according to the cytokine being tested, add 10 μ l of anti-IL-1 β antibody for the IL-1 β test, 10 μ l anti-IL-6 antibody for the IL-6 test, or 10 μ l anti-TNF- α antibody for TNF- α test.

An amount of 50 μ l streptavidin-HRP was then added to the sample and standard wells, mixed, and covered with a sealer, and incubated for 60 minutes at 37°C. Afterwards, the sealers were removed, the plates were washed 5 times with a wash buffer (soaking the wells with at least 0,35 ml wash buffer for 30 seconds to 1 minute for each wash), and blotted with paper towels. Each well was added 50 μ l each of substrate solution A and solution B, covered with new sealers, and incubated for 10 minutes at 37°C in the dark. Consequently, 50 μ l of stop solution is added to each well, changing the blue color into yellow. Within ten minutes following the addition of a stop solution, the optical density value for each well is then read by a microplate reader set to 450 nm. Data analysis is then done.

3. RESULTS

In this study, there was a sum of 40 patients who were divided into two groups (20 COVID-19 patients and 20 controls), in which data collection was carried out randomly according to the predetermined inclusion and exclusion criteria. The 20 subjects in the control group consisted of 50% (10 subjects) of gingivitis, where 50% (5 subjects) were aged between 21-30 years, 20% (2 subjects) were aged 31-40 years and 30% (3 subjects) were aged 41-50 years. The periodontitis subjects in the control group (10 subjects) consisted of 30% (3 subjects) aged 21-30 years, 20% (2 subjects) aged 31-40 years, 40% (4 subjects) aged 41-50 years, and 10% (1 subject) aged 51-60 years.

In the previous COVID-19 patient group with a sum of 20 subjects, half of the subjects (10 people) experiencing gingivitis consisted of 90% (9 subjects) aged 21-30 years and 10% (1 subject) aged 51-60 years. The other half of the previous COVID-19 patient group (10 people) experiencing periodontitis consisted of 50% (5 subjects) aged 21-30 years, 20% (2 subjects) aged 31-40 years, 10% (1 subject) aged 41-50 years, and 10% (1 subject) aged 61-70 years. The comorbidities of each subject were not recorded.

Table 1. Independent sample tested for gingivitis and periodontitis.

-	-	Mean (ng/L)	Std Deviation	Sig
IL-1 β	The control group with gingivitis	447.8143	47.38173	0.057
	COVID-19 patients with gingivitis	390.1222	60.49859	0.057
	The control group with periodontitis	182.8500	83.86997	1.000
	COVID-19 patients with periodontitis	182.8400	47.07463	1.000
IL-6	The control group with gingivitis	88.0857	12.04138	0.135
	COVID-19 patients with gingivitis	76.0111	17.02487	0.135
	The control group with periodontitis	33.1700	18.91243	0.567
	COVID-19 patients with periodontitis	37.2300	11.22884	0.567
TNF- α	The control group with gingivitis	48.9286	6.41293	0.341
	COVID-19 patients with gingivitis	43.7778	12.54370	0.341
	The control group with periodontitis	11.8600	7.15871	0.666
	COVID-19 patients with periodontitis	12.9600	3.27930	0.666

Based on long COVID-19 symptoms experienced by 20 COVID-19 patients, it was found that 45% (9 subjects) were patients with chronic COVID-19 because they had symptoms that persisted for more than 12 weeks after being declared negative for infection with the SARS-CoV-2 virus, while 55% (11 subjects) were post-acute COVID. Fatigue was the symptom with the highest percentage, namely 35% (7 subjects), 30% (6 subjects) experienced cough, 20% (4 subjects) experienced headaches, and 15% (5 subjects) experienced other symptoms such as sore throat, difficulty concentrating, flu, nausea, to shortness of breath.

OHI-s scores of each patient were tested statistically and were not significantly different. In the gingivitis group, 16 samples were used and were tested for normality using the Shapiro-Wilk test. Several samples in the gingivitis group were removed due to contamination. Thus, the protein levels were too low to be analyzed by ELISA test. Normality tests indicated that the data were distributed normally. The statistical test used for the gingivitis group was an independent T-test (Table 1). The periodontitis samples were used without any subtraction in the sample amount and tested for normality using the Shapiro-Wilk test, which showed that the data were distributed normally. The statistical test used for the periodontitis group was also an independent T-test (Table 1).

The Levene Test indicated that IL-1 β , IL-6, and TNF- α significance levels in the gingivitis group were 0.057, 0.135, and 0.341, respectively, which implies that there were no substantial differences between IL-1 β , IL-6 and TNF- α levels among COVID-19 patients with gingivitis and controls with gingivitis. The average IL-1 β , IL-6, and TNF- α levels were also higher in controls than in COVID-19 patients.

The significance levels for IL-1 β , IL-6, and TNF- α in the periodontitis group were 1.00, 0.567, and 0.666, respectively, which also implies that there are no substantial differences between the IL-1 β , IL-6 and TNF- α levels among COVID-19 patients with periodontitis and controls with periodontitis. However, higher levels of IL-6 and TNF- α were elevated in COVID-19 patients compared to controls.

4. DISCUSSION

In addition to systemic manifestations, COVID-19

correlates with local oral cavity signs, most commonly dysgeusia, taste loss, and mucosal lesions. Other periodontal tissues involving clinical manifestations still have an undecided relation to COVID-19 [17]. These oral cavity manifestations, their plausible relationship with oral disease, and the probable mechanisms of hyperinflammation underlying these two diseases signify an association between COVID-19 and oral disease [18]. Long-term inflammation in patients with COVID-19 may give rise to a pathological response in periodontal tissues, precipitating fibrinogen degradation, coagulation cascade, and altering bacterial flora, which could prompt or aggravate periodontal disease [19].

Amplified systemic inflammation could result in periodontal disease, which is a multifactorial infectious disease mediated by immune responses. Increased levels of cytokines in periodontal tissues and serum are found in periodontal disease patients juxtaposed to healthy controls [20]. This assertion is further supported by one case-control study by Anand *et al.*, which ascertained that COVID-19 patients had higher average values of plaque scores, amount of mobile teeth, gingival bleeding score, probing depth, recession, and clinical attachment loss as opposed to controls, denoting a significant association between COVID-19 and periodontal disease [21]. In one review study, Kanchana Sukumar stated that the levels of serum proinflammatory cytokine levels in COVID-19 patients were also elevated, particularly IFN- γ , IFN- γ -induced protein 10, IL-1 β , IL-6, IL-12, and monocyte chemoattractant protein (MCP-1) [22]. In two other meta-analyses studies, higher levels of IL-6 are detected in complicated COVID-19 cases than in those with non-severe conditions [23].

There are a number of undetermined matters associated with long COVID, such as its etiology, prevention, and treatment. It is known that older age, obesity, gender, hypertension, immunosuppression, asthma, and severe acute-phase disease are all associated with increased long COVID risk. Psychological stress is also associated with respiratory tract infection severity and duration. Stress may activate the hypothalamic-pituitary-adrenal axis, consequently dysregulating the immune system. Thus, psychological stress may be a risk factor for long COVID. [24].

Long-COVID is characterized by persistent immune activation that includes elevated levels of IL-1, IL-6 and TNF, which supports the two hypotheses of the immunopathogenesis of Long-COVID, namely the constant immune response against incessant viral antigens and/ or chronic reprogramming of immune cells. A cohort by Schultheiß *et al.* has proven these two hypotheses, substantiating the notion that long COVID is associated with heightened plasma levels of IL-1 β , IL-6 and TNF- α [11]. It is then presumed that long COVID, which inflicts systemic inflammation by increasing cytokines related to periodontal diseases (*e.g.*, IL-1 β , IL-6, and TNF- α), would influence periodontal disease severity. Hence, this study is conducted to discern this presumption.

Interleukin-1 β is a significant pro-inflammatory cytokine that plays a role in infection and injury. They contribute to host reactions toward pathogens and aggravate damage in chronic diseases and tissue injuries [25]. The other proinflammatory cytokines, IL-6 and TNF- α , promote leukocyte recruitment, activation, and differentiation, along with B-cell maturation and T-helper cell subset expansion. Both of these cytokines are associated as an essential part of the acute COVID-19 immune response. For the most part, IL-6 is more predictive of poor COVID-19 outcomes (*e.g.*, respiratory failure, the need for mechanical ventilation) [26].

In periodontal disease pathogenesis, the IL-6 family is identified as a principal factor. In which several studies stated that SARS-Cov-2 infection, in particular, induces IL-6 production up to the extent of 1000-fold above the standard range in several cases. Elevated IL-6 levels can precipitate adverse clinical outcomes and are associated with cytokine storm pathogenesis associated with both periodontitis as well as COVID-19, further denoting the plausible association between the two diseases [23,27]. Interleukin-6 also plays a role in innate and adaptive immunity by initiating the differentiation of B lymphocytes into antibody-producing cells. IL-8, on the other hand, acts as a neutrophil chemoattractant [28]. Additionally, TNF- α is recognized to induce tissue inflammation and endothelial activation. In uncontrolled circumstances, TNF- α could induce various inflammatory diseases involving multiple organ systems [29].

But despite this finding, this study showed that the levels of proinflammatory cytokines were not higher in COVID-19 patients compared to controls in the gingivitis group. Interleukin-1 β levels were not found to be higher either in the periodontitis group. This study also revealed no substantial differences between IL-1 β , IL-6, and TNF- α between COVID-19 patients and controls in both gingivitis and periodontitis groups. Several samples in the gingivitis group were withdrawn due to low levels of protein. However, this subtraction in the gingivitis group leads to a limitation of the study results.

CONCLUSION

Observations found in this study put forward a plausible connection linking periodontal disease and long COVID. Hypothetically, long COVID could exert an influence on periodontal disease, escalating systemic inflammation by increasing certain cytokines, such as IL-1 β , IL-6, and TNF- α ,

in which there are also correlated to periodontal diseases. In this study, we found increased levels of IL-6 and TNF- α in periodontitis patients with long COVID compared to controls. However, higher levels of IL-1, IL-6 and TNF were not found in long-COVID patients with gingivitis, nor were IL levels in the periodontitis group. These could be due to a small number of subjects or the subtraction of some samples. A further clinical study with many subjects is needed to assess the relationship between these two diseases. Moreover, the discovery of long COVID impact on periodontal diseases clinically could raise awareness among clinicians to be more mindful of COVID-19's long-term impact on the dentistry field, particularly on periodontal health.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

This research was approved by the Ethics Committee for Health Research, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia.

HUMAN AND ANIMAL RIGHTS

No animals were used in this research. All procedures performed in studies involving human participants were in accordance with the 1975 Declaration of Helsinki, as revised in 2013.

CONSENT FOR PUBLICATION

Informed consent was obtained from all participants.

STANDARDS OF REPORTING

CONSORT guidelines were followed.

AVAILABILITY OF DATA AND MATERIAL

The data and supportive information are available within the article.

FUNDING

This research received funding from a grant from the Faculty of Dentistry, Trisakti University, Jakarta, Indonesia.

CONFLICT OF INTEREST

Nicola De Angelis is a member of the editorial advisory board of the journal The Open Dentistry Journal.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- [1] Vernuccio F, Lombardo FP, Cannella R, *et al.* Thromboembolic complications of COVID-19: The combined effect of a pro-coagulant pattern and an endothelial thrombo-inflammatory syndrome. Clin Radiol 2020; 75(11): 804-10. [<http://dx.doi.org/10.1016/j.crad.2020.07.019>] [PMID: 32829885]
- [2] Humagain M, Humagain R, Rokaya D. Dental practice during COVID-19 in Nepal: A descriptive cross-sectional study. JNMA J Nepal Med Assoc 2020; 58(230): 764-9. [<http://dx.doi.org/10.31729/jnma.5022>] [PMID: 34504378]
- [3] Rokaya D, Koontongkaew S. Can coronavirus disease-19 lead to temporomandibular joint disease? Open Access Maced J Med Sci 2020; 8(T1): 142-3.

- [http://dx.doi.org/10.3889/oamjms.2020.5003]
- [4] Gunepin M, Derache F, Trousselard M, Salsou B, Risso JJ. Impact of chronic stress on periodontal health. *J Oral Med Oral Surg* 2018; 24(1): 44-50. [http://dx.doi.org/10.1051/mbcb/2017028]
- [5] Paul G, Mahajan RK, Mahajan R, Gautam P, Paul B. Systemic manifestations of COVID-19. *J Anaesthesiol Clin Pharmacol* 2020; 36(4): 435-42. [http://dx.doi.org/10.4103/joacp.JOACP_359_20] [PMID: 33840920]
- [6] Mehandru S, Merad M. Pathological sequelae of long-haul COVID. *Nat Immunol* 2022; 23(2): 194-202. [http://dx.doi.org/10.1038/s41590-021-01104-y] [PMID: 35105985]
- [7] Davis HE, McCorkell L, Vogel JM, Topol EJ. Long COVID: major findings, mechanisms and recommendations. *Nat Rev Microbiol* 2023; 21(3): 133-46. [http://dx.doi.org/10.1038/s41579-022-00846-2] [PMID: 36639608]
- [8] Crook H, Raza S, Nowell J, Young M, Edison P. Long covid—mechanisms, risk factors, and management. *BMJ* 2021; 374: n1648. [http://dx.doi.org/10.1136/bmj.n1648] [PMID: 34312178]
- [9] Raveendran AV, Jayadevan R, Sashidharan S. Long COVID: An overview. *Diabetes Metab Syndr* 2021; 15(3): 869-75. [http://dx.doi.org/10.1016/j.dsx.2021.04.007] [PMID: 33892403]
- [10] Queiroz MAF, Neves PFM, Lima SS, *et al.* Cytokine profiles associated with acute COVID-19 and long COVID-19 syndrome. *Front Cell Infect Microbiol* 2022; 12(June): 922422. [http://dx.doi.org/10.3389/fcimb.2022.922422] [PMID: 35846757]
- [11] Schultheiß C, Willscher E, Paschold L, *et al.* The IL-1 β , IL-6, and TNF cytokine triad is associated with post-acute sequelae of COVID-19. *Cell Rep Med* 2022; 3(6): 100663. [http://dx.doi.org/10.1016/j.xcrm.2022.100663] [PMID: 35732153]
- [12] Könönen E, Gursoy M, Gursoy U. Periodontitis: A multifaceted disease of tooth-supporting tissues. *J Clin Med* 2019; 8(8): 1135. [http://dx.doi.org/10.3390/jcm8081135] [PMID: 31370168]
- [13] Gofur NP. Impact of SARS-CoV-2 on periodontal tissue manifestation. *J Int Oral Health* 2020; 12(8): 90. [http://dx.doi.org/10.4103/jioh.jioh_205_20]
- [14] Fabri GMC. Potential link between COVID-19 and periodontitis: Cytokine storm, immunosuppression, and dysbiosis. *Ohdm* 2020; 19(December): 1-5.
- [15] Taba M, Kinney J, Kim AS, Giannobile WV. Diagnostic biomarkers for oral and periodontal diseases. *Dent Clin North Am* 2005; 49(3): 551-71. [http://dx.doi.org/10.1016/j.cden.2005.03.009]
- [16] Gupta S, Mohindra M, Singla M, *et al.* The clinical association between Periodontitis and COVID-19. *Clin Oral Invest* 2022; 26(2): 1361-74. [http://dx.doi.org/10.1007/s00784-021-04111-3] [PMID: 34448073]
- [17] Srinivasan M. Taste dysfunction and long COVID-19. *Front Cell Infect Microbiol* 2021; 11: 716563. [http://dx.doi.org/10.3389/fcimb.2021.716563] [PMID: 34336725]
- [18] Brandini DA, Takamiya AS, Thakkar P, Schaller S, Rahat R, Naqvi AR. Covid-19 and oral diseases: Crosstalk, synergy or association? *Rev Med Virol* 2021; 31(6): e2226. [http://dx.doi.org/10.1002/rmv.2226] [PMID: 33646645]
- [19] Drozdik A. COVID-19 and SARS-CoV-2 infection in periodontology: A narrative review *J Periodont Res* 2022; 1-9. [http://dx.doi.org/10.1111/jre.13034]
- [20] Ramadan DE, Hariyani N, Indrawati R, Ridwan RD, Diyatri I. Cytokines and chemokines in periodontitis. *Eur J Dent* 2020; 14(3): 483-95. [http://dx.doi.org/10.1055/s-0040-1712718] [PMID: 32575137]
- [21] Anand PS, Jadhav P, Kamath KP, Kumar SR, Vijayalaxmi S, Anil S. A case-control study on the association between periodontitis and coronavirus disease (COVID-19). *J Periodontol* 2022; 93(4): 584-90. [http://dx.doi.org/10.1002/JPER.21-0272] [PMID: 34347879]
- [22] Sukumar K, Tadepalli A. Nexus between COVID-19 and periodontal disease. *J Int Med Res* 2021; 49(3) [http://dx.doi.org/10.1177/03000605211002695] [PMID: 33745336]
- [23] Darestani MN, Akbari A, Yaghobee S, Taheri M, Akbari S. COVID-19 pandemic and periodontal practice: The immunological, clinical, and economic points of view. *BioMed Res Int* 2022; 2022: 1-10. [http://dx.doi.org/10.1155/2022/3918980] [PMID: 35047633]
- [24] Wang S, Quan L, Chavarro JE, *et al.* Associations of depression, anxiety, worry, perceived stress, and loneliness prior to infection with risk of post-COVID-19 conditions *JAMA Psychiatry* 2022; 79(11): 1081-91. [http://dx.doi.org/10.1001/jamapsychiatry.2022.2640]
- [25] Lopez-Castejon G, Brough D. Understanding the mechanism of IL-1 β secretion. *Cytokine Growth Factor Rev* 2011; 22(4): 189-95. [http://dx.doi.org/10.1016/j.cytogfr.2011.10.001] [PMID: 22019906]
- [26] Peluso MJ, Lu S, Tang AF, *et al.* Markers of immune activation and inflammation in individuals with postacute sequelae of severe acute respiratory syndrome coronavirus 2 infection. *J Infect Dis* 2021; 224(11): 1839-48. [http://dx.doi.org/10.1093/infdis/jiab490] [PMID: 34677601]
- [27] Qi M, Sun W, Wang K, *et al.* Periodontitis and COVID-19: Immunological characteristics, related pathways, and association. *Int J Mol Sci* 2023; 24(3): 3012. [http://dx.doi.org/10.3390/ijms24033012] [PMID: 36769328]
- [28] Srimaneepong V, Rokaya D, Thunyakitpisal P, Qin J, Saengkiattiyut K. Corrosion resistance of graphene oxide/silver coatings on Ni-Ti alloy and expression of IL-6 and IL-8 in human oral fibroblasts. *Sci Rep* 2020; 10(1): 3247. [http://dx.doi.org/10.1038/s41598-020-60070-x] [PMID: 32094428]
- [29] Karki R, Sharma BR, Tuladhar S, *et al.* Synergism of TNF- α and IFN- γ triggers inflammatory cell death, tissue damage, and mortality in SARS-CoV-2 infection and cytokine shock syndromes. *Cell* 2021; 184(1): 149-168.e17. [http://dx.doi.org/10.1016/j.cell.2020.11.025] [PMID: 33278357]