

Get More with  
SINTA Insight

Go to Insight



## PADJADJARAN JOURNAL OF DENTISTRY

UNIVERSITAS PADJADJARAN

✱ P-ISSN : 19790201 < > E-ISSN : 25496212 📁 Subject Area : Health, Science



0.75  
Impact Factor



1226  
Google Citations



Sinta 2  
Current  
Accreditation

🔗 [Google Scholar](#)

🔗 [Garuda](#)


🌐 [Website](#)

🌐 [Editor URL](#)

### History Accreditation

2018 2019 2020 2021 2022 2023 2024 2025 2026

### Citation Per Year By Google Scholar



### Journal By Google Scholar

	All	Since 2018
Citation	1226	1135
h-index	13	12
i10-index	24	22

## Garuda

## Google Scholar

### [Antibacterial potential of celery leaf extract toothpaste on the growth of Streptococcus mutans ATCC 25175](#)

Faculty of Dentistry Universitas Padjadjaran 📖 [Padjadjaran Journal of Dentistry Vol 35, No 1](#)  
(2023): March 2023 1-4

📅 2023 📄 DOI: 10.24198/pjd.vol35no1.30624 🔄 Accred : Sinta 2

### [Effect of varnish coating agent application on surface hardness of glass ionomer cement after immersion in carbonated drinks](#)

Faculty of Dentistry Universitas Padjadjaran 📖 [Padjadjaran Journal of Dentistry Vol 35, No 1](#)  
(2023): March 2023 47-52

📅 2023 📄 DOI: 10.24198/pjd.vol35no1.33139 🔄 Accred : Sinta 2

### [The effectiveness of using local exhaust ventilation, HEPA filter, and dental aerosol suction on indoor air quality](#)

Faculty of Dentistry Universitas Padjadjaran 📖 [Padjadjaran Journal of Dentistry Vol 35, No 1](#)  
(2023): March 2023 23-28

📅 2023 📄 DOI: 10.24198/pjd.vol35no1.33566 🔄 Accred : Sinta 2

### [Relationship between mother's oral health literacy level with oral hygiene behavior and self-reported oral health status in sociodemographic scope](#)

Faculty of Dentistry Universitas Padjadjaran 📖 [Padjadjaran Journal of Dentistry Vol 35, No 1](#)  
(2023): March 2023 40-46

📅 2023 📄 DOI: 10.24198/pjd.vol35no1.35411 🔄 Accred : Sinta 2

### [Effectiveness of mixed tea plant extract gel with chitosan on fibroblast cells after tooth extraction of Wistar rats](#)

Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1](#)  
(2023): March 2023 10-15  
📅 2023    📄 DOI: 10.24198/pjd.vol35no1.36563    🏷️ Accred : Sinta 2

[Prevention and control of COVID-19 infection at the community health center during the new normal era](#)

Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1](#)  
(2023): March 2023 70-79  
📅 2023    📄 DOI: 10.24198/pjd.vol35no1.39563    🏷️ Accred : Sinta 2

[Relationship between oral health service quality and patient satisfaction at community health centers in Kupang city during COVID-19 pandemic](#)

Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1](#)  
(2023): March 2023 86-90  
📅 2023    📄 DOI: 10.24198/pjd.vol35no1.40111    🏷️ Accred : Sinta 2

[The effect of xerostomia on the quality of life of controlled and uncontrolled type 2 diabetes mellitus patients](#)

Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1](#)  
(2023): March 2023 5-9  
📅 2023    📄 DOI: 10.24198/pjd.vol35no1.41996    🏷️ Accred : Sinta 2

[Aloe vera extract wound healing sheet and free-range chicken egg albumin \(gallus domesticus\) accelerate angiogenesis on gingival incision wound in rats](#)

Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1](#)  
(2023): March 2023 16-22  
📅 2023    📄 DOI: 10.24198/pjd.vol35no1.42775    🏷️ Accred : Sinta 2

[The association of cariogenic foods consumption frequency on the occurrence of dental caries in children with autism spectrum disorders](#)


Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1](#)  
(2023): March 2023 34-39  
📅 2023    📄 DOI: 10.24198/pjd.vol35no1.42892    🏷️ Accred : Sinta 2

[View more ...](#)

Get More with  
SINTA Insight

[Go to Insight](#)

Citation Per Year By Google Scholar



Journal By Google Scholar

	All	Since 2018
Citation	1226	1135
h-index	13	12
i10-index	24	22



e ISSN 2549-6212  
p ISSN 1979-0201

# PADJADJARAN JOURNAL OF DENTISTRY

<http://jurnal.unpad.ac.id/pjd>

**Padjadjaran  
J. Dent**

**Vol. 34**

**No. 3**

**P. 184-309**

**Bandung  
November 2022**

e ISSN 2549-6212  
p ISSN 1979-0201

# PADJADJARAN JOURNAL OF DENTISTRY

<http://jurnal.unpad.ac.id/pjd>

Padjadjaran  
J. Dent

Vol. 30

No. 1

P. 138-208

Bandung  
March 2018

Accredited  
by





HOME LOGIN REGISTER SEARCH CURRENT ARCHIVES ANNOUNCEMENTS ABOUT

Home > About the Journal > Editorial Team

## Chief Editor

- Prof Sunardhi Widyaputra, drg, MS, PhD, Scopus ID= 6602995626; Department of Oral Biology, Faculty of Dentistry Universitas Padjadjaran, Indonesia

## Handling Editor

- Prof. Dr. Nina Djustiana, drg, MKes, Scopus ID= 57189578833; Department of Dental Materials, Science, and Technology, Faculty of Dentistry Universitas Padjadjaran, Indonesia
- Dr Anne Agustina Suwargiani, drg, MKM, Scopus ID= 57203020093; Department Ilmu Kesehatan Gigi Masyarakat, Fakultas Kedokteran Gigi Universitas Padjadjaran, Indonesia
- Zulia Hasratiningsih, drg, MDSc, Scopus ID= 37045476800; Departemen Ilmu Teknologi dan Material Kedokteran Gigi, Fakultas Kedokteran Gigi Universitas Padjadjaran, Indonesia, Indonesia
- Dr. Netty Suryanti, drg, MARS., Scopus ID= 57210117266; Department of Community Dental Health, Faculty of Dentistry, Padjadjaran University, Indonesia

## Editorial Board

- Dr. Ali Mohammed, Scopus ID= 57652411300; Faculty of Medicine, Dentistry and Health Sciences, The University of Melbourne, Australia, Australia
- Cortino Sukotjo, DDS, PhD, MMSc, FACP, Scopus ID= 6508194317; Department of Restorative Dentistry and Advanced Prosthodontics, College of Dentistry, University of Illinois, United States
- Prof. Mariko Naito, Scopus ID= 57204325415; Department of Oral Epidemiology, Graduate School of Biomedical and Health Sciences Hiroshima University, Japan
- Drg Rizky Indrameikha Sugianto, MPH, PhD, Scopus ID= 57201006215, Postdoctoral Researcher, Hannover Medical School,, Germany
- drg. Niekla Survia Andiesta, BDS, MDS, Scopus ID= 57202599268; Division of Children and Community Oral Health, School of Dentistry, International Medical University, Malaysia
- Prof. Mohamed Ebrahim Parker, Scopus ID= 7403672513; Department of Diagnostic Sciences Radiology, Maxillofacial and Forensic Sciences, University of The Western Cape, South Africa
- Prof. Kotaro Tanimoto, Scopus ID= 57191990083; Graduate School of Biomedical and Health Sciences Dentistry & Oral Health Sciences, Hiroshima University, Japan
- Prof. Dr. Mohammad Tariqur Rahman, Scopus ID = 55457946600, Dean Office Faculty of Dentistry, universiti malaya, Malaysia
- Prof. Yoshizo Matsuka, Scopus ID= 7003862097; Department of Stomatognathic Function and Occlusal Reconstruction, Graduate School of Biomedical Sciences Tokushima University, Japan
- Prof. Dr. Zamros Yuzadi Mohd Yusof, Scopus ID = 22939737100, Department of Community Oral Health & Clinical Prevention Faculty of Dentistry, Universiti Malaya, Malaysia, Malaysia
- Associate Professor Dr Akram Hassan, Scopus ID= 55832848700; Department of Periodontics, School of Dental Sciences Universiti Sains Malaysia, Malaysia
- Dr. Solachuddin Jauhari Arief, DDS., PhD. Ichwan, Scopus ID= 6504103591; PAPRSB Institute of Health Sciences, International Islamic University, Brunei Darussalam
- Prof. Dr. Arlette Suzy Puspa Pertiwi Setiawan, drg., Sp KGA., M.Si., Scopus ID= 56044838600; Department of Pediatric, Faculty of Dentistry Padjadjaran University,, Indonesia
- Dr Arief Cahyanto, MT., Ph.D, Scopus ID= 55532851800; Department of Dentistry Material Science and Technology, Faculty of Dentistry, Padjadjaran University, Indonesia, Indonesia
- Dr Elizabeth Fitriana Sari, Scopus ID= 57219228212 Departemen Ilmu Penyakit Mukut, Fakultas Kedokteran Gigi Universitas Padjadjaran,, Indonesia
- Dr. Rasmi Rikmasari, drg., Sp.Pro., Subsp. OGST (K), Scopus ID= 57191990083; Department of Prosthodontics, Faculty of Dentistry, Padjadjaran University, Indonesia, Indonesia
- Dr. Sri Tjahajawati, drg., M.Kes.AIFM., Scopus ID= 57197722254; Department of Oral Biology, Faculty of Dentistry Universitas Padjadjaran, Indonesia

## MANUSCRIPT TEMPLATE



## FOCUS AND SCOPE

## AUTHOR GUIDELINES

## PEER REVIEW

## PUBLICATION ETHICS

## SCREENING FOR PLAGIARISM

## EDITORIAL BOARD

## REVIEWER

## COPYRIGHT TRANSFER FROM

## AUTHOR FEES

## JOURNAL HISTORY

## ARCHIVES

## ONLINE SUBMISSION

## CONTACT

## ACCOLADE



## VISITORS

ID	57257	CN	6181
US	4451	IN	4119
PH	2159	MY	1335
TH	734	EG	674
TR	673	GB	588
Newest:	GD	You:	IN
Today:			12
Month:			334
Total:			90665
			Supercounters.com

## MAPS



- **Dr. Hendra Dian Adhita Dharsono**, drg, Sp.KG., Subsp.KE[K], Scopus ID= 57204917449; Department of Dental Conservation, Faculty of Dentistry, Padjadjaran University, Indonesia
- **Fahmi Oscandar**, drg., M.Kes., SpOF., SubSp OFK (K), Ph.D. (Cr.Img), Scopus ID= 57199734614; Departemen Radiologi Kedokteran Gigi, Fakultas Kedokteran Gigi, Universitas Padjadjaran, Indonesia
- **Dr. Endang Sjamsudin**, drg, Sp.BMMF, Subsp.TMF-TMJ[K], Scopus ID= 57192257503; Department of Oral Surgery, Faculty of Dentistry, Padjadjaran University, Indonesia
- **R. Tantry Maulina**, drg., M.Kes., Ph.D., Scopus ID= 57191972242; Department of Oral Surgery, Faculty of Dentistry, Padjadjaran University, Indonesia, Indonesia
- **Dr. Avi Laviana**, drg., Sp.Ort., Subsp.DDTK(K), Scopus ID= 57211331865; Doctor of Orthodontics Department of Orthodontics Faculty of Dentistry, Padjadjaran University, Indonesia
- **Prof. Dr. Irna Sufiawati**, drg., Sp.PM., Subsp.Inf[K], Scopus ID= 56081844700; Departemen Ilmu Penyakit Mukut, Fakultas Kedokteran Gigi Universitas Padjadjaran, Indonesia, Indonesia
- **Amaliya**, drg., M.Sc., Ph.D., Scopus ID= 56584444300; Department of Periodontics, Faculty of Dentistry, Padjadjaran University, Indonesia
- **Lusi Epsilawati**, drg., Sp.RKG., Subsp.Rad.P(K), M.Kes., Scopus ID= 55523245700; Department of Radiology, Faculty of Dentistry, Padjadjaran University, Indonesia
- **Vita Mulya Passa Novianti**, drg., Sp.Pros., Scopus ID= 57217103247; Department of Prosthodontics, Faculty of Dentistry, Padjadjaran University, Indonesia
- **Aldilla Miranda**, drg., Sp.Perio[K], Scopus ID= 57205063638; Department of Periodontics, Faculty of Dentistry Universitas Padjadjaran, Indonesia

## Managing Editor

- **Siti Mariam**, Orcid ID: 0000-0003-0304-6875, Administrasi Jurnal Kedokteran Gigi Universitas Padjadjaran, Unit Publikasi Ilmiah Fakultas Kedokteran Gigi Universitas Padjadjaran, Indonesia
- **Hari Muhdori**, Orcid ID: 0000-0001-8263-4637; Administrasi Jurnal Padjadjaran Journal of Dentla Researchers and Students, Unit Publikasi Ilmiah, Fakultas Kedokteran Gigi Universitas Padjadjaran, Indonesia
- **Robby Wahyu Akbar**, Orcid ID: 0009-0008-1554-6221; Administrasi Jurnal Padjadjaran Journal of Dentistry, Fakultas Kedokteran Gigi Universitas Padjadjaran, Indonesia



00554361 Visitor Stat



Padjadjaran Journal of Dentistry is licensed under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/)



## RECOMMENDED APPS



## USER

Username

Password

☐ Remember me

Login

PAuS Login

## JOURNAL CONTENT

Search :

Keywords...

Search Scope

All

Search

## Browse

- By Issue
- By Author
- By Title
- Other Journals
- Categories

## KEYWORDS

E. faecalis Indonesia Oral health care  
P. gingivalis Porphyromonas  
gingivalis S. mutans S. sanguinis  
Streptococcus mutans  
Streptococcus sanguinis aloe  
vera antibacterial activity  
attitude calcium children dental  
hypnosis knowledge oral health  
periodontitis plaque index  
practice wound healing



## Content of this issue

### Volume 34, Number 3, November 2022

1.	Flora Ng Lai Yi Eriska Riyanti Meirina Gartika	The effect of acidulated phosphate fluoride (APF) gel against erosion on enamel microstructure of primary teeth which soaked with <i>Citrus aurantifolia</i>	184-191
2.	Saskia Lenggogeni Nasroen Florence Meliawaty Tichvy Tammama Putri Indriani Prasdiaz Tias Choirunnisa Ani Melani Maskoen	Analysis of FOXE1 rs4460498 and GSTP-1 I105V associated with non syndromic cleft lip and palate among Deutero Malay Subrace in Indonesia	192-201
3	Felisha Febriane Balafif Mieke Hemiawati Satari Diah Dhianawaty	Antifungal effect of ethyl acetate fraction of <i>Sarang semut</i> ( <i>Myrmecodia pendens</i> Merr.& Perry) against <i>Candida albicans</i>	202-209
4	Muhamad Tio Dwi Hertanto Wahyu Hidayat Netty Suryanti	Implementation of community dental and oral health care before COVID-19 pandemic to new normal period at Public Health Care	210-222
5	Meta Maulida Damayanti Ajeng Kartika Sari Annisa Rahmah Furqaani	Effectiveness of natural lip balm cinnamon ( <i>Cinnamomum burmannii</i> ) in accelerating the incision wound healing process in <i>rattus norvegicus</i>	223-231
6	Indah Puti Rahmayani Sabirin Amanda Debora Polii Rahmadaniah Khaerunnisa Euis Reni Yuslianti Ris Kristiana	Effect of 20% ethanol extract of Subang pineapple gel on oral mucosa incisional wound length	232-238
7	Megananda Hiranya Putri Euis Julaeah Eliza Herijulianti Neneng Nurjanah	Formulation of Mouthwash for Gingivitis from Combination Infusion of Salam leaves ( <i>Eugenia Polyantha</i> Wight) and Betel leaf ( <i>Piper betle</i> . L)	239-248
8	Amy Nindia Carabelly Maharani Lallyza Apriasari Irnamanda Dwipura Selviana Rizky Pramitha Viranda Sutanti Annisa Kurniasari Riznasyarielia Nikmatun Nafisah	Correlation between presence of bacillus species and OHI-s in recurrent aphthous stomatitis patients	249-254
9	Alisha Zuchraini Pandji Yurika Ambar Lita Winny Yohana	Correlation between human chronological age and dentin volume of maxillary canine	255-263
10	Wiwiek Poedjiastoeti Muhamad Novo Perwira Lubis Yessy Ariesanti Intan Farizka Jackson Dipankara Samroeng Ingiam	Alveolar bones density assessment of dental implant sites using cone-beam computed tomography	264-270
11	Muhamat Muhtar S. Abdurrohman Rahmawati Sri Praptiningsih Yulistinawati Yulistinawati	The effect of brown algae ( <i>Sargassum</i> sp.) gel on the number of osteoclasts in periodontitis rats	271-276





**Content of this issue**  
**Volume 34, Number 3, November 2022**

12	Munifah Abdat Dani Laulea Fitri Talat Hasan Al-Gunaid	Positive Correlation Between Maternal Self-Efficacy with The Children Oral Hygiene Behavior	277-283
13	Trimurni Abidin Gieska Lailarahma Darmawan Fatimah Dewi Dalimunthe	The effect of oligosaccharide chitosan irrigating agent on penetration of two type endodontic sealers into the dentinal tubules: a confocal laser scanning microscopy study	284-292
14	Syahla Fatimah Azzahra Anne Agustina Suwargiani Anna Muryani	Differences in the motivation for seeking oral health treatment among parents of kindergarten students	293-301
15	Endang Sjamsudin Arina Al Khaq Abel Tasman Yuza	Distribution frequency of Post Odontectomy Complications under General Anesthesia in Universitas Padjadjaran Dental and Oral Hospital (RSGM Unpad) Bandung	302-309

## Alveolar bones density assessment of dental implant sites using cone-beam computed tomography

Wiwiek Poedjiastoeti<sup>1\*</sup>, Muhamad Novo Perwira Lubis<sup>2</sup>, Yessy Ariesanti<sup>1</sup>,  
Intan Farizka<sup>2</sup>, Jackson Dipankara<sup>1</sup>, Samroeng Inqlam<sup>3</sup>

<sup>1</sup>Departement of Oral and Maxillofacial Surgery, Faculty of Dental Medicine, Universitas Trisakti, Jakarta, Indonesia

<sup>2</sup>Departement of Oral and Maxillofacial Radiology, Faculty of Dental Medicine, Universitas Trisakti, Jakarta, Indonesia

<sup>3</sup>Department of Oral and Maxillofacial Radiology, Faculty of Dentistry, Thammasat University, Pathumtani, Bangkok, Thailand

### ABSTRACT

**Introduction:** a preoperative assessment of bone density plays a vital role in the success of dental implant treatment. the maxilla and mandibular alveolar bone had a variety of bone densities. Therefore, dental imaging is an important step before placing a dental implant. Recently, Cone beam computed tomography (CBCT) is widely used in dental medicine and also recommended by AAOMR for preoperative implant placement. The aims of this study is to analyzed the alveolar bone density of the dental implant sites for dental implant planning using CBCT. **Methods:** ninety-three CBCT data were retrieved from the database of the department of oral and maxillofacial radiology at Dental Hospital Faculty of Dentistry Universitas Trisakti and examined. The recipient sites for dental implant placement were determined based on CBCT data using implant planning software (i-Dixel). The alveolar bones value is recorded in grayscale value (GV). **Results:** a great variety of alveolar bone density was observed ranging from 134-891 GV. One-way ANOVA was conducted, and statistically significant was only shown at the anterior mandibular region and molar mandibular region. However, no statistical differences were observed when comparing the male and female groups using the T-test. **Conclusion:** Critical evaluation through dental CBCT can be made before dental implant placement in the alveolar bones. the lowest alveolar bone density was observed in maxillary molar dental implant sites with mean 322 GV and highest alveolar bone density was shown in mandibular anterior dental implant sites with mean 558.8 GV. Determining the bone density at the dental implant site before dental implant placement is crucial. Primary stability and secondary stability is determined by bone density. Therefore, higher value of alveolar bone density is needed for dental implant placement.

**Keywords:** alveolar bone density assessment; dental implant; cone-beam computed tomography.

p-ISSN: 1979-0201; e-ISSN: 2549-6212; Available from: <http://jurnal.unpad.ac.id/pid/article/view/42383>

DOI: [10.24198/pjd.vol34no3.42383](https://doi.org/10.24198/pjd.vol34no3.42383)

Submission: Oct 12, 2022; Accepted: Nov 30, 2022; Published online: Nov 30, 2022

---

*\*Corresponding author: Wiwiek Poedjiastoeti, Department of Oral and Maxillofacial Surgery, Faculty of Dental Medicine, Universitas Trisakti, Jl. Kyai Tapa No. 260, Jakarta, 11440, e-mail: [wiwiek@trisakti.ac.id](mailto:wiwiek@trisakti.ac.id), Phone: 021-5655786*

## INTRODUCTION

Tooth loss has been a problem for humans over the centuries. Since the 19<sup>th</sup> century, Dental implant has been used to replace missing teeth.<sup>1</sup> To date, the dental implant has been widely used because of its advantages, such as more protection for the surrounding teeth and the alveolar bone. On the other hand, rehabilitation treatment using dental implants has been reported to increase mastication function and quality of life.<sup>2</sup> Therefore, a dental implant can be an essential treatment option for replacing missing teeth.<sup>3</sup>

Dental implants have a wide variety of success rates in dentistry. The quantity and quality of bone availability are the primary determinants in predicting the success rate of the dental implant. However, different sites in the jaw have different success rates.<sup>4</sup> The functional quality of alveolar bones is determined by bone density. Bone density has a critical effect on surgical technique, healing time, and time of loading during prosthodontic reconstructions.<sup>5,6</sup> The golden site for dental implant placement is in the site with high bone density. Otherwise, the success rate decreases when the bone has a lower density or is porous.<sup>4</sup> Assessing the bone tissue has two objectives: first, is the bone tissue sufficient for dental implant placement (as a diagnostic tool), and second, to predict the success rate of the dental implant (prognostic tool).<sup>7</sup>

Poor bone density is considered a risk factor for biological complications of the implant, associated with lack of primary stability and impaired healing or osseointegration, which can lead to early implant loss.<sup>7</sup> There are two strategies to evaluate bone quality and quantity. The direct estimation strategy incorporates ex vivo examinations (i.e., dry skulls or corpses) or sample/biopsy recovered for examination from creatures or human subjects and in vivo examinations done on living subjects.

The indirect estimation strategy depends on radiographic imaging, like registered tomography (CT) or cone-beam computed tomography (CBCT). These estimation methods give a three-dimensional depiction of bony structures. They are viewed as a precise diagnostic instrument that enables the assessment of the morphology, bone quality, and volume of the leftover alveolar ridge.<sup>8</sup>

The American Academy of Oral and Maxillofacial Radiology (AAOMR) currently advocates using sectional imaging for all dental implant treatment.<sup>9</sup> CBCT provided volumetric data of the jaw with relatively low radiation doses and costs with high resolution.<sup>6,10</sup> Overall dental implant placement (implant size, angle, and type) can directly influence the alveolar bone quantity. On the other hand, the stability of dental implants is associated with bone quality. Therefore, a comprehensive assessment of both factors is essential.<sup>11</sup> Ivanova et al.<sup>12</sup> Concluded primary stability and secondary stability is determined by bone density.<sup>12</sup> Baltayan et al.<sup>7</sup> found a strong correlation between alveolar bone density and primary stability.<sup>13</sup>

Poor bone quality associated with lack of primary stability.<sup>7</sup> Therefore, to better understand the alveolar bone density of the edentulous area for dental implant planning using CBCT, we studied the edentulous area of the implant site on 93 patients who underwent preoperative evaluation using CBCT in our department in January 2018 - December 2021. The aims of this study is to analyzed the alveolar bone density of the dental implant sites for dental implant planning using CBCT.

## METHODS

In a cross-sectional study, ninety-three CBCT data of patients randomly selected from the department of oral and maxillofacial radiology database at Dental Hospital Faculty of Dentistry Universitas trisakti (RSGMP-FKG Usakti) between January 2018 - December 2021 was evaluated. The CBCT data must meet the following inclusion criteria: CBCT data in patients aged a minimum of 20 years old and the radiograph taken between January 2018-December 2021, edentulous areas with the potential for dental implant placement in the jaw and the exclusion criteria is CBCT radiographs with poor quality.

Measurements Morita 3D Cone-Beam Computed Tomography (3D Accuitomo 170, J. Morita Co., Japan) was used to obtain patient 3D scan data at 90 KvP, 5 mA, and 80 µm voxel size exposure setting. Cross sectional view with 1.00 mm slice thickness was obtained. All CBCT data were evaluated by two calibrated oral and maxillofacial radiologists.




Figure 1. Bone density measurement. (A1) fixture simulation lower molar region (A2) bone density assessment lower molar region (A3) Sagittal view showing implant fixture simulation at lower molar region (A4) Coronal view lower molar region (B1) fixture simulation upper incisive region (B2) bone density assessment upper incisive region (B3) Sagittal view showing implant fixture simulation at upper incisive region (B4) Coronal view of upper incisive region.

Inter-observer agreement was calculated to verify the reliability of measurements. Bone density Head reorientation on CBCT images was undergone prior to bone density measurement. Bone density was measured using the software AIS-3D App by choosing an implant planning

feature build on the software. The next step was to determine the region of interest or tooth area to be measured (incisive, premolar, molar in both jaws) followed by implant insertion in the selected region, and the bone density mean value inside the implant volume will be calculated (Figure 1).

Table 1. Interpretation of Cohen's Kappa ( $\kappa$ )

Cohen's Kappa ( $\kappa$ )	Strength of Agreement
<0.00	Poor
0.01-0.20	Slight
0.21-0.40	Fair
0.41-0.60	Moderate
0.61-0.80	Substantial
0.81-1.00	Almost perfect

### Data analysis

We recorded the alveolar bone density data in Microsoft Excel then all statistical analyses conducted using SPSS Statistics for Windows V22 (IBM, USA). First, inter-observer reliability was observed using Cohen's Kappa ( $\kappa$ ) statistical analysis. The interpretation of Cohen's Kappa ( $\kappa$ ) can be seen in table 1.<sup>14</sup> Independent sample T-test was used to compare the mean difference between the male and female groups.

One-way ANOVA was used to compare the mean difference among age groups. Tukey post hoc analysis was used to compare all possible age group pairing when One-way ANOVA showed a significant difference. The study protocol was approved by the Komisi Etik Penelitian Kesehatan Fakultas Kedokteran Gigi Universitas Trisakti

(Ethical clearance committee, No 024/S3/KEPK/FKG/7/2022)

### RESULTS

Two observers evaluated the collected CBCT image data of 93 subjects. First, we divided the alveolar bones into maxillary and mandibular and then differentiated them into three regions: anterior, premolar, and molar.

The kappa coefficients of inter-observer A and B were 0.92, 0.91, and 0.87, respectively, for the maxillary anterior region, maxillary premolar region, and maxillary molar region, indicating almost perfect agreement. However, the kappa coefficients for the mandibular anterior region bone density

Table 2. Inter-observer kappa coefficients

The Alveolar bones regions	Observer A to B
Maxillary Anterior Region	0.92
Maxillary Premolar Region	0.91
Maxillary Molar Region	0.87
Mandibular Anterior Region	0.70
Mandibular Premolar Region	0.95
Mandibular Molar Region	0.90

The result showed that mean maxillary anterior region bone density was  $362.4 \pm 107.35$  GV, maxillary premolar region bone density was  $326.6 \pm 77.12$  GV, maxillary molar region

bone density was  $322.0 \pm 103.21$  GV, mandibular anterior, premolar, and molar region density were  $557.8 \pm 160.29$  GV,  $402.8 \pm 111.06$  GV, and  $408.6 \pm 95.87$  GV respectively (Table 3).

Table 3. The alveolar bones density

The Alveolar bones regions	Min	Max	Mean	Std. Deviation
Maxillary Anterior Region	195	587	362.4	107.35
Maxillary Premolar Region	172	582	326.6	77.12
Maxillary Molar Region	134	745	322.0	103.21
Mandibular Anterior Region	319	891	557.8	160.29
Mandibular Premolar Region	257	848	402.8	111.06
Mandibular Molar Region	173	681	408.6	95.87

Note: all result value describes in grayscale value (GV)

The distribution of male and female patients was 42 and 51 people. The mean maxillary anterior region bone density was 348.5 GV, 377.6 GV for males and females, and maxillary premolar region bone density was 313.8 GV, and 341.6 GV for males and females, maxillary molar region bone density was 296 GV and 342,4 GV. Therefore, the mandibular anterior region bone density was 518.5

and 656.2 for males and females, the mandibular premolar region bone density was 373.1 GV and 428.8 GV for males and females, and the mandibular molar region bone density was 407.6 GV and 409.5 GV for male and female. However, no statistically significant difference was observed when comparing the male and female group bone density (Table 4).

Table 4. Bone density analysis based on gender and age group.

The Alveolar bones Region	Mean $\pm$ SD (GV) based on gender		p	Mean $\pm$ SD (GV) based on Age Group			p
	Male	Female		21-45	46-60	>61	
Maxillary anterior region	348.5 $\pm$ 117.4	377.6 $\pm$ 98.4	0.52	314.5 $\pm$ 69.1	366.2 $\pm$ 122.4	440.6 $\pm$ 35.5	0.25
Maxillary premolar region	313.8 $\pm$ 79.2	341.6 $\pm$ 73.9	0.26	294.2 $\pm$ 73.4	329.6 $\pm$ 84.5	356 $\pm$ 53.6	0.21
Maxillary molar region	296 $\pm$ 62.8	324.4 $\pm$ 124.9	0.27	302.7 $\pm$ 31	343.1 $\pm$ 175.5	318.6 $\pm$ 55.7	0.76
Mandibular anterior region	515.5 $\pm$ 139.6	656.2 $\pm$ 186.4	0.15	324.5 $\pm$ 7.7	570.2 $\pm$ 150.5	649.7 $\pm$ 102.3	0.04*
Mandibular premolar region	373.1 $\pm$ 90.4	428.8 $\pm$ 123.3	0.17	389.7 $\pm$ 55.7	412.2 $\pm$ 154.9	399.5 $\pm$ 54.8	0.90
Mandibular molar region	407.6 $\pm$ 94.4	409.5 $\pm$ 98.9	0.94	451.8 $\pm$ 66	412.8 $\pm$ 105.1	350.3 $\pm$ 80	0.01*

Note : Grayscale Value (a number that represents the amount of attenuation of the X-ray beam by the material contained in each voxel or structural unit of the tomographic volume )

\*p<0.05 = statistically significant difference

The median age was 50.34, ranging from 21 to 82 years old. We divided the age group into three groups, ranging from 21-45 (young adult),

46-60 (middle adult), and >61 (elderly) according to WHO age group classification in 2015. The one-way ANOVA test was carried out.

Table 5. Multiple comparison using Tukey post hoc test

The Alveolar bones Region	Age Group	Age Group	p	95% Confidence Interval	
				Lower bound	Upper bound
Maxillary Anterior Region	21-45	46-60	0.581	-181.54	78.11
	21-45	>61	0.231	-314.31	61.98
	46-60	>61	0.518	-243.73	94.83
Maxillary Premolar Region	21-45	46-60	0.458	-197.33	36.43
	21-45	>61	0.194	-147.08	23.48
	46-60	>61	0.666	-100.85	48.15
Maxillary Molar Region	21-45	46-60	0.747	-178.83	98.01
	21-45	>61	0.951	-147.69	115.91
	46-60	>61	0.879	-102.34	151.39
Mandibular Anterior Region	21-45	46-60	0.088	-526.46	34.96
	21-45	>61	0.038*	-632.76	-17.73
	46-60	>61	0.599	-296.94	137.94
Mandibular Premolar Region	21-45	46-60	0.898	-148.47	103.54
	21-45	>61	0.984	-151.90	132.40
	46-60	>61	0.966	-113.29	138.72
Mandibular Molar Region	21-45	46-60	0.384	-31.61	109.51
	21-45	>61	0.012*	19.02	183.96
	46-60	>61	0.113	-11.38	136.46

\*p<0.05 statistically significant difference

No significant differences were found between the maxillary anterior region, maxillary premolar region, maxillary molar region, and mandibular premolar region ( $p=0.255$ ,  $p=0.216$ ,  $p=0.760$ , and  $p=0.903$ , respectively) compared to age groups (Table 4). However, the mandibular anterior and mandibular molar regions showed significant differences compared to age groups with  $p=0.45$  and  $p=0.16$ . The multiple comparisons using Tukey post hoc clarify the differences only seen in the young adult group compared to elderly groups in the mandibular anterior and molar region (Table 5).

## PEMBAHASAN

Bone density assessment is indispensable to achieving the best dental implant treatment planning and the primary stability of the dental implant.<sup>15</sup> Bone density is one predictor of successful dental implant treatment. Therefore, an accurate evaluation of the alveolar bone density is essential. CBCT is three-dimensional dental imaging widely used recently in dental medicine. One of the several advantages of CBCT dental imaging enables the assessment of bone density.<sup>9</sup> This study was conducted to assess the

alveolar bone density of dental implant sites using the dental CBCT. A higher value of inter-observer reliability was observed in our study at table 2. Nevertheless, the mandibular anterior region has shown substantial agreement. This study's decrease in inter-observer reliability is thought to be because the available data for the mandibular anterior region is limited.

The result is that a slight difference in the assessment significantly influences the reliability. In line with the previous study, the agreement value of our result ranges from 0.70-0.95, and the previous study was 0.61-0.92.<sup>16</sup> Similar result was seen in a study conducted by Alkhader et al.<sup>17</sup> they showed the Cohen's kappa value was range from 0.77-0.93. The reliability of inter-observer reliability can be improved by practice and training in diagnostic imaging.<sup>14</sup>

The alveolar bone density was greater in females compared to males (Table 4). However, no statistically significant differences between male and female alveolar bone density were found. No data available from the previous study can be compared to our study. This is due to our CBCT data being recorded in grayscale value. Therefore, we try to compare the result with available data recorded in Hounsfield Unit (HU)



using Computed Tomography (CT). This decision was made based on a study that provided a strong correlation between GV of CBCT and HU of CT.<sup>18</sup> In contrast to our study, Cassetta et al.<sup>19</sup> found a statistically significant difference in alveolar bone density between gender, and they concluded that the density of the male alveolar bone was greater than that of females.<sup>19</sup>

A study conducted by Kim et al. was in line with Cassetta result; they found a statistical difference in the alveolar bone density between gender. However, in the study by Kim et al.<sup>20</sup> they showed a greater alveolar bone density was seen in females compared to males. Several risk factors may affect the density of the alveolar bone, such as gender, periodontal disease, osteoporosis,<sup>21</sup> tooth loss, menopause, and age.<sup>22</sup>

Our study showed that a great variety of alveolar bone density was found in the maxilla or mandibular. Statistically significant differences between age and alveolar bone density were observed in young adults compared to the elderly. Based on our result, age seems to affect alveolar bone density (Table 4). Other authors found different results that report that age does not affect the alveolar bone density value.<sup>21</sup>

Previously three-dimensional alveolar bones were evaluated using CT, including the alveolar bones density assessment. Consider the patient who received x-ray exposure. CT is not ideal for some diagnostic purposes in dental medicine due to the high dose of radiation, poor resolution, longer scanning time, difficult interpretation, high-cost charge, and limited availability.<sup>23</sup> To overcome the limitation, Arai et al.<sup>24</sup> (1999) in Japan and Mozzo et al.<sup>25</sup> (1998) in Italy at introduced cone-beam computed tomography (CBCT) for oral and maxillofacial application. The CBCT fulfils the needs, such as reducing unnecessary exposure to the patient, high resolution, rapid scan time, dose reduction, and reduced image artefact.<sup>26</sup>

The alveolar bone density plays a vital role in the mechanical immobilization of the dental implant during healing, distribution, and transition of stresses from the prosthesis to the implant-bone interface after healing. The availability of alveolar bone density determines the dental implant design, surgical approach, healing time, and initial progressive loading during prosthetic construction.<sup>27</sup> With the reason and data provided

above, preoperative evaluation of the alveolar bone density prior to dental implant placement at the edentulous area planned to a received dental implant is mandatory.

The limitation of our study is we did not compare the alveolar bone density based on the left and right regions of the jaw and lack of data available from our database for the mandibular anterior region. Hopefully, in the future, a study could be conducted to overcome the limitation.

## CONCLUSION

The highest alveolar bone density value was found on the mandibular anterior region, followed by the mandibular molar, mandibular premolar, maxillary anterior, maxillary premolar and maxillary molar regions. These conditions were noted in both gender and age groups. In the age group above 61 years old, the highest density of alveolar bone differences was found in the mandibular anterior region; however, the lowest density was noted in the mandibular molar region. The grayscale value in CBCT can be used to determine the alveolar bone density of dental implant sites.

## REFERENCES

1. Alghamdi HS, Jansen JA. The development and future of dental implants. *Dent Mater J*. 2020; 39(2): 167-172. DOI: [10.4012/dmj.2019-140](https://doi.org/10.4012/dmj.2019-140).
2. Elani HW, Starr JR, Da Silva JD, Gallucci GO. Trends in dental implant use in the U.S., 1999-2016, and projections to 2026. *J Dent Res*. 2018; 97(13): 1424-30. DOI: [10.1177/0022034518792567](https://doi.org/10.1177/0022034518792567).
3. Sulijaya B, Kuswandani SO, Soeroso Y. Contemporary guided bone regeneration therapy for unaesthetic anterior peri-implantitis case. *Dent J*. 2016; 49(4): 181-4. DOI: [10.20473/j.djmk.v49.i4.p181-184](https://doi.org/10.20473/j.djmk.v49.i4.p181-184).
4. Attar BM, Alaei S, Badrian H, Davoudi A. Clinical and radiological evaluation of implants placed with osteotome sinus lift technique: 19-month follow-up. *Ann Maxillofac Surg*. 2016; 6(2): 190-4. DOI: [10.4103/ams.ams.7.16](https://doi.org/10.4103/ams.ams.7.16).
5. Arina YMDa, Ferdiansyah F, Rubianto M. The evaluation of mandibular bone density in chronic periodontitis models. *Dent J*. 2018;

- 51(4): 210-5. DOI: [10.20473/j.djmk.v51.i4.p210-215](https://doi.org/10.20473/j.djmk.v51.i4.p210-215)
6. Eskandarloo A, Arabi R, Bidgoli M, Yousefi F, Poorolajal J. Association between marginal bone loss and bone quality at dental implant sites based on evidence from cone beam computed tomography and periapical radiographs. *Contemp Clin Dent*. 2019; 10(1): 36-41. DOI: [10.4103/ccd.ccd.185.18](https://doi.org/10.4103/ccd.ccd.185.18).
7. Chrcanovic BR, Albrektsson T, Wennerberg A. bone quality and quantity and dental implant failure: a systematic review and meta-analysis. *Int J Pros*. 2017; 30(3): 219-37. DOI: [10.11607/ijp.5142](https://doi.org/10.11607/ijp.5142).
8. Fokas G, Vaughn VM, Scarfe WC, Bornstein MM. Accuracy of linear measurements on CBCT images related to presurgical implant treatment planning: a systematic review. *Clin Oral Implants Res*. 2018; 29(16): 393-415. DOI: [10.1111/clr.13142](https://doi.org/10.1111/clr.13142).
9. Weiss R 2nd, Read-Fuller A. Cone beam computed tomography in oral and maxillofacial surgery: an evidence-based review. *Dent J (Basel)*. 2019; 7(2): 52. DOI: [10.3390/dj7020052](https://doi.org/10.3390/dj7020052).
10. Jacobs R, Salmon B, Codari M, Hassan B, Bornstein MM. Cone beam computed tomography in implant dentistry: recommendations for clinical use. *BMC Oral Health*. 2018; 18(1): 88. DOI: [10.1186/s12903-018-0523-5](https://doi.org/10.1186/s12903-018-0523-5).
11. Wakimoto M, Matsumura T, Ueno T, Mizukawa N, Yanagi Y, Iida S. Bone quality and quantity of the anterior maxillary trabecular bone in dental implant sites. *Clinical oral implants research*. 2012; 23(11): 1314-9. DOI: [10.1111/j.1600-0501.2011.02347.x](https://doi.org/10.1111/j.1600-0501.2011.02347.x).
12. Igarashi C, Theramballi YG, Kobayashi K. Inter-observer reliability in cone-beam computed tomography assessment of the retromolar canal: A practical plan to improve diagnostic imaging. *Imaging Sci Dent*. 2022; 52(2): 181-6. DOI: [10.5624/isd.20210289](https://doi.org/10.5624/isd.20210289).
13. Sreerama R, Kolluru KC, Gottumukkala V, Innampudi CK, Konathala JR, Krishnaveni G. Assessment of the effect of bone density on implant stability: a clinical study. *J Pharm Bioallied Sci*. 2021; 13(1): S297-S300. DOI: [10.4103/jpbs.JPBS.794.20](https://doi.org/10.4103/jpbs.JPBS.794.20).
14. Kim JH, Abdala-Junior R, Munhoz L, Cortes ARG, Watanabe PCA, Costa C et al. Comparison between different cone-beam computed tomography devices in the detection of mechanically simulated peri-implant bone defects. *Imaging Sci Dent*. 2020; 50(2): 133-9. DOI: [10.5624/isd.2020.50.2.133](https://doi.org/10.5624/isd.2020.50.2.133)
15. Alkhader M, Hudieb M, Khader Y. Predictability of bone density at posterior mandibular implant sites using cone-beam computed tomography intensity values. *Eur J Dent*. 2017; 11(3): 311-6. DOI: [10.4103/ejd.ejd.14.17](https://doi.org/10.4103/ejd.ejd.14.17)
16. Razi T, Niknami M, Alavi Ghazani F. Relationship between Hounsfield Unit in CT scan and gray scale in CBCT. *J Dent Res Dent Clin Dent Prospects*. 2014; 8(2): 107-10. DOI: [10.5681/joddd.2014.019](https://doi.org/10.5681/joddd.2014.019).
17. Cassetta M, Sofan AA, Altieri F, Barbato E. Evaluation of alveolar cortical bone thickness and density for orthodontic mini-implant placement. *J Clin Exp Dent*. 2013; 5(5): 245-52. DOI: [10.4317/jced.51228](https://doi.org/10.4317/jced.51228)
18. Kim JH, Lim YJ, Kim B, Lee J. How do parameters of implant primary stability correspond with ct-evaluated bone quality in the posterior maxilla? A correlation analysis. *Materials (Basel)*. 2021; 14(2): 270. DOI: [10.3390/ma14020270](https://doi.org/10.3390/ma14020270)
19. Yu B, Wang CY. Osteoporosis and periodontal diseases - An update on their association and mechanistic links. *Periodontol 2000*. 2022; 89(1): 99-113. DOI: [10.1111/prd.12422](https://doi.org/10.1111/prd.12422).
20. Takaishi Y, Arita S, Honda M. Assessment of alveolar bone mineral density as a predictor of lumbar fracture probability. *Adv Ther*. 2013; 30(5): 487-502. DOI: [10.1007/s12325-013-0028-1](https://doi.org/10.1007/s12325-013-0028-1)
21. Saati S, Kaveh F, Yarmohammadi S. Comparison of cone beam computed tomography and multi slice computed tomography image quality of human dried mandible using 10 anatomical landmarks. *J Clin Diagn Res*. 2017; 11(2): ZC13-ZC16. DOI: [10.7860/JCDR/2017/20637.9253](https://doi.org/10.7860/JCDR/2017/20637.9253).
22. Venkatesh E, Elluru SV. Cone beam computed tomography: basics and applications in dentistry. *J Istanbul Univ Fac Dent*. 2017; 51(3 Suppl 1): S102-S121. DOI: [10.17096/jiufd.00289](https://doi.org/10.17096/jiufd.00289).
23. Sirisha Attili HS, Sitaram Prasad Kasina VHC Kumar, Srilatha Balusu, Sarat CB. To evaluate the bone mineral density in mandible of edentulous patients using computed tomography: an in vivo study. *J Int Oral Health*. 2015; 7(4): 22-6.

# Alveolar bones density assessment of dental implant sites using cone-beam computed tomography

*by* M Novo Perwira Lubis FKG

---

**Submission date:** 04-Dec-2023 02:05PM (UTC+0700)

**Submission ID:** 2247188315

**File name:** bones\_density\_assessment\_of\_dental\_implant\_sites\_using\_CBCT.pdf (254.1K)

**Word count:** 4118

**Character count:** 21575

## Alveolar bones density assessment of dental implant sites using cone-beam computed tomography

Wiwiek Poedjiastoeti<sup>1\*</sup>, Muhamad Novo Perwira Lubis<sup>2</sup>, Yessy Ariesanti<sup>1</sup>,  
Intan Farizka<sup>2</sup>, Jackson Dipankara<sup>1</sup>, Samroeng Ingiam<sup>3</sup>

<sup>1</sup>Departement of Oral and Maxillofacial Surgery, Faculty of Dental Medicine, Universitas Trisakti, Jakarta, Indonesia

<sup>2</sup>Departement of Oral and Maxillofacial Radiology, Faculty of Dental Medicine, Universitas Trisakti, Jakarta, Indonesia

<sup>3</sup>Department of Oral and Maxillofacial Radiology, Faculty of Dentistry, Thammasat University, Pathumtani, Bangkok, Thailand

### ABSTRACT

**Introduction:** a preoperative assessment of bone density plays a vital role in the success of dental implant treatment. the maxilla and mandibular alveolar bone had a variety of bone densities. Therefore, dental imaging is an important step before placing a dental implant. Recently, Cone beam computed tomography (CBCT) is widely used in dental medicine and also recommended by AAOMR for preoperative implant placement. The aims of this study is to analyzed the alveolar bone density of the dental implant sites for dental implant planning using CBCT. **Methods:** ninety-three CBCT data were retrieved from the database of the department of oral and maxillofacial radiology at Dental Hospital Faculty of Dentistry Universitas Trisakti and examined. The recipient sites for dental implant placement were determined based on CBCT data using implant planning software (i-Dixel). The alveolar bones value is recorded in grayscale value (GV). **Results:** a great variety of alveolar bone density was observed ranging from 134-891 GV. One-way ANOVA was conducted, and statistically significant was only shown at the anterior mandibular region and molar mandibular region. However, no statistical differences were observed when comparing the male and female groups using the T-test. **Conclusion:** Critical evaluation through dental CBCT can be made before dental implant placement in the alveolar bones. the lowest alveolar bone density was observed in maxillary molar dental implant sites with mean 322 GV and highest alveolar bone density was shown in mandibular anterior dental implant sites with mean 558.8 GV. Determining the bone density at the dental implant site before dental implant placement is crucial. Primary stability and secondary stability is determined by bone density. Therefore, higher value of alveolar bone density is needed for dental implant placement.

**Keywords:** alveolar bone density assessment; dental implant; cone-beam computed tomography.

p-ISSN: 1979-0201; e-ISSN: 2549-6212; Available from: <http://jurnal.unpad.ac.id/pjd/article/view/42383>

DOI: [10.24198/pjd.vol34no3.42383](https://doi.org/10.24198/pjd.vol34no3.42383)

Submission: Oct 12, 2022; Accepted: Nov 30, 2022; Published online: Nov 30, 2022

\*Corresponding author: Wiwiek Poedjiastoeti, Department of Oral and Maxillofacial Surgery, Faculty of Dental Medicine, Universitas Trisakti, Jl. Kyai Tapa No. 260, Jakarta, 11440, e-mail: [wiwiek@trisakti.ac.id](mailto:wiwiek@trisakti.ac.id), Phone: 021-5655786



## INTRODUCTION

Tooth loss has been a problem for humans over the centuries. Since the 19<sup>th</sup> century, Dental implant has been used to replace missing teeth.<sup>1</sup> To date, the dental implant has been widely used because of its advantages, such as more protection for the surrounding teeth and the alveolar bone. On the other hand, rehabilitation treatment using dental implants has been reported to increase mastication function and quality of life.<sup>2</sup> Therefore, a dental implant can be an essential treatment option for replacing missing teeth.<sup>3</sup>

Dental implants have a wide variety of success rates in dentistry. The quantity and quality of bone availability are the primary determinants in predicting the success rate of the dental implant. However, different sites in the jaw have different success rates.<sup>4</sup> The functional quality of alveolar bones is determined by bone density. Bone density has a critical effect on surgical technique, healing time, and time of loading during prosthodontic reconstructions.<sup>5,6</sup> The golden site for dental implant placement is in the site with high bone density. Otherwise, the success rate decreases when the bone has a lower density or is porous.<sup>4</sup> Assessing the bone tissue has two objectives: first, is the bone tissue sufficient for dental implant placement (as a diagnostic tool), and second, to predict the success rate of the dental implant (prognostic tool).<sup>7</sup>

Poor bone density is considered a risk factor for biological complications of the implant, associated with lack of primary stability and impaired healing or osseointegration, which can lead to early implant loss.<sup>7</sup> There are two strategies to evaluate bone quality and quantity. The direct estimation strategy incorporates ex vivo examinations (i.e., dry skulls or corpses) or sample/biopsy recovered for examination from creatures or human subjects and in vivo examinations done on living subjects.

The indirect estimation strategy depends on radiographic imaging, like registered tomography (CT) or cone-beam computed tomography (CBCT). These estimation methods give a three-dimensional depiction of bony structures. They are viewed as a precise diagnostic instrument that enables the assessment of the morphology, bone quality, and volume of the leftover alveolar ridge.<sup>8</sup>

The American Academy of Oral and Maxillofacial Radiology (AAOMR) currently advocates using sectional imaging for all dental implant treatment.<sup>9</sup> CBCT provided volumetric data of the jaw with relatively low radiation doses and costs with high resolution.<sup>6,10</sup> Overall dental implant placement (implant size, angle, and type) can directly influence the alveolar bone quantity. On the other hand, the stability of dental implants is associated with bone quality. Therefore, a comprehensive assessment of both factors is essential.<sup>11</sup> Ivanova et al.<sup>12</sup> Concluded primary stability and secondary stability is determined by bone density.<sup>12</sup> Baltayan et al.<sup>7</sup> found a strong correlation between alveolar bone density and primary stability.<sup>13</sup>

Poor bone quality associated with lack of primary stability.<sup>7</sup> Therefore, to better understand the alveolar bone density of the edentulous area for dental implant planning using CBCT, we studied the edentulous area of the implant site on 93 patients who underwent preoperative evaluation using CBCT in our department in January 2018 - December 2021. The aims of this study is to analyzed the alveolar bone density of the dental implant sites for dental implant planning using CBCT.

## METHODS

In a cross-sectional study, ninety-three CBCT data of patients randomly selected from the department of oral and maxillofacial radiology database at Dental Hospital Faculty of Dentistry Universitas Trisakti (RSGMP-FKG Usakti) between January 2018 - December 2021 was evaluated. The CBCT data must meet the following inclusion criteria: CBCT data in patients aged a minimum of 20 years old and the radiograph taken between January 2018-December 2021, edentulous areas with the potential for dental implant placement in the jaw and the exclusion criteria is CBCT radiographs with poor quality.

Measurements Morita 3D Cone-Beam Computed Tomography (3D Accuitomo 170, J. Morita Co., Japan) was used to obtain patient 3D scan data at 90 KvP, 5 mA, and 80 µm voxel size exposure setting. Cross sectional view with 1.00 mm slice thickness was obtained. All CBCT data were evaluated by two calibrated oral and maxillofacial radiologists.




Figure 1. Bone density measurement. (A1) fixture simulation lower molar region (A2) bone density assessment lower molar region (A3) Sagittal view showing implant fixture simulation at lower molar region (A4) Coronal view lower molar region (B1) fixture simulation upper incisive region (B2) bone density assessment upper incisive region (B3) Sagittal view showing implant fixture simulation at upper incisive region (B4) Coronal view of upper incisive region.

Inter-observer agreement was calculated to verify the reliability of measurements. Bone density Head reorientation on CBCT images was undergone prior to bone density measurement. Bone density was measured using the software AIS-3D App by choosing an implant planning

feature build on the software. The next step was to determine the region of interest or tooth area to be measured (incisive, premolar, molar in both jaws) followed by implant insertion in the selected region, and the bone density mean value inside the implant volume will be calculated (Figure 1).

Table 1. Interpretation of Cohen's Kappa ( $\kappa$ )

Cohen's Kappa ( $\kappa$ )	Strength of Agreement
<0.00	Poor
0.01-0.20	Slight
0.21-0.40	Fair
0.41-0.60	Moderate
0.61-0.80	Substantial
0.81-1.00	Almost perfect

### Data analysis

We recorded the alveolar bone density data in Microsoft Excel then all statistical analyses conducted using SPSS Statistics for Windows V22 (IBM, USA). First, inter-observer reliability was observed using Cohen's Kappa ( $\kappa$ ) statistical analysis. The interpretation of Cohen's Kappa ( $\kappa$ ) can be seen in table 1.<sup>14</sup> Independent sample T-test was used to compare the mean difference between the male and female groups.

One-way ANOVA was used to compare the mean difference among age groups. Tukey post hoc analysis was used to compare all possible age group pairing when One-way ANOVA showed a significant difference. The study protocol was approved by the Komisi Etik Penelitian Kesehatan Fakultas Kedokteran Gigi Universitas Trisakti

(Ethical clearance committee, No 024/S3/KEPK/ FKG/7/2022)

### RESULTS

Two observers evaluated the collected CBCT image data of 93 subjects. First, we divided the alveolar bones into maxillary and mandibular and then differentiated them into three regions: anterior, premolar, and molar.

The kappa coefficients of inter-observer A and B were 0.92, 0.91, and 0.87, respectively, for the maxillary anterior region, maxillary premolar region, and maxillary molar region, indicating almost perfect agreement. However, the kappa coefficients for the mandibular anterior region bone density



Table 2. Inter-observer kappa coefficients

The Alveolar bones regions	Observer A to B
Maxillary Anterior Region	0.92
Maxillary Premolar Region	0.91
Maxillary Molar Region	0.87
Mandibular Anterior Region	0.70
Mandibular Premolar Region	0.95
Mandibular Molar Region	0.90

The result showed that mean maxillary anterior region bone density was  $362.4 \pm 107.35$  GV, maxillary premolar region bone density was  $326.6 \pm 77.12$  GV, maxillary molar region

bone density was  $322.0 \pm 103.21$  GV, mandibular anterior, premolar, and molar region density were  $557.8 \pm 160.29$  GV,  $402.8 \pm 111.06$  GV, and  $408.6 \pm 95.87$  GV respectively (Table 3).

Table 3. The alveolar bones density

The Alveolar bones regions	Min	Max	Mean	Std. Deviation
Maxillary Anterior Region	195	587	362.4	107.35
Maxillary Premolar Region	172	582	326.6	77.12
Maxillary Molar Region	134	745	322.0	103.21
Mandibular Anterior Region	319	891	557.8	160.29
Mandibular Premolar Region	257	848	402.8	111.06
Mandibular Molar Region	173	681	408.6	95.87

Note: all result value describes in grayscale value (GV)

The distribution of male and female patients was 42 and 51 people. The mean maxillary anterior region bone density was 348.5 GV, 377.6 GV for males and females, and maxillary premolar region bone density was 313.8 GV, and 341.6 GV for males and females, maxillary molar region bone density was 296 GV and 342.4 GV. Therefore, the mandibular anterior region bone density was 518.5

and 656.2 for males and females, the mandibular premolar region bone density was 373.1 GV and 428.8 GV for males and females, and the mandibular molar region bone density was 407.6 GV and 409.5 GV for male and female. However, no statistically significant difference was observed when comparing the male and female group bone density (Table 4).

Table 4. Bone density analysis based on gender and age group.

The Alveolar bones Region	Mean $\pm$ SD (GV) based on gender		p	Mean $\pm$ SD (GV) based on Age Group			p
	Male	Female		21-45	46-60	>61	
Maxillary anterior region	348.5 $\pm$ 117.4	377.6 $\pm$ 98.4	0.52	314.5 $\pm$ 69.1	366.2 $\pm$ 122.4	440.6 $\pm$ 35.5	0.25
Maxillary premolar region	313.8 $\pm$ 79.2	341.6 $\pm$ 73.9	0.26	294.2 $\pm$ 73.4	329.6 $\pm$ 84.5	356 $\pm$ 53.6	0.21
Maxillary molar region	296 $\pm$ 62.8	324.4 $\pm$ 124.9	0.27	302.7 $\pm$ 31	343.1 $\pm$ 175.5	318.6 $\pm$ 55.7	0.76
Mandibular anterior region	515.5 $\pm$ 139.6	656.2 $\pm$ 186.4	0.15	324.5 $\pm$ 7.7	570.2 $\pm$ 150.5	649.7 $\pm$ 102.3	0.04*
Mandibular premolar region	373.1 $\pm$ 90.4	428.8 $\pm$ 123.3	0.17	389.7 $\pm$ 55.7	412.2 $\pm$ 154.9	399.5 $\pm$ 54.8	0.90
Mandibular molar region	407.6 $\pm$ 94.4	409.5 $\pm$ 98.9	0.94	451.8 $\pm$ 66	412.8 $\pm$ 105.1	350.3 $\pm$ 80	0.01*

Note : Grayscale Value (a number that represents the amount of attenuation of the X-ray beam by the material contained in each voxel or structural unit of the tomographic volume )

\*p<0.05 = statistically significant difference

The median age was 50.34, ranging from 21 to 82 years old. We divided the age group into three groups, ranging from 21-45 (young adult),

46-60 (middle adult), and >61 (elderly) according to WHO age group classification in 2015. The one-way ANOVA test was carried out.

Table 5. Multiple comparison using Tukey post hoc test

The Alveolar bones Region	Age Group	Age Group	P	95% Confidence Interval	
				Lower bound	Upper bound
3 Maxillary Anterior Region	21-45	46-60	0.581	-181.54	78.11
	21-45	>61	0.231	-314.31	61.98
	46-60	>61	0.518	-243.73	94.83
Maxillary Premolar Region	21-45	46-60	0.458	-197.33	36.43
	21-45	>61	0.194	-147.08	23.48
	46-60	>61	0.666	-100.85	48.15
Maxillary Molar Region	21-45	46-60	0.747	-178.83	98.01
	21-45	>61	0.951	-147.69	115.91
	46-60	>61	0.879	-102.34	151.39
Mandibular Anterior Region	21-45	46-60	0.088	-526.46	34.96
	21-45	>61	0.038*	-632.76	-17.73
	46-60	>61	0.599	-296.94	137.94
Mandibular Premolar Region	21-45	46-60	0.898	-148.47	103.54
	21-45	>61	0.984	-151.90	132.40
	46-60	>61	0.966	-113.29	138.72
Mandibular Molar Region	21-45	46-60	0.384	-31.61	109.51
	21-45	>61	0.012*	19.02	183.96
	46-60	>61	0.113	-11.38	136.46

\*p<0.05 statistically significant difference

No significant differences were found between the maxillary anterior region, maxillary premolar region, maxillary molar region, and mandibular premolar region ( $p=0.255$ ,  $p=0.216$ ,  $p=0.760$ , and  $p=0.903$ , respectively) compared to age groups (Table 4). However, the mandibular anterior and mandibular molar regions showed significant differences compared to age groups with  $p=0.45$  and  $p=0.16$ . The multiple comparisons using Tukey post hoc clarify the differences only seen in the young adult group compared to elderly groups in the mandibular anterior and molar region (Table 5).

## PEMBAHASAN

Bone density assessment is indispensable to achieving the best dental implant treatment planning and the primary stability of the dental implant.<sup>15</sup> Bone density is one predictor of successful dental implant treatment. Therefore, an accurate evaluation of the alveolar bone density is essential. CBCT is three-dimensional dental imaging widely used recently in dental medicine. One of the several advantages of CBCT dental imaging enables the assessment of bone density.<sup>9</sup> This study was conducted to assess the

alveolar bone density of dental implant sites using the dental CBCT. A higher value of inter-observer reliability was observed in our study at table 2. Nevertheless, the mandibular anterior region has shown substantial agreement. This study's decrease in inter-observer reliability is thought to be because the available data for the mandibular anterior region is limited.

The result is that a slight difference in the assessment significantly influences the reliability. In line with the previous study, the agreement value of our result ranges from 0.70-0.95, and the previous study was 0.61-0.92.<sup>16</sup> Similar result was seen in a study conducted by Alkhader et al.<sup>17</sup> they showed the Cohen's kappa value was range from 0.77-0.93. The reliability of inter-observer reliability can be improved by practice and training in diagnostic imaging.<sup>14</sup>

The alveolar bone density was greater in females compared to males (Table 4). However, no statistically significant differences between male and female alveolar bone density were found. No data available from the previous study can be compared to our study. This is due to our CBCT data being recorded in grayscale value. Therefore, we try to compare the result with available data recorded in Hounsfield Unit (HU)



using Computed Tomography (CT). This decision was made based on a study that provided a strong correlation between GV of CBCT and HU of CT.<sup>18</sup> In contrast to our study, Cassetta et al.<sup>19</sup> found a statistically significant difference in alveolar bone density between gender, and they concluded that the density of the male alveolar bone was greater than that of females.<sup>19</sup>

A study conducted by Kim et al. was in line with Cassetta result; they found a statistical difference in the alveolar bone density between gender. However, in the study by Kim et al.<sup>20</sup> they showed a greater alveolar bone density was seen in females compared to males. Several risk factors may affect the density of the alveolar bone, such as gender, periodontal disease, osteoporosis,<sup>21</sup> tooth loss, menopause, and age.<sup>22</sup>

Our study showed that a great variety of alveolar bone density was found in the maxilla or mandibular. Statistically significant differences between age and alveolar bone density were observed in young adults compared to the elderly. Based on our result, age seems to affect alveolar bone density (Table 4). Other authors found different results that report that age does not affect the alveolar bone density value.<sup>21</sup>

Previously three-dimensional alveolar bones were evaluated using CT, including the alveolar bones density assessment. Consider the patient who received x-ray exposure. CT is not ideal for some diagnostic purposes in dental medicine due to the high dose of radiation, poor resolution, longer scanning time, difficult interpretation, high-cost charge, and limited availability.<sup>23</sup> To overcome the limitation, Arai et al.<sup>24</sup> (1999) in Japan and Mozzo et al.<sup>25</sup> (1998) in Italy at introduced cone-beam computed tomography (CBCT) for oral and maxillofacial application. The CBCT fulfils the needs, such as reducing unnecessary exposure to the patient, high resolution, rapid scan time, dose reduction, and reduced image artefact.<sup>26</sup>

The alveolar bone density plays a vital role in the mechanical immobilization of the dental implant during healing, distribution, and transition of stresses from the prosthesis to the implant-bone interface after healing. The availability of alveolar bone density determines the dental implant design, surgical approach, healing time, and initial progressive loading during prosthetic construction.<sup>27</sup> With the reason and data provided

above, preoperative evaluation of the alveolar bone density prior to dental implant placement at the edentulous area planned to a received dental implant is mandatory.

The limitation of our study is we did not compare the alveolar bone density based on the left and right regions of the jaw and lack of data available from our database for the mandibular anterior region. Hopefully, in the future, a study could be conducted to overcome the limitation.

## CONCLUSION

The highest alveolar bone density value was found on the mandibular anterior region, followed by the mandibular molar, mandibular premolar, maxillary anterior, maxillary premolar and maxillary molar regions. These conditions were noted in both gender and age groups. In the age group above 61 years old, the highest density of alveolar bone differences was found in the mandibular anterior region; however, the lowest density was noted in the mandibular molar region. The grayscale value in CBCT can be used to determine the alveolar bone density of dental implant sites.

## REFERENCES

1. Alghamdi HS, Jansen JA. The development and future of dental implants. *Dent Mater J.* 2020; 39(2): 167-172. DOI: [10.4012/dmj.2019-140](https://doi.org/10.4012/dmj.2019-140).
2. Elani HW, Starr JR, Da Silva JD, Gallucci GO. Trends in dental implant use in the U.S., 1999-2016, and projections to 2026. *J Dent Res.* 2018; 97(13): 1424-30. DOI: [10.1177/0022034518792567](https://doi.org/10.1177/0022034518792567).
3. Sulijaya B, Kuswandani SO, Soeroso Y. Contemporary guided bone regeneration therapy for unaesthetic anterior peri-implantitis case. *Dent J.* 2016; 49(4): 181-4. DOI: [10.20473/j.djmk.v49.i4.p181-184](https://doi.org/10.20473/j.djmk.v49.i4.p181-184).
4. Attar BM, Alaei S, Badrian H, Davoudi A. Clinical and radiological evaluation of implants placed with osteotome sinus lift technique: 19-month follow-up. *Ann Maxillofac Surg.* 2016; 6(2): 190-4. DOI: [10.4103/ams.ams.7.16](https://doi.org/10.4103/ams.ams.7.16).
5. Arina YMDa, Ferdiansyah F, Rubianto M. The evaluation of mandibular bone density in chronic periodontitis models. *Dent J.* 2018;

- 51(4): 210-5. DOI: [10.20473/j.djmk.v51.i4.p210-215](https://doi.org/10.20473/j.djmk.v51.i4.p210-215)
6. Eskandarloo A, Arabi R, Bidgoli M, Yousefi F, Poorolajal J. Association between marginal bone loss and bone quality at dental implant sites based on evidence from cone beam computed tomography and periapical radiographs. *Contemp Clin Dent*. 2019; 10(1): 36-41. DOI: [10.4103/ccd.ccd.185.18](https://doi.org/10.4103/ccd.ccd.185.18).
7. Chrcanovic BR, Albrektsson T, Wennerberg A. bone quality and quantity and dental implant failure: a systematic review and meta-analysis. *Int J Pros*. 2017; 30(3): 219-37. DOI: [10.11607/ijp.5142](https://doi.org/10.11607/ijp.5142).
8. Fokas G, Vaughn VM, Scarfe WC, Bornstein MM. Accuracy of linear measurements on CBCT images related to presurgical implant treatment planning: a systematic review. *Clin Oral Implants Res*. 2018; 29(16): 393-415. DOI: [10.1111/clr.13142](https://doi.org/10.1111/clr.13142).
9. Weiss R2nd, Read-Fuller A. Cone beam computed tomography in oral and maxillofacial surgery: an evidence-based review. *Dent J (Basel)*. 2019; 7(2): 52. DOI: [10.3390/dj7020052](https://doi.org/10.3390/dj7020052).
10. Jacobs R, Salmon B, Codari M, Hassan B, Bornstein MM. Cone beam computed tomography in implant dentistry: recommendations for clinical use. *BMC Oral Health*. 2018; 18(1): 88. DOI: [10.1186/s12903-018-0523-5](https://doi.org/10.1186/s12903-018-0523-5).
11. Wakimoto M, Matsumura T, Ueno T, Mizukawa N, Yanagi Y, Iida S. Bone quality and quantity of the anterior maxillary trabecular bone in dental implant sites. *Clinical oral implants research*. 2012; 23(11): 1314-9. DOI: [10.1111/j.1600-0501.2011.02347.x](https://doi.org/10.1111/j.1600-0501.2011.02347.x).
12. Igarashi C, Theramballi YG, Kobayashi K. Inter-observer reliability in cone-beam computed tomography assessment of the retromolar canal: A practical plan to improve diagnostic imaging. *Imaging Sci Dent*. 2022; 52(2): 181-6. DOI: [10.5624/isd.20210289](https://doi.org/10.5624/isd.20210289).
13. Sreerama R, Kolluru KC, Gottumukkala V, Innampudi CK, Konathala JR, Krishnaveni G. Assessment of the effect of bone density on implant stability: a clinical study. *J Pharm Bioallied Sci*. 2021; 13(1): S297-S300. DOI: [10.4103/jpbs.JPBS.794.20](https://doi.org/10.4103/jpbs.JPBS.794.20).
14. Kim JH, Abdala-Junior R, Munhoz L, Cortes ARG, Watanabe PCA, Costa C et al. Comparison between different cone-beam computed tomography devices in the detection of mechanically simulated peri-implant bone defects. *Imaging Sci Dent*. 2020; 50(2): 133-9. DOI: [10.5624/isd.2020.50.2.133](https://doi.org/10.5624/isd.2020.50.2.133)
15. Alkhader M, Hudieb M, Khader Y. Predictability of bone density at posterior mandibular implant sites using cone-beam computed tomography intensity values. *Eur J Dent*. 2017; 11(3): 311-6. DOI: [10.4103/ejd.ejd.14.17](https://doi.org/10.4103/ejd.ejd.14.17)
16. Razi T, Niknami M, Alavi Ghazani F. Relationship between Hounsfield Unit in CT scan and gray scale in CBCT. *J Dent Res Dent Clin Dent Prospects*. 2014; 8(2): 107-10. DOI: [10.5681/joddd.2014.019](https://doi.org/10.5681/joddd.2014.019).
17. Cassetta M, Sofan AA, Altieri F, Barbato E. Evaluation of alveolar cortical bone thickness and density for orthodontic mini-implant placement. *J Clin Exp Dent*. 2013; 5(5): 245-52. DOI: [10.4317/jced.51228](https://doi.org/10.4317/jced.51228)
18. Kim JH, Lim YJ, Kim B, Lee J. How do parameters of implant primary stability correspond with ct-evaluated bone quality in the posterior maxilla? A correlation analysis. *Materials (Basel)*. 2021; 14(2): 270. DOI: [10.3390/ma14020270](https://doi.org/10.3390/ma14020270)
19. Yu B, Wang CY. Osteoporosis and periodontal diseases - An update on their association and mechanistic links. *Periodontol 2000*. 2022; 89(1): 99-113. DOI: [10.1111/prd.12422](https://doi.org/10.1111/prd.12422).
20. Takaishi Y, Arita S, Honda M. Assessment of alveolar bone mineral density as a predictor of lumbar fracture probability. *Adv Ther*. 2013; 30(5): 487-502. DOI: [10.1007/s12325-013-0028-1](https://doi.org/10.1007/s12325-013-0028-1)
21. Saati S, Kaveh F, Yarmohammadi S. Comparison of cone beam computed tomography and multi slice computed tomography image quality of human dried mandible using 10 anatomical landmarks. *J Clin Diagn Res*. 2017; 11(2): ZC13-ZC16. DOI: [10.7860/JCDR/2017/20637.9253](https://doi.org/10.7860/JCDR/2017/20637.9253).
22. Venkatesh E, Elluru SV. Cone beam computed tomography: basics and applications in dentistry. *J Istanbul Univ Fac Dent*. 2017; 51(3 Suppl 1): S102-S121. DOI: [10.17096/jiufd.00289](https://doi.org/10.17096/jiufd.00289).
23. Sirisha Attili HS, Sitaram Prasad Kasina VHC Kumar, Srilatha Balusu, Sarat CB. To evaluate the bone mineral density in mandible of edentulous patients using computed tomography: an in vivo study. *J Int Oral Health*. 2015; 7(4): 22-6.

# Alveolar bones density assessment of dental implant sites using cone-beam computed tomography

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

14%

INTERNET SOURCES

12%

PUBLICATIONS

6%

STUDENT PAPERS

## PRIMARY SOURCES

- |   |   |    |
|---|---|----|
| 1 | Submitted to Fakultas Kedokteran Gigi Universitas Trisakti<br>Student Paper   | 3% |
| 2 | <a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a><br>Internet Source   | 3% |
| 3 | <a href="http://maxfac.no">maxfac.no</a><br>Internet Source   | 2% |
| 4 | <a href="http://journals.lww.com">journals.lww.com</a><br>Internet Source   | 1% |
| 5 | McPartland, J.M.. "Counterstrain and traditional osteopathic examination of the cervical spine compared", Journal of Bodywork & Movement Therapies, 199704<br>Publication         | 1% |
| 6 | Jacobs, Reinhilde, and Marc Quirynen. "Dental cone beam computed tomography: justification for use in planning oral implant placement", Periodontology 2000, 2014.<br>Publication | 1% |



7	<a href="https://pubmed.ncbi.nlm.nih.gov">pubmed.ncbi.nlm.nih.gov</a> Internet Source	1 %
8	<a href="https://worldwidescience.org">worldwidescience.org</a> Internet Source	1 %
9	<a href="https://jurnal.pdgi.or.id">jurnal.pdgi.or.id</a> Internet Source	1 %
10	Submitted to iGroup Student Paper	1 %
11	<a href="https://mafiadoc.com">mafiadoc.com</a> Internet Source	1 %
12	Leonard Morar, Grigore Băciuț, Mihaela Băciuț, Simion Bran, Horațiu Colosi, Avram Manea, Oana Almășan, Cristian Dinu. "Analysis of CBCT Bone Density Using the Hounsfield Scale", Prosthesis, 2022 Publication	1 %
13	<a href="https://etd.uwc.ac.za">etd.uwc.ac.za</a> Internet Source	1 %
14	<a href="https://inabj.org">inabj.org</a> Internet Source	1 %
15	<a href="https://www.frontiersin.org">www.frontiersin.org</a> Internet Source	1 %



Exclude quotes On

Exclude matches

< 15 words

Exclude bibliography On

# Alveolar bones density assessment of dental implant sites using cone-beam computed tomography

## GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/0

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7