

Instantly Check Your Writing

Check your grammar, spelling, and punctuation instantly with Grammarly

Grammarly

[Learn](#)

Journal of International Dental and Medical Research

COUNTRY

Turkey



Universities and research institutions in Turkey



Media Ranking in Turkey

SUBJECT AREA AND CATEGORY

Dentistry
Dentistry
(miscellaneous)

PUBLISHER

Ektodermal Displazi
Grubu

H-INDEX

15

PUBLICATION TYPE

Journals

ISSN

1309100X

COVERAGE

2009-2022

SCOPE






Information not localized




Join the conversation about this journal

FIND SIMILAR JOURNALS ?

<p>1</p> <p>Journal of International Oral Health</p> <p>IND</p> <p>40%</p> <p>similarity</p>	<p>2</p> <p>Pesquisa Brasileira em Odontopediatria e Clínica</p> <p>BRA</p> <p>38%</p> <p>similarity</p>	<p>3</p> <p>Journal of International Society of Preventive and</p> <p>IND</p> <p>37%</p> <p>similarity</p>	<p>4</p> <p>Indian Journal of Dental Research</p> <p>IND</p> <p>36%</p> <p>similarity</p>
--	--	--	---





← Show this widget in your own website

Just copy the code below and paste within your html code:

``



Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Metrics based on Scopus® data as of April 2023

R **Rrahman FERIZI** 2 months ago

Could You please clarify with the " Beall's List" (<https://beallslist.net/standalone-journals/#update>) to "remove" this Journal (JIDMR) from this "un-fictive" list, to "protect" Your Database aspects, because we can't get any response from the Journal (JIDMR) management even though we have asked them this?!

Best regards!

reply



Melanie Ortiz 2 months ago


SCImago Team

Dear Rrahman,
Thank you for contacting us. Unfortunately, SCImago cannot help you with your request. SJR is committed to help decision-making through scientometric indicators.
Best Regards, SCImago Team

M **Michael Kim** 4 months ago

Dear SCImago Team,
I think Scopus should review the criteria of this magazine to see if this is a predator journal or not?
For some of the following reasons:
1. Some articles do not have a DOI index
2. The grammar in articles is very bad
3. The magazine's home page is not clear, the domain name may be traded to another owner
4. The quality of reviews is not really good, leading to the time from submission to publication of

Journal of
International
Dental and Medical
Research



Editorial Board of JIDMR

Prof. Dr. Izzet YAVUZ
Editor-in-Chief and General Director

Advisory Board

Prof. Dr. Refik ULKU **Editor for Medicine**
Prof. Dr. Zulkuf AKDAG **Editor for Biomedical Research**
Prof. Dr. Ozkan ADIGUZEL **Associate Editor**

Gajanan Kiran KULKARNI (CANADA)
Betul KARGUL (TURKEY)
Diah Ayu MAHARANI (INDONESIA)
Francisco Cammarata-Scalisi (Venezuela)
Myroslav Goncharuk-Khomyn (UKRAINE)

Ferranti WONG (UK)
Zeki AKKUS (TURKEY)
Michele CALLEA (ROME, ITALY)
Zelal ULKU (TURKEY)

Moschos A. PAPAD
Lindawati S KUSDH
Yasemin YAVUZ (TL)
Yuliya MAKEDONO
Nik Noriah Nik HUS

Editorial Board

Abdel Fattah BADAWI (EGYPT)

Abdurrahman ONEN (TURKEY)

Ahmet YALINKAYA (TURKEY)

Ahmet DAG (TURKEY)

Ali Al-ZAAG (IRAQ)

Ali BUMIN (TURKEY)

Ali FADEL (EGYPT)

Guvenc BASARAN (TURKEY)

Güven ERBİL (TURKEY)

Halimah AWANG (MALAYSIA)

Halit AKBAS (TURKEY)

Heloisa Fonseca MARAO (BRAZIL)

Hilal TURKER (TURKEY)

Huseyin ASLAN (TURKEY)

Nezahat AKPOLAT (TURKEY)

Nihal HAMAMCI (TURKEY)

Nik Noriah Nik HUSSEIN (MA)

Nicola Pranno (ROME)

Nurten AKDENİZ (TURKEY)

Nurten ERDAL (TURKEY)

Orhan TACAR (TURKEY)

Ali GUR (TURKEY)	Igor BELYAEV (SWEDEN)	Ozant ONCAG (TURKEY)
Ali Riza ALPOZ (TURKEY)	Ilhan INCI (ZURICH)	Ozgur UZUN (TURKEY)
Ali Riza Tunçdemir (TURKEY)	Ilker ETIKAN (TURKEY)	Ozkan ADIGUZEL (TURKEY)
Allah Bakhsh HAAFIZ (USA)	Isil TEKMEK (TURKEY)	Rafat Ali SIDDIQUI (PAKISTA
Alpaslan TUZCU (TURKEY)	Isin ULUKAPI (TURKEY)	Refik ULKU (TURKEY)
Alpen ORTUG (TURKEY)	Jalen DEVECIOGLU KAMA (TURKEY)	Sabiha Zelal ULKU (TURKEY
Armelia Sari WIDYARMAN (INDONESIA)	Kemal CIGDEM (TURKEY)	Sabri BATUN (TURKEY)
Ashish AGGARWAL (INDIA)	Kemal NAS (TURKEY)	Sadullah KAYA (TURKEY)
Ayşe GUNAY (TURKEY)	Kewal KRISHAN (INDIA)	Saul Martins PAIVA (BRAZIL)
Aziz YASAN (TURKEY)	King Nigel MARTYN(HONG KONG, CHINA)	Sedat AKDENİZ (TURKEY)
Awiruth KLAISIRI (THAILAND)	Kursat ER (TURKEY)	Seher GUNDUZ ARSLAN (TL
Balasubramanian MADHAN (INDIA)	Levent ERDINC (TURKEY)	Selahattin ATMACA (TURKEY
Benik HARUTUNYAN (ARMENIA)	Luca TESTARELLI (ROME)	Selahattin TEKES (TURKEY)
Betul KARGUL (TURKEY)	Lucianne Cople MAIA (BRAZIL)	Serdar ERDINE (TURKEY)
Betul URREHMAN (UAE)	Luciane Rezende COSTA (BRAZIL)	Serdar ONAT (TURKEY)
Bugra OZEN (TURKEY)	Marri Sai ARCHANA (INDIA)	Sergio Adriane Bezerra DE M
Carlos Menezes AGUIAR (BRAZIL)	Manoj KUMAR (INDIA)	Serhan AKMAN (TURKEY)
Cemil SERT (TURKEY)	Marcelo Rodrigues AZENHA (BRAZIL)	Sertac PEKER (TURKEY)
Chiramana SANDEEP (INDIA)	Marcia Cancado FIGUEIREDO (BRAZIL)	Seyed Amir Danesh SANI (UŞ
Christine Bettina STAUDT (SWITZERLAND)	Marco MONTANARI (ITALY)	Seyit Burhaneddin ZINCIRCIC
Cihan AKGUL (TURKEY)	Margaret TZAPHLIDOU (GREECE)	Shailesh LELE (INDIA)
Claudia DELLAVIA (ITALY)	Maria Elisa Oliveira dos SANTOS (BRAZIL)	Sinerik N. AYRAPETYAN (AR
Diah Ayu MAHARANI (INDONESIA)	Medi GANIBEGOVIC (BOSNIA and HERZEGOVINA)	Smaragda KAVADIA (GREEK
Dinesh Rokaya (NEPAL)	Mehmet DOGRU (TURKEY)	Sossani SIDIROPOULOU (GI
Edoardo BAUNER (ROMA)	Mehmet Emin ERDAL (TURKEY)	Stefano Di CARLO (ROME)
Emmanuel Joao N. Leal da SILVA (BRAZIL)	Mehmet Sinan DOGAN (TURKEY)	Sunit Kr. JUREL (INDIA)
Emin Caner TUMEN (TURKEY)	Mehmet Zulkuf AKDAG (TURKEY)	Stephen D. SMITH (USA)
Emrullah BAHSI (TURKEY)	Meral ERDİNC (TURKEY)	Susumu TEREKAWA (JAPAN
Ertunc Dayı (TURKEY)	Michele CALLEA (ITALY)	Suha TURKASLAN (TURKEY
Fadel M. ALI (EGYPT)	Mohamed TREBAK (USA)	Suleyman DASDAG (TURKE'
Fahinur ERTUGRUL (TURKEY)	Mohammad Khursheed Alam (KSA)	Taskin GURBUZ (TURKEY)
Feral OZTURK (TURKEY)	Mohammad SUHRON (INDONESIA)	Ufuk ALUCLU (TURKEY)
Feridun BASAK (TURKEY)	Mohammed Mustahsen URREHMAN (UAE)	Ugur KEKLIKCI (TURKEY)
Ferranti WONG (UNITED KINGDOM)	Moschos A. PAPADOPOULOS (GREECE)	Xiong-Li YANG (CHINA)
Feyzi Çelik (TURKEY)		Vatan KAVAK (TURKEY)

Feyzullah Uçmak (TURKEY)	Mostaphazadeh AMROLLAH (IRAN)	Yasar YILDIRIM (TURKEY)
Figen SEYMEN (TURKEY)	M.S. Rami REDDY (INDIA)	Yasemin YAVUZ (TURKEY)
Filippo BATTELLI (ITALY)	Muhammad FAHIM (INDIA)	Yavuz SANISOGLU (TURKEY)
Filiz Acun KAYA (TURKEY)	Mukadder ATMACA (TURKEY)	Yu LEI (USA)
Flavio Domingues Das NEVES (BRAZIL)	Murat AKKUS (TURKEY)	Yuri LIMANSKI (UKRAINE)
Folakemi OREDUGBA (NIGERIA)	Murat SOKER (TURKEY)	Zafer C. CEHRELI (TURKEY)
Francesca De ANGELIS (ITALY)	Mustafa KELLE (TURKEY)	Zeki AKKUS (TURKEY)
Gaetano PAOLONE (ITALY)	Mustafa ZORTUK (TURKEY)	Zuhal KIRZIOGLU (TURKEY)
Gajanan Kiran KULKARNI (CANADA)	Muzeyyen YILDIRIM (TURKEY)	Zurab KOMETIANI (GEORGIA)
Gamze AREN (TURKEY)	Neval Berrin ARSERIM (TURKEY)	Zurab S. KHABADZE (RUSSIA)
Gauri LELE (INDIA)		
Gonul OLMEZ (TURKEY)		
Gulsen YILMAZ (TURKEY)		
Gulten UNLU (TURKEY)		



Journal of International Dental and Medical Research

Research



Contents of JIDMR 2022 – Vol.15 – No.2

29/06/2022 /

**Journal of International Dental and
Medical Research**

ISSN: 1309-100X

Journal of
International
Dental and Medical
Research



2022 - Vol. 15 – No. 2

<http://www.jidmr.com>

Download Cover page



[<http://www.jidmr.com/journal/wp-content/uploads/2022/06/Cover15-2.pdf>]

Current Issue of JIDMR

Table of Contents 2022 Vol.15–No.2



[http://www.jidmr.com/journal/wp-content/uploads/2022/06/0_2_15_22_Table_of_Cc

DENTISTRY

EXPERIMENTAL ARTICLE

1. Roughness and Wear on Enamel Treated with a Microabrasive Agent

Paola-Andrea Rojas-Martínez, Daniela Álvarez-Quiceno, Lenin-Rafael Torres-Osorio, Alberto-Carlos Cruz-González

Full Text

PDF

Pages 461-467

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/1-D22_1746_Alberto_Carlos_Cruz_Gonzalez_Colombia.pdf]

EXPERIMENTAL ARTICLE

2. Basic Fibroblast Growth Factor Expression after Gingival Mesenchymal Stem Cell's Metabolite Provision in Lipopolysaccharide induce inflammatory bone resorption in vivo

Dr. Delany, Alexander Delany, Alexander Delany, Alexander Delany, Alexander Delany

Dwi Kanmawati, Alexander Patera Nugraha, Alivianda Zahrina, Nastiti Faradilla Ramadhani, Wibi Riawan,

Igo Syaiful Ihsan, Diah Savitri Ernawati, Ida Bagus Narmada, Tania Saskianti, Fianza Rezkita, Andari Sarasati, Albertus Putera Nugraha, Tengku Natasha Eleena Binti Tengku Ahmad Noor

Full Text **PDF**

Pages 468-471

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/2-D22_1778_Alexander_Patera_Nugraha_Indonesia.pdf]

EXPERIMENTAL ARTICLE

3. Effect of Cementation Materials and Sonic Application on Push-Out Bond Strength of Fiber Posts to Root Canal Dentin

Pornsawan Sirianothaikul, Wanhip Plooksawasdi, Juthatip Aksornmuang

Full Text **PDF**

Pages 472-478

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/3-D22_1809_Juthatip_Aksornmuang_Thailand.pdf]

EXPERIMENTAL ARTICLE

4. Characterization of Bovine Sponge Amnion (BSA) by a Novel Process for Dental Treatment

Octarina, Elly Munadzirroh, Fathilah Abdul Razak, Basril Abbas, Helal Soekartono, Titien Hary Agustantina, Meircurius Dwi Condro Surboyo

Full Text **PDF**

Pages 479-484

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/4-10-4-Characterization-D22_1783_Dian_Agustin_Wahjuningrum_Indonesia.pdf]

EXPERIMENTAL ARTICLE

5. Antimicrobial Efficacy of Chlorhexidine and Hyaluronic Acid Mouthwashes on Streptococcus Viridans: An In-Vitro Study

Omar Mohamed Shehatta, Ahmed Luay Osman, Walid Shaaban Elsayed, Sessa Reddy, Jovita Dsouza, Hossam Abdelmagyd, Devapriya Finney Shadroch

Full Text **PDF**

Pages 485-489

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/5-D21_1653_Jovita_Dsouza_UAE.pdf]

EXPERIMENTAL ARTICLE

6. The Effectiveness of Tobacco Leaf Effervescent Tablets (Nicotiana Tabacum L.) 75% against Surface Roughness and Acrylic Color Change

Dewi Kristiana, Ady Soesetijo, A. Gunadi, Rahardyan Parnaadji, Amiyatun Naini, Surartono Dwiatmoko, Delvi Fitriani

Full Text **PDF**

Pages 490-497

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/6-D22_1759_Dewi_Kristiana_Indonesia.pdf]

EXPERIMENTAL ARTICLE

7. Antibacterial Activity of *Salvadora Persica* and Benzylisothiocyanate against *Prevotella Intermedia* and *Eikenella Corrodens* Incorporated into Periodontal Chips

Fouad H. AL-Bayaty, Muhammad Azly Abdul Razak, Saba F. Hussain, Tengku Z. Mulok, Khalid Almas, Steph Smith, and Mahmood A. Abdulla

Full Text 

Pages 498-504

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/7-D22_1800_Fouad_H_AL_Bayaty_Malaysia.pdf]

EXPERIMENTAL ARTICLE

8. Expression of HSP 70, Nf-Kb and TNF- α in Inflammation Response Post Dental Pulp Tissue Extirpation

Galih Sampoerno, Anuj Bhardwaj, Eddo Supriyanto, Salwa Aulia Rahmawati, Wanda Salsabilla

Full Text 

Pages 505-510

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/8-10-6-hsp70-D22_1783_Dian_Agustin_Wahjuningrum_Indonesia.pdf]

EXPERIMENTAL ARTICLE

9. Evaluation of Dentin Microcracks after Root Canal Instrumentation using Three Thermally Treated Rotary Nickel Titanium Files

Johnson J., Abd-Elaal H., Elsewify T., El-Sayed W., Eid B.

Full Text 

Pages 511-515

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/9-D22_1811_Tarek_Medhat_Elsewify_Egypt.pdf]

EXPERIMENTAL ARTICLE

10. Accuracy of Volume of Interest Determination with Cone-Beam Computed Tomography in Periodontitis due to Psychological Stress

Baby Prabowo, Ani Melani Maskoen, Arief Budiarto, Fourier Dzar Eljabbar Latief, Arni Diana Fitri, Arif Rachman

Full Text 

Pages 516-520

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/10-D22_1770_Arif_Rachman_Indonesia.pdf]

EXPERIMENTAL ARTICLE

11. Morphological Assessment of the Gum after Transplantation of a Free Mucosal Graft Subjected to Processing by Er;Cr;YSGG 2780nm Laser (Experimental Study on Mini Pigs)

T.V. Furtsev, V.A. Khorzhevsky, A.A. Dernovoy, M.M. Nikolaenko, B.F. Cherkashin, A.I. Volynkina

Full Text **PDF**

Pages 521-525

[<http://www.jidmr.com/journal/wp-content/uploads/2022/06/11->

D22_1749_Furtsev_Taras_Vladimirovich_Russia.pdf]

EXPERIMENTAL ARTICLE

12. Evaluation of Bubble Tea Drink on Metabolism Streptococcus Mutans Atcc 25175

Juni Handajani, Britney Amadea Christianty, Annisa Fadiya Rahmawati

Full Text **PDF**

Pages 526-533

[<http://www.jidmr.com/journal/wp-content/uploads/2022/06/12->

D22_1708_Juni_Handajani_Indonesia.pdf]

EXPERIMENTAL ARTICLE

13. Surface roughness of monolithic ceramic and ceramic-like CAD/CAM materials

Sadullah Uctasli, Banu Yıldırım, Ünsun Çetin

Full Text **PDF**

Pages 534-537

[<http://www.jidmr.com/journal/wp-content/uploads/2022/06/13->

D21_1441_Sadullah_Uctasli_Turkey.pdf]

EXPERIMENTAL ARTICLE

14. Mixed Polymethylmethacrylate and Hydroxyapatite as a Candidate of Synthetic Graft Materials for Cleft Palate

Tania Saskianti, Aisyah Novianti, Diina Sahar, Alexander Patera Nugraha, Tansza Setiana Putri, Masami Kanawa, Mega Moeharyono Puteri, Ardianti Maartrina Dewi, Diah Savitri Ernawati

Full Text **PDF**

Pages 538-543

[<http://www.jidmr.com/journal/wp-content/uploads/2022/06/14->

D22_1742_Tania_Saskianti_Indonesia.pdf]

EXPERIMENTAL ARTICLE

15. Cytotoxicity and Cell Migration effect of Crude Spilanthes Acmella Ethanolic and Water Extract

Fatonah Charu, Sissada Tannukit, Nattapon Rotpenpian, Suwanna Jitpukdeebodindra

Full Text **PDF**

Pages 544-551

[<http://www.jidmr.com/journal/wp-content/uploads/2022/06/15->

D22_1808_Suwanna_Jitpukdeebodindra_Thailand-2.pdf]

EXPERIMENTAL ARTICLE

16. Molecular Docking Analysis of Quercetin and Diclofenac as Cox-2 Potential Inhibitors

Christian Khoswanto, Siswandono

Full Text **PDF**

Pages 552-555

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/16-D22_1773_Christian_Khoswanto_Indonesia.pdf]

EXPERIMENTAL ARTICLE

17. Canal Transportation and Centering Ability of Thermally-Treated NiTi Files

Alarfaj B., Elsewify T., El-Sayed W., Eid B.

Full Text **PDF**

Pages 556-560

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/17-D22_1812_Tarek_Medhat_Elsewify_Egypt-2.pdf]

EXPERIMENTAL ARTICLE

18. Expression of Bmp-2 in the Alveolar Rats' Extraction Socket after High Sucrose Diet

Christian Khoswanto

Full Text **PDF**

Pages 561-563

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/18-D22_1805_Christian_Khoswanto_Indonesia.pdf]

EXPERIMENTAL ARTICLE

19. Role of Ziziphus mauritiana Lam. on Oral Candidiasis and its Relation to the Antibody Response, Blood Electrolyte, and Liver Profile

Utmi Arma, Widyawati Widyawati, Abu Bakar, Basri A. Gani

Full Text **PDF**

Pages 564-573

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/19-D22_1797_Abu_Bakar_Indonesia.pdf]

EXPERIMENTAL ARTICLE

20. Effect of Strengthening Methods on Flexural Properties of Polymethylmethacrylate

Titiphong Phasookjai, Pisaisit Chaijareenont, Pimduen Rungsiyakull

Full Text **PDF**

Pages 574-580

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/20-D22_1810_Pimduen_Rungsiyakull_Thailand.pdf]

EXPERIMENTAL ARTICLE

21. Reconstruction and Optimized Expression of a Synthetic Secretory Leukocyte Protease Inhibitor (SLPI) Gene in Escherichia coli BL21

Elly Munadzirroh, Evi Umayah Ulfa, One Asmarawati, Ni Nyoman Tri Puspangsih, Mudyawati Kamaruddin, Muhammad Riza Hafidz Bahtiar, Arlita Rifyanti Yasminda

Full Text **PDF**

Pages 581-586

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/21-10-8-Reconstruction-D22_1783_Dian_Agustin_Wahjuningrum_Indonesia.pdf]

EXPERIMENTAL ARTICLE

22. Hyperbaric Oxygen Therapy Reduces Inflammation in Nickel Hypersensitivity

Meinar Nur Ashrin, Rima Parwati Sari, Dwi Andriani

Full Text **PDF**

Pages 587-593

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/22-D21_1672_Meinar_Nur_Ashrin_Indonesia.pdf]

EXPERIMENTAL ARTICLE

23. The Oral Administration of Capsaicin Regulate Prostaglandin E2 and Endorphin Serum Level after 24 Hours of Orthodontic Tooth Movement in Wistar Rats (*Rattus Novergicus*)

Ervina Restiwulan Winoto, Ida Bagus Narmada, I Ketut Sudiana, Hari Basuki, Retno Pudji Rahayu, Siswandono, Alexander Patera Nugraha

Full Text **PDF**

Pages 594-599

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/23-D21_1682_Ida_Bagus_Narmada_Indonesia.pdf]

EXPERIMENTAL ARTICLE

24. Fractionated Ethanol Extract of Red Ginger (*Zingiber officinale* var. *rubrum*) as Anti-Inflammatory Drug: An In-Silico Study

Fitrul A.E. Farhana, Prawati Nuraini, Seno Pradopo

Full Text **PDF**

Pages 600-605

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/24-D21_1704_Prawati_Nuraini_Indonesia.pdf]

EXPERIMENTAL ARTICLE

25. Dysplastic Model of Oral Squamous Cell Carcinoma in Male Wistar Rat: Chemically Induction with Dimethyl Benz(A) anthrance (DMBA)

Ahyar Riza, Denny Satria, Alfath Nugroho, Jeremias Lumban Toruan

Full Text **PDF**

Pages 606-610

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/25-D21_1683_Ahyar_Riza_Indonesia.pdf]

EXPERIMENTAL ARTICLE

26. The Effectiveness of Moringa Leaf Extract (*Moringa Oleifera*) Against *Porphyromonas gingivalis* Bacteria in Periodontitis Cases Through IL-1 Cytokine Analysis

Irene Edith Rieuwpassa, Sri Ramadany, Sumintarti, Israyani, Harun Achmad, Rini Sitanaya, Hans Lesmana, Arni Irawaty Djais, Arrang Sesioria, Harmawaty, Nur

Full Text **PDF**

Pages 611-617

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/26-D22_1771_Harun_Achmad_Indonesia.pdf]

EXPERIMENTAL ARTICLE

27. Synthesis of Cellulose Fiber from Coconut Coir as Potential Application of Dental Flowable Composite Filler

Twii Agnita Cevanti, Mahardika F. Rois, Nur Shiyama Purnama Sari, Steella Ilham Isnaini, Sella Ramadhani Alya Sasono, Ghaly Muhammad Bahreisy Firdaus, Heru Setyawan, Adioro Soetojo,
Ira Widjiastuti

Full Text **PDF**

Pages 618-622

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/27-10-1-Sintetis-D22_1783_Dian_Agustin_Wahjuningrum_Indonesia.pdf]

CLINICAL ARTICLE

28. The use of Lysozyme Toothpaste to Prevent Early Childhood Caries (Ecc) in 2 Years Old Children

Essie Octiara, Heriandi Sutadi, Yahwardiah Siregar, Ameta Primasari

Full Text **PDF**

Pages 623-629

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/28-D21_1700_Essie_Octiara_Indonesia.pdf]

CLINICAL ARTICLE

29. Efficacy of Osseodensification versus Expander Technique for Alveolar Ridge Expansion: A 3-Years Randomised Controlled Trial

Fakhreldin H. Abdel-Rahman, Eman A. Yousef, Mohamed A. Tawfik, Ola M. Maria

Full Text **PDF**

Pages 630-640

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/29-D22_1757_Ola_M_Maria_Canada.pdf]

CLINICAL ARTICLE

30. Development and Implementation of Digital Technologies in Dental Practice

Yuliya. A. Makedonova, Ludmila M. Gavrikova, Maria V.Kabytova, Denis Yu.Dyachenko, Oksana N. Kurkina, Svetlana V. Dyachenko, Artem Yu. Tkalin, Lev V. Zhuravlev

Full Text **PDF**

Pages 641-648

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/30-D22_1803_Yuliya_A_Makedonova_Russia.pdf]

CLINICAL ARTICLE

31. Knowledge, Attitude and Perception of Private Dental Practitioners Towards Medical Emergencies in Klang Valley, Malaysia

Nor Azura Binti Ahmad Tarmidzi, Nalisha Binti Mohamed Ramli, Norashikin Amran, Husna Najihah Binti Norazmi, Nur Saniah Binti Mohd Arifin

Full Text 

Pages 649-655

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/31-D22_1802_Nor_Azura_Ahmad_Tarmidzi_Malaysia.pdf]

CLINICAL ARTICLE

32. Reduction of Excessive Overjet in Pediatric Malocclusion Using Myofunctional Therapy Accompanied by Electromyography Activity Evaluation in Orofacial Muscles

Harun Achmad, Intan Sari Areni, Sri Ramadany, Nur Hildah Inayah, Rosdiana Agustin, Reza Ardiansya

Full Text 

Pages 656-668

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/32-D22_1766_Harun_Achmad_Indonesia.pdf]

CLINICAL ARTICLE

33. Impact of Oral Health and Oral Health Behavior on Happiness Among Young Adults in Mangalore City – An Explorative Study

Avinash B R, Ramya Shenoy, Mithun Pai B H, Rajesh G

Full Text 

Pages 669-674

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/33-D22_1767_Mithun_Pai_BH_India.pdf]

CLINICAL ARTICLE

34. Root Canal Morphology of Permanent Lower Premolars in Qatari Population: A Cone-Beam Computed Tomography Study

Amal Diab, Fatima Al-Sheeb, Nadya Marouf, Mashael Lari, Hanan Diab

Full Text 

Pages 675-679

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/34-D22_1813_Amal_Issam_Diab_Qatar.pdf]

CLINICAL ARTICLE

35. Mucocele Excision Using Electrocautery in Pediatric Patient: A Case Report

Felita Putri Sutjipto, Ardianti Maartrina Dewi, Siti Nur Lestari, Firli Cahaya Khairani, Udijanto Tedjosongko, Soegeng Wahlujo, Ilvana Ardiwirastuti, Alit Rahma Estu, Achmad Nadian Permana, Puspita Ayuningtyas

Full Text 

Pages 680-684

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/35-D22_1823_Ardianti_Maartrina_Dewi_Indonesia.pdf]

CLINICAL ARTICLE

36. Modified Periodontal Risk Parameters (MPRP) for Periodontal Management by Risk Assessment (PEMBRA): A Pilot Study

Mahyunah Masud, Nurkhairunnisa A. Kadir, Nur Asma Mahirah Mahlil, Izyan Hazwani Baharuddin, Nila Kasuma

Full Text 

Pages 685-690

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/36-D22_1764_Mahyunah_Masud_Malaysia.pdf]

CLINICAL ARTICLE

37. Comparison of Digital and Manual Determination of Maxillary Sinus Volume: A CBCT Study

Hamad Abdullah Alshiddi, Abdulrahman Khalid Alkhaldi, Khalid Almas, Mohammad Abdulaziz Alsaati, Saad Saeed Alzahrani, Mansour Fahad Aljubair, Intesar Siddiqui, Steph Smith

Full Text 

Pages 691-698

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/37-D22_1814_Khalid_Almas_K.S.A..pdf]

CLINICAL ARTICLE

38. The Application of Teledentistry : an Alternative Dental Service in Pandemic Era

Nila Kasuma, Purwita Nurwidyastuti, Citra Lestari, Harfindo Nismal, Murniwati, Deli Mona, Gian Ernesto

Full Text 

Pages 699-706

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/38-D22_1739_Fauzan_Yusro_Indonesia.pdf]

CLINICAL ARTICLE

39. Peeragogy Teaching Method in Oral Pathology Course

Nur Suraiya Izzaty Sabari, Nur Najihah Roslan, Nuramirah Azizan, Farinawati Yazid, Norliwati Ibrahim

Full Text 

Pages 707-711

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/39-D21_1652_Norliwati_Ibrahim_Malaysia.pdf]

CLINICAL ARTICLE

40. The Influence of Social Determinant and Chronic Disease on Self-Reported Tooth Loss in Indonesian Middle-Aged and Elderly

Sri Susilawati, Fidya Meditia Putri, Kurnia Wahyudi, Tince Arniati Jovina, Azlina Nuur Sanjaya, Anne Agustina Suwargiani

Full Text 

Pages 712-719

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/40-D22_1739_Fauzan_Yusro_Indonesia.pdf]

CLINICAL ARTICLE

41. Knowledge and Perception about Dental Implants among Undergraduate Dental Students and Interns in UAE

Azzah Eissa AlHemrani, Jayadevan Sreedharan, Salem Hassan Abu Fanas, Jovita Dsouza, Sesha Reddy, Hossam Abdelmagyd

Full Text 

Pages 720-727

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/41-D21_1679_Jovita_Dsouza_UAE.pdf]

CLINICAL ARTICLE

42. Head and neck cancer awareness exploratory survey in undergraduate students

Masita Mandasari, Ambar Kusuma Astuti, Ratna Kumala Indrastiti, Nathaniel Simon Treister, Jesslyn Karsono, Sonia Adhelia Sulisfianty

Full Text 

Pages 728-734

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/42-D21_1690_Masita_Mandasari_Indonesia.pdf]

CLINICAL ARTICLE

43. Comparative Evaluation of Powered and Manual Toothbrushes in Reducing Plaque and Gingivitis in Institutionalized Orphan Children

Madiraju Guna Shekhar, Sajith Abraham, Sheethal Joy, Nazargi Mahabob

Full Text 

Pages 735-740

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/43-D22_1753_Guna_Shekhar_M_SA.pdf]

CLINICAL ARTICLE

44. Knowledge and Perceptions Regarding Tobacco Harm Reduction among Dentists in West Java, Indonesia

Aliyya Puteri Ramadhani, Budhi Cahya Prasetyo, Amaliya Amaliya, Konstantinos E. Farsalinos

Full Text 

Pages 741-750

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/44-D22_1747_Amaliya_Amaliya_Indonesia.pdf]

CLINICAL ARTICLE

45. Effectiveness of Asynchronous VS Synchronous Online Histology Learning in Covid-19 Era Among Dental Students

Nor Asmaq Mohd Said, Siti Nur Sabrina Ahmad, Farinawati Yazid, Nurul Shaqinah Nasruddin, Norliwati Ibrahim, Siak Kai Hun, Azizah Ahmad Fauzi

Full Text **PDF**

Pages 751-756

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/45-D22_1761_Azizah_binti_Ahmad_Fauzi_Malaysia.pdf]

CLINICAL ARTICLE

46. The Oral Biology Parameter of the Diabetes Mellitus Type-2 Patients Relate to the Oral Candida Species Development

Sri Rezeki, Rachmi Fanani Hakim, Sunnati Sunnati, Salvinia Salvinia, Basri A. Gani

Full Text **PDF**

Pages 757-765

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/46-D21_1701_Sri_Rezeki_Indonesia.pdf]

CLINICAL ARTICLE

47. Characteristics and Predicting Factors of Repeat Dental Treatment Under General Anesthesia in Patients with Special Health Care Needs: A Retrospective Study

Pornpailin Kasemkhun, Apiwan Smutkeeree, Varangkanar Jirattanasopha

Full Text **PDF**

Pages 766-770

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/47-D22_1782_Pornpailin_Kasemkhun_Thailand.pdf]

CLINICAL ARTICLE

48. A Retrospective Study on the Labial Maxillary and Mandibular Alveolar Bone after Anterior Retraction

Hilda Fitria Lubis, Aditya Rachmawati, Stephani Tanius

Full Text **PDF**

Pages 771-775

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/48-D22_1825_Hilda_Fitria_Lubis_Indonesia.pdf]

CLINICAL ARTICLE

49. Needs Assessment of Oral Health Care Supplies for Dependent Older Adults in Phitsanulok Province, Thailand

Sarinlavan Somboonchan, Piyanart Chatiketu, Ramya Shenoy, Pimduen Rungsiyakull

Full Text **PDF**

Pages 776-782

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/49-D22_1786_Pimduen_Rungsiyakull_Thailand.pdf]

CLINICAL ARTICLE

50. Dentist Perceptions on Dental-Radiodiagnosis Using Smartphone: Crossectional study

Supriyadi, Swasthi Prastyarini, Lusi Hidayati, Pujiana Endah Lestari

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/50-D22_1719_Supriyadi_Indonesia.pdf]

CLINICAL ARTICLE

51. Role of Ultrasound in the Detection of Lesions of the Parotid Gland in HIV Patients

Victor Dsouza, Ramya Shenoy, Bastian TS

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/51-D22_1792_Ceena_Denny_E_India.pdf]

CLINICAL ARTICLE

52. Dentocraniofacial Cephalometric and Clinical Parameter Comparison of Obstructive Sleep Apnea (OSA) and Control Group Patient Age 9–12 Years before Orthodontic Treatment

Ignatia Wulandari, Miesje Karmiati Purwanegara, Fauziah Fardizza, Krisnawati
Tarman

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/52-D22_1775_Ignatia_Wulandari_Indonesia.pdf]

CLINICAL ARTICLE

53. Self-perception oral malodor and risk predictors in a group of Thai dental patients

Supawadee Naorungroj, Supitcha Talungchit, Kaewkwan Tanthai, Benjaporn
Tantipichiankul, Ratirat Klawpai, Kanchanit Sittiyuno, Keerarat Aompornnararat

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/53-D22_1796_Supawadee_Naorungroj_Thailand.pdf]

CLINICAL ARTICLE

54. The Differences between Panoramic Mandibular Indexes in Young and Older Women on Panoramic Radiograph

Bramma Kiswanjaya, Shafira Gina Andriana, Heru Suryonegoro

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/54-D22_1776_Bramma_Kiswanjaya_Indonesia.pdf]

CLINICAL ARTICLE

55. Assessment of Denture Hygiene Knowledge and Attitude in Al Madinah AlMunawwarah

Full Text **PDF**

Pages 814-819

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/55-D22_1798_Rania_M_Moussa_KSA.pdf]

CASE REPORT

56. Odontogenic Myxofibroma: A Case Report

Chonakan Thitiyuk, Siripatra Patchanee, Narissaporn Chaiprakit

Full Text **PDF**

Pages 820-823

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/56-D22_1751_Narissaporn_Chaiprakit_Thailand.pdf]

CASE REPORT

57. A Management of Anterior Crossbite with Removable Posterior Bite Riser, Composite Inclined Plane, or Fixed Appliance

Winny Yohana, Risti Saptarini Primanti

Full Text **PDF**

Pages 824-828

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/57-D22_1705_Winny_Yohana_Indonesia.pdf]

CASE REPORT

58. Two Case Reports of Intravenous Midazolam Sedation in Patients with Down Syndrome Receiving Dental Treatment

Farah Natashah Mohd, Abdul Hadi Said, Mohd Nizamuddin Ismail

Full Text **PDF**

Pages 829-833

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/58-D22_1769_Farah_Natashah_Mohd_Malaysia.pdf]

CASE REPORT

59. Review of Surgical Exposure of Impacted Upper Canine in Complementing Orthodontic Treatment: A Case Report

Winny Yohana

Full Text **PDF**

Pages 834-838

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/59-D21_1676_Winny_Yohana_Indonesia.pdf]

REVIEW

60. Clinical Bond Failures Rate of Adhesive Precoated Orthodontic Brackets: A Systematic Review

Kitiporn Chaimaungchuen, Suwannee Luppanapornlarp, Irin Sirisoontorn

REVIEW

61. The Effect of Parents' Socioeconomic Factors on Their Willingness to Take Care of Their Children's Oral Health in Early Childhood

Dyah Nawang Palupi Pratamawari, Dewi Atikasari, Taufan Bramantoro

REVIEW

62. Bacterial Adhesion and the Role of Oral Mouthwashes in Orthodontics: A Literature Review

Valeria Díaz Cabrera, Mariano Ortiz Pizarro, Katherine Yolanda Lozano Peralta, Martin Arturo Vilela Estrada, Christian Richard Mejía Álvarez, Victor Serna Alarcón

REVIEW

63. Effect of Hyaluronic Acid Gel as an Adjunct Therapy After Scaling Root Planing in Chronic Periodontitis: A Rapid Review

Ina Hendiani, Siti Sopiati, Tyssa Maira Putrifasha

REVIEW

64. Prosthodontic Management of Ridge Resorption: An Updated Review

Sebahate Hamiti Alidema, Jadranka Bundevska, Sotir Maja, Goce Dimoski, Rajmonda Halili

REVIEW

65. Effectiveness of Twin Block Device as Upper Airway Correction in Pediatric Patients with Class II Malocclusion and Its Relationship with Muscle Contraction: A Systematic Review

Harun Achmad, Rini Sitanaya, Hans Lesmana, Arni Irawaty Djais, Rosdiana Agustin

Full Text **PDF**

Pages 873-884

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/65-D22_1785_Harun_Achmad_Indonesia.pdf]

REVIEW

66. An Analysis of Genetic Risk Factors (Chromosomes 3 and 9) and Mutation of Spike COVID-19 in the Severity and Transmission Factor

Farida Anwari, Fery Setiawan, Arif Rahman Nurdianto, Theresia Indah Budhy, Heribertus AB Tena, Heni Puspitasari

Full Text **PDF**

Pages 885-890

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/66-10-3-Analysis-D22_1783_Dian_Agustin_Wahjuningrum_Indonesia.pdf]

REVIEW

67. Comparison of Shear Bond Strength Between Uncoated and Precoated Orthodontic Brackets: A Systematic Review

Watcharawee Thanetchaloempong, Dhanabhol Riowruangsanggoon, Irin Sirisoontorn

Full Text **PDF**

Pages 891-895

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/67-D22_1726_Irin_Sirisoontorn_Thailand-1.pdf]

REVIEW

68. The Role of Static Magnetic Healing Abutment in Osteoblastic Differentiation to Reduce Marginal Crestal Bone Loss

Leonard Christiaan Nelwan, Sindy Cornelia Nelwan, Asti Meizarini, Nunthawan Nowwarote

Full Text **PDF**

Pages 896-898

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/68-10-10-The-Role-D22_1783_Dian_Agustin_Wahjuningrum_Indonesia.pdf]

REVIEW

69. Computer Aided Drug Discovery Utilization in Conservative Dentistry

Anastasia Elsa Prahasti, Tamara Yuanita, Retno Pudji Rahayu

Full Text **PDF**

Pages 899-903

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/69-10-5-Computer-D22_1783_Dian_Agustin_Wahjuningrum_Indonesia.pdf]

REVIEW

70. Hydrogel Scaffold in Pulp Dentin Complex Regeneration

Elline, Kun Ismiyatin, Theresia Indah Budhy

Full Text **PDF**

Pages 904-908

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/70-10-7-Hydrogel-D22_1783_Dian_Agustin_Wahjuningrum_Indonesia.pdf]

REVIEW

71. The Association between Periodontitis and Atherosclerotic Cardiovascular Disease: A Scoping Review

Agus Susanto, Dyah Nindita Carolina, Siti Sopiati, Ina Hendiani

Full Text **PDF**

Pages 909-915

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/71-D22_1793_D22_1795_Agus_Susanto_Indonesia.pdf]

REVIEW

72. Full Coverage Crowns for Primary anterior Teeth – A Literature Review

Parichat Pengpue, Pisaisit Chaijareenont, Varisara Sirimaharaj, Wannapa Chinadet

Full Text **PDF**

Pages 919-921

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/72-D22_1807_Wannapa_Chinadet_Thailand.pdf]

REVIEW

73. The Correlation between Oral Health Condition in Down Syndrome Children with Physical Fitness: A Literature Review

Mega Moeharyono Puteri, Tania Saskianti, Alit Rahma Estu, Barnabas Bornado, Brian Maulani, Nita Naomi

Full Text **PDF**

Pages 922-925

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/73-10-2-The-correlation-D22_1783_Dian_Agustin_Wahjuningrum_Indonesia.pdf]

REVIEW

74. Clinical Periodontal Parameter of Smokers with Periodontitis in Asia Following Periodontal Therapy: An Update Systematic Review and Meta-Analysis

Mahira Aviandiva, Dhas Salsabila Putri, Benso Sulijaya, Sri Lelyati C Masulili, Natalina, Robert Lessang

Full Text **PDF**

Pages 926-934

[http://www.jidmr.com/journal/wp-content/uploads/2022/06/74-D21_1688_Benso_Sulijaya_Indonesia.pdf]

MEDICINE

EXPERIMENTAL ARTICLE

75. Antimicrobial Effect of Structural Change on the

**75. Angiopathy as a Cause of Structural Organ Changes under
Experimental Conditions in Diabetes Mellitus**

Lesia Mateshuk-Vatseba, Iryna Savka, Maksym Tsytoivskyi, Nazar Blyshchak,
Solomiya Fedevych

Full Text **PDF**

Pages 935-941

[[http://www.jidmr.com/journal/wp-content/uploads/2022/06/75-](http://www.jidmr.com/journal/wp-content/uploads/2022/06/75-M21_1698_Tsytoivskyi_Maksym_Ukraine.pdf)

[M21_1698_Tsytoivskyi_Maksym_Ukraine.pdf](http://www.jidmr.com/journal/wp-content/uploads/2022/06/75-M21_1698_Tsytoivskyi_Maksym_Ukraine.pdf)]

EXPERIMENTAL ARTICLE

**76. The Stemness and Quiescence Maintenance of Human Iliac Bone
Marrow Mesenchymal Stem Cells by Resveratrol: An In Vitro Study**

Chairul Yahya, Mohammad Saifur Rohman, Mohammad Hidayat, Alexander
Patera Nugraha, Fedik Abdul Rantam

Full Text **PDF**

Pages 942-949

[[http://www.jidmr.com/journal/wp-content/uploads/2022/06/76-](http://www.jidmr.com/journal/wp-content/uploads/2022/06/76-FTD22_1834_Mohammad_Saifur_Rohman_Indonesia.pdf)

[FTD22_1834_Mohammad_Saifur_Rohman_Indonesia.pdf](http://www.jidmr.com/journal/wp-content/uploads/2022/06/76-FTD22_1834_Mohammad_Saifur_Rohman_Indonesia.pdf)]

EXPERIMENTAL ARTICLE

**77. Poly L-Lactic Acid (PLLA)-Collagen Coating Chitosan as a Spring-Loaded
Silo Candidate for Gastroschisis**

Prihartini Widiyanti, Fahreza Rachmat Yoviansyah, Djony Izak Rudyarjo

Full Text **PDF**

Pages 950-954

[[http://www.jidmr.com/journal/wp-content/uploads/2022/06/77-](http://www.jidmr.com/journal/wp-content/uploads/2022/06/77-MD22_1740_Prihartini_Widiyanti_Indonesia.pdf)

[MD22_1740_Prihartini_Widiyanti_Indonesia.pdf](http://www.jidmr.com/journal/wp-content/uploads/2022/06/77-MD22_1740_Prihartini_Widiyanti_Indonesia.pdf)]

CLINICAL ARTICLE

**78. Identification of Lung Cancer Metastasis from Patient Records: An
Explanatory Meta-Diagnosis**

Zeynep Ertem

Full Text **PDF**

Pages 955-958

[[http://www.jidmr.com/journal/wp-content/uploads/2022/06/78-](http://www.jidmr.com/journal/wp-content/uploads/2022/06/78-M22_1801_Zeynep_Ertem_US.pdf)

[M22_1801_Zeynep_Ertem_US.pdf](http://www.jidmr.com/journal/wp-content/uploads/2022/06/78-M22_1801_Zeynep_Ertem_US.pdf)]

share page



Characterization of Bovine Sponge Amnion (BSA) by a Novel Process for Dental Treatment

Octarina¹, Elly Munadzir^{2*}, Fathilah Abdul Razak³, Basril Abbas⁴, Helal Soekartono²,
Titien Hary Agustantina², Meircurius Dwi Condro Surboyo⁵

1. Doctoral Program, Faculty of Dental Medicine, Universitas Airlangga, Surabaya 60132, Indonesia.

2. Department of Dental Material, Faculty of Dental Medicine, Universitas Airlangga, Surabaya 60132, Indonesia.

3. Department of Oral and Craniofacial Sciences, Faculty of Dentistry, Universitas of Malaya, Kuala Lumpur 50603, Malaysia.

4. Centre for Application of Isotopes and Radiation, Jl. Lebak Bulus Raya no. 49, Jakarta 12070, Indonesia.

5. Department of Oral Medicine, Faculty of Dental Medicine, Universitas Airlangga, Surabaya 60132, Indonesia.

Abstract

One of dental treatments in dentistry is tooth extraction. This treatment will cause injury after extraction and biomaterial is needed to accelerate wound healing. Bovine amniotic membrane is a biomaterial with collagen content that can accelerate wound healing. However, in the process of application, the membrane will tear easily and is difficult to apply.

The purpose of this study was to fabricate the bovine amniotic membrane into bovine sponge amnion (BSA) and characterize it. BSA was derived from bovine amniotic membrane with the addition of gelatin which then underwent freeze-drying process and lyophilization. BSA was then characterized using Fourier Transform Infrared Spectroscopy (FTIR), scanning electron microscopy (SEM), and swelling test.

The result of characterization using FTIR indicated that amide A, amide B, amide I, amide II, and triple helix of collagen groups were shown in the FTIR graphs. The surface characterization using SEM showed a mean porosity size of 304.866 μm with a porosity proportion of 64%. The swelling analysis showed the proportion of swelling was 1320 times.

The BSA has an ideal pore size, functional group and swelling ability for biomaterial candidate in dentistry.

Experimental article (J Int Dent Med Res 2022; 15(2): 479-484)

Keywords: Biomaterials; biomedical; bovine sponge amnion; bovine amniotic membrane; human & health.

Received date: 15 January 2022

Accept date: 02 March 2022

Introduction

Post extraction complications was ranked second for dental problems in Indonesia.¹ Complications that occur after tooth extraction will cause large wounds that have the potential to cause critically size bone defects which causes bone healing is not optimal.² The wound healing process is a complex process that involves bio-cellular and biochemical activities that run continuously.³ The wound healing process will involve activation and synchronization of intracellular, intercellular, and extracellular

mechanisms, including coagulation and inflammatory events, accumulation of fibrous tissue, collagen deposition, epithelialization, wound contraction, tissue granulation, and remodelling.⁴ Wounds caused by extraction require materials that can accelerate wound healing.⁵ One of the requirements for wound healing materials is that they can interact well with biological systems.⁶

Amniotic membrane is a biomaterial that has anti-inflammatory and antimicrobial properties that play a role in repairing and regenerating damaged tissue. Amniotic membrane has growth factors such as epidermal growth factor (EGF), transforming growth factor alpha (TGF- α), keratinocyte growth factor (KGF), hepatocyte growth factor (HGF), basic fibroblast growth factor (bFGF), transforming growth factor beta (TGF- β), keratinocyte growth factor receptor (KGFR), and hepatocyte growth factor receptor (HGFR).⁷ The extracellular matrix of the amniotic membrane consists of collagen types I, II, III, IV,

*Corresponding author:

Elly Munadzir
Department of Dental Material
Faculty of Dental Medicine, Universitas Airlangga.
St. Prof. Dr. Moestopo 47 Surabaya, Indonesia 60132.
Telp. 6231-5030255.
E-mail: elly-m@fkg.unair.ac.id

laminin, nidogen, fibronectin, and proteoglycans.⁸ One source of the amniotic membrane is bovine. Some of the advantages of bovine amniotic membrane are the ease of legality, ethics, and that it is not affected by ritual beliefs as in the human amniotic membrane.⁹ Bovine amniotic membrane can be produced in large quantities on farms, with good food quality so as to produce an amniotic membrane with good elasticity and thickness. The bovine amniotic membrane is rich in growth factors and collagen, which can accelerate the re-epithelialization process in wound healing.¹⁰

The limitation of the amniotic membrane is its low mechanical properties so that it is easily degraded.¹¹ This property also causes the amniotic membrane to tear easily upon application.¹² Ease of degradation of the amniotic membrane, will affect the integrity of the structure which is important when healing the extraction wounds with long-term treatment.¹³ To solve this problem, modification of the amniotic membrane is required.¹⁴ In dental treatment, the amniotic membrane can be combined with gelatin. Gelatin acts as a binder to form a structure and improve mechanical properties without losing the characteristics of the amniotic membrane.¹⁵ Fabrication the amniotic membrane in the form of a sponge is carried out to provide ease of use. The amniotic membrane with freeze-drying treatment will form a porous structure that resembles a sponge.¹⁶ The porous structure of the sponge helps the gas exchange and absorption of wound exudate. This structure also plays a role as a place for cell growth and vascularization as well as carrying medicaments to accelerate the wound healing process.^{17,18}

Material characterization will provide information about the physical and chemical properties of the material. The physical and chemical properties of a material can be influenced by the size and components of the material.¹⁹ The functional group, surface morphology, and swelling are important characteristics of a biomaterial to be used for biomedical applications. The bovine amniotic membrane contains collagen which plays a role in wound regeneration. For ease of application, it is fabricated into a bovine sponge amnion. Based on this, this study aimed to fabricate a bovine amniotic membrane in the form of a bovine sponge amnion (BSA) then analyze its characteristics using Fourier Transform Infrared

Spectroscopy (FTIR), scanning electron microscopy (SEM), and the percentage of swelling as candidates for dental treatment.²⁰

Materials and methods

Fabrication of bovine amnion sponge specimen

This research is a laboratory experimental in vitro study Bovine amniotic membrane was obtained from the Installation Center for Biomaterials Bank Network Dr. Soetomo Surabaya. The fresh bovine amniotic membrane was cleaned in order to wash away the blood clots that stucked on the membrane and rinsed with saline solution containing antibiotics for four times.




Figure 1. Bovine sponge amnion (BSA) fabrication scheme.

The inner amniotic membrane was separated from the chorion (Figure 1A) and put into deep freezer at -80°C for 24 hours (Figure 1B). The membrane was then freeze-dried for eight hours (Figure 1C). After the freeze-drying process was carried out, the amniotic membrane would become a dry thin sheet. The next step was cutting the dry thin sheet of bovine amniotic membrane into small pieces (Figure 1D), then crushed with a mortar and pestle until it became smaller pieces of particles (Figure 1E). The particles were sieved with a sieving machine until it was $250\text{ }\mu\text{m}$ in size (Figure 1F). The bovine amniotic membrane was then added with water

(H₂O) in a ratio of 1:1 to form the amniotic slurry (Figure 1G). The amniotic slurry was then mixed with bovine gelatin in 1:1 ratio and processed by freeze-drying method for 2×24 hours (Figure 1H and 1I). When the amniotic sponge was obtained, the sponge was cut into a size of 1.5×5 mm (Figure 1J).

Fourier transform infrared spectroscopy (FTIR) analysis

FTIR analysis was conducted at the Center for Isotopes and Radiation at the Nuclear Power Agency (PAIR BATAN). The sample of bovine sponge amnion was added with KBr powder (*Sigma Aldrich, Germany*) in a ratio of 1:50, crushed using a mortar and pestle. After it was mixed homogeneously, the sample was put into a sample compartment and then tested using an FTIR device for characterization. Functional group BSA analysis was using IR Prestige-21 (*Shimadzu, Japan*). The analysis in this study used the measurement of the percentage of transmittance with a resolution of 4.0 and at a wavelength of 4000 to 400 cm⁻¹.

Scanning electron microscope (SEM) analysis

SEM analysis was conducted in Faculty of Mathematics and Natural Sciences, Bandung Institute of Technology (FMIPA ITB). BSA samples were prepared with a gold coating before testing. The prepared samples were then put into chamber and vacuumed. The analysis of sample surface, morphology, and components were characterized using SEM (*JEOL, JSM 6510-LA*). The pore size in this study was measured 30 times in different places with a 50x SEM magnification with Image-J software. The percentage of porosity was analyzed by using SEM images with a magnification of 100x with Origin Pro 2021 software.

Swelling analysis

Swelling analysis was conducted at DMTCORE FKG USAKTI. In the preliminary test, the swelling presentation on the sample was carried out for 1, 10, 15, 20 minutes, and 24 hours. However, the results of the swelling presentation were not significantly different. In this study, the 10-minute immersion showed the best results and used for swelling analysis calculation.

The initial weight (W_w) of the BSA, was weighed before immersion using an analytical balance (*Fujitsu FSAR-210, Japan*). BSA samples were immersed in 5 mL phosphate buffer solution

(PBS). After immersion, the samples were gently placed on filter paper with a diameter of 9 mm (*Whatman, US*) for three seconds to absorb excess water to measure the final weight (W_d). Swelling analysis was calculated using following equation:

$$\text{Swelling} = \frac{W_w - W_d}{W_d} \times 100\%$$

Results

FTIR analysis

As shown in Figure 2 that OH, Amide A, Amide B, Amide I, Amide II, Amide III, and triple helix of collagen functional group peaks can be observed in the graphs (OH = 3307.92 cm⁻¹; Amide Peak A = 3074.53 cm⁻¹; Amide peak B = 2929.87 cm⁻¹; Amide peak I = 1674.21 cm⁻¹; Amide Peak II = 1584.84 cm⁻¹; Amide Peak III = 1240.23 cm⁻¹; Triple helix of collagen: 1454.53 cm⁻¹).




Figure 2. Analysis of bovine sponge amnion using FTIR.

The amide A absorption band occurred in the presence of an N-H strain bond from the amide group associated with a hydrogen bond and OH group. The absorption of the amide-I protein absorption band is associated with the C = O stretching mode. Amide-II protein absorption band is associated with the N – H bending mode and the C – N stretching mode. The amide band III is produced from a combination of the C – N stretching phase and the N – H plane bending, with some contribution from the C – C stretching and the C = O bending vibration.

Discussion

In this study, a bovine amniotic membrane was fabricated with the addition of gelatin to form a bovine sponge amnion. The gelatin used was a ready-made bovine gelatin. The addition of gelatin in this study was carried out with the aim of bonding the biomaterial so that it became solid, producing a firm sponge that did not tear easily during biomedical applications. This study also aimed to analyze the interaction between collagen and gelatin at the time of manufacture. This material characterization can be analyzed using FTIR, SEM, and swelling procedures.

Collagen is a protein substance in the form of fiber which is the main part of connective tissue needed for wound healing, scar tissue formation, and formation of bone matrices. Collagen is a key component in wound healing. Initially, type III collagen is more dominant, which then is replaced by type I collagen. After 20 days, the granulation tissue is replaced by collagen and bone begins to form at the base and margins of the extraction socket.²⁰ The bovine amniotic membrane has the main composition of collagen which plays a role in wound healing. This study was conducted to analyze what components were contained during the fabrication of BSA. The organic components contained in BSA are indicated by the peak of the functional group which can be analyzed using FTIR.

FTIR analysis was carried out to ensure that the compound produced was gelatin by comparing the results of the sample spectrum and the gelatin standard. The structure of gelatin, like most proteins, has carbonyl, amine, and hydroxyl group. Gelatin gives rise to the typical IR absorption of amide A at wavenumbers 3600–2300 cm^{-1} , amide I at 1636–1661 cm^{-1} , amide II at 1560–1335 cm^{-1} , and amide III at 1300–1200 cm^{-1} (Maryam *et al.* 2019).²¹ This FTIR image supports the findings of this study and shows that the gelatin and collagen in the bovine amniotic membrane bind well in the presence of the amide III functional group.²² Collagen band characteristics shown in the graphs imply that collagen is retained at each decellularization process.⁴

Triple helix of collagen is type I collagen. In particular, among the various types of collagens, type I collagen is the most abundant extracellular matrix component and can be used as a scaffold,

cell migration promotion, wound healing, and tissue regeneration. Resorbable collagen membranes have been used in an integrated manner in tissue regeneration and guided bone regeneration procedures due to its biocompatibility and ability to promote wound healing. The composition of collagen in the fabrication of BSA shows that this biomaterial has potential as a material used for dental applications.

SEM analysis

The SEM analysis results showed irregular pore of BSA samples with magnification was 50x and 100x (Figure 3A and 3B). The pore is a combination of macro and micro pore sizes. The results of this study showed the smallest pore was 133.101 μm and the largest was 605.803 μm with an average pore of 304.866 μm . The porosity percentage in BSA showed 64% porosity (Figure 3C). Pore structure in biomaterial is needed and important in wound healing process where gas permeability, absorption of wound exudates, cell development, exchange of nutrients, and metabolites will occur inside the pores.¹⁸ The microstructure, pore size, and porosity percentage of a biomaterial can be analyzed using SEM. The combination of macro and micro pores sizes in this study can be caused or influenced by the fabrication method. Freeze-drying method creates gelatin sponges with large and clearly visible pores.²³




Figure 3. SEM overview of the morphology and microstructure of bovine sponge amnion. (A) 50x magnification, (B) 100x magnification; (C) Image of 3D color fill surface overview of bovine sponge amnion porosity percentage.

Image-J is a software for analyzing digital images that has been widely used in medicine and biology.²⁴ In this study, the magnification was at 50x SEM image to calculate pore size. These results are in line with the study results of Murphy *et al.* (2010) that the ideal mean size of the pores for cell proliferation and migration in bone tissue formation is 325 μm . Minimum pore size for bone growth is in the optimal range of 100–135 μm . The pore size exceeding 300 μm will play a role in the process of vascularization.²⁵ Combined pores larger than 300 μm lead to direct osteogenesis while pores smaller than 300 μm can promote osteochondral ossification. As a scaffold, the pores on sponges have an oval structure with an ideal diameter ranging from 100 to 500 μm .²⁶

To allow cell growth to occur, the scaffold must have a porous structure with a porosity value of 80-90% to support regeneration. The greater the percentage of scaffold porosity, the faster the cell proliferation and differentiation.²⁷ In this study, the percentage porosity analysis was less than 80%. This can be affected by the freeze temperature and the freeze-drying method used.²⁸ This porosity percentage value can be influenced by the location where SEM images were taken. The definition of the size of the porosity is limited to the pixel size of the SEM images. Image pixel size affects the smallest measurable scale value and causes large deviations in the mean of porosity percentage. Biomaterials used for biomedicine generally have soft pores that formed a three-dimensional network.²⁹ Analysis with the use of Micro-computed X-ray tomography (Micro-CT) can improve the accuracy of the calculation of the percentage of porosity in biomaterials with porous structures. This analysis can also provide three-dimensional characterization of complex multi-porous microarchitectures.³⁰ After all, SEM results in this study indicate that this material has a combination of macro and micro size pore structures that can act as a material for wound healing in dental treatment.

	Before Immersion (gr)	After Immersion (gr)	Swelling percentage (%)
Specimen 1	0.0019	0.0270	1321
Specimen 2	0.0017	0.0223	1211
Specimen 3	0.0017	0.0260	1429
Mean \pm SD			1320 \pm 108

Table 1. The percentage of swelling after immersion in PBS solution.

The swelling ability plays an important role in tissue healing process, where the transfer of nutrients and body fluids into the biomaterial will be maximized if a biomaterial has good swelling

capability.³¹ In this study, when the sample was put into the PBS solution, the sample immediately expanded and after ten minutes, the mean swelling percentage reached 1320%. This result is in line with that conducted by Angulo and Sobral (2016) who used a incorporation of gelatin to fabricate scaffolds for tissue engineering. Gelatin has the ability to increase the swelling ability of the biomaterial.³² BSA contains collagen which causes this biomaterial to have high hydrophilicity properties. Hydrophilicity is the ability of collagen to bind water through hydrogen bonds, so that collagen is more easily dissolved in water. If the hydrogen bonds meet water (H_2O), then there is an intermolecular attraction at the H atom which is attached to the very electronegative atoms (N, O, and F).³³ A study by Cheng *et al.* (2017) shows that collagen sponges show a higher water absorption rate this can be caused with the presence of a porous collagen sponge structure and water retention of the collagen fibers.³⁴

Research conducted by Guclu *et al* (2017) increases the swelling capacity of a biomaterial that can occur up to 1200% to 1900% as a candidate biomaterial for bone healing.³⁵⁻³⁷ This shows that swelling is also an important feature for developing suitable tissue engineering constructs for bone defect regeneration.²⁷ The study by Indrawati *et al.* (2019) shows that BSA increase the bone morphogenetic protein-2 (BMP-2), as a marker of osteogenesis from the first day to the seventh day after the application in post-extraction of rat teeth.¹² This swelling presentation has been able to make BSA as a material for wound healing not only of soft tissue but also of hard tissue.

Conclusions

Based on the results of this study, it shows the potential characteristics of BSA as a biomaterial that can improve wound healing in dental treatment. Further studies are recommended for in vivo animal studies to analyze the effect of biomaterials on wound healing and regeneration after tooth extraction.

Acknowledgments

The author would like to thank the Installation of Tissue Bank and Cells of RSUD Dr. Soetomo Surabaya for the bovine sponge amnion (BSA). The author also thanks all laboratories that have supported this study, especially the DMTCORE Laboratory of Trisakti University. Last but not least, the author would like to thank Ms. Agnes Victoria, S.K.G for the lovely illustrations in this paper.

Declaration of Interest

The authors declare that there are no conflicts of interest.

Ethical policy and institutional review board statement

Ethical clearance had been obtained from the Ethics Commission of the Faculty of Dental Medicine, Airlangga University, Surabaya (No. 293/KKEPK.FKG/XII/2016).

Swelling analysis

In this study, the 10-minute immersion showed the results of the swelling percentage of 1321%, 1211%, and 1429% which are shown in Table 1.

References

- Riskesdas. Laporan Riskesdas 2018. Badan Penelitian dan Pengembangan Kesehatan. Kementerian Kesehatan Republik Indonesia. 2018:181-222
- Walsh DP, Raftery RM, Chen G, Heise A, O'Brien FJ, Cryan SA. Rapid healing of a critical-sized bone defect using a collagen-hydroxyapatite scaffold to facilitate low dose, combinatorial growth factor delivery. *J Tissue Eng Regen Med*. 2019;13:1843-53.
- Eming S, Paul M, Tomic-Camic M. Wound repair and regeneration: Mechanisms, signaling, and translation. *Sci Transl Med*. 2014;6:1-36.
- Ballesteros ACV, Puello HRS, Lopez-Garcia JA, Bernal-Ballen A, Mosquera DLN, Forero DMM, et al. Bovine decellularized amniotic membrane: Extracellular matrix as scaffold for mammalian skin. *Polymers (Basel)*. 2020;12.
- Pei Y, Ye D, Zhao Q, Wang X, Zhang C, Huang W, et al. Effectively promoting wound healing with cellulose/gelatin sponges constructed directly from a cellulose solution. *J Mater Chem B*. 2015;3:7518-28.
- O'Brien FJ. Biomaterials & scaffolds for tissue engineering. *Mater Today*. 2011;14:88-95.
- Koizumi N, Inatomi T, Sotozono C, Fullwood NJ, Quantock AJ, Kinoshita S. Growth factor mRNA and protein in preserved human amniotic membrane. *Curr Eye Res*. 2000;20:173-7.
- Niknejad H, Peirovi H, Jorjani M, Ahmadiani A, Ghanavi J, Seifalian M, et al. Properties Of The Amniotic Membrane For Potential Use In Tissue Engineering. *Eur Cells Mater*. 2008:88-99.
- Putra NHD, Matulatan F, Wibowo MD, Danardono E. Effects of dried bovine amniotic membrane as prosthetics of abdominal fascial defect closure observed by the expression of platelet-derived growth factor in Rattus norvegicus wistar strain. *Syst Rev Pharm*. 2020;11:987-91.
- Yang Z, Cui Q, An R, Wang J, Song X, Shen Y, et al. Comparison of microbiomes in ulcerative and normal mucosa of recurrent aphthous stomatitis (RAS)-affected patients. *BMC Oral Health*. 2020;20:1-8.
- Cai M, Zhang J, Guan L, Zhao M. Novel implantable composite biomaterial by fibrin glue and amniotic membrane for ocular surface reconstruction. *J Mater Sci Mater Med*. 2015;26.
- Indrawati DW, Munadzirah E, Sulisetyawati TIB, El Fadhlallah PM. Sponge amnion potential in post tooth extraction wound healing by interleukin-6 and bone morphogenetic protein-2 expression analysis: An animal study. *Dent Res J (Isfahan)*. 2019;16:283-8.
- Dimitriou R, Mataliotakis GI, Calori GM, Giannoudis P V. The role of barrier membranes for guided bone regeneration and restoration of large bone defects: Current experimental and clinical evidence. *BMC Med*. 2012;10:1-24.
- Dadkhah Tehrani F, Firouzeh A, Shabani I, Shabani A. A Review on Modifications of Amniotic Membrane for Biomedical Applications. *Front Bioeng Biotechnol*. 2021;8:1-25.
- Zhang Q, Qian C, Xiao W, Zhu H, Guo J, Ge Z, et al. Development of a visible light, cross-linked GelMA hydrogel containing decellularized human amniotic particles as a soft tissue replacement for oral mucosa repair. *RSC Adv*. 2019;9:18344-52.
- Kang M, Choi S, Cho Lee AR. Effect of freeze dried bovine amniotic membrane extract on full thickness wound healing. *Arch Pharm Res*. 2013;36:472-8.
- Cheng H, Li C, Jiang Y, Wang B, Wang F, Mao Z, et al. Facile preparation of poly sacharide-based sponges and its potential application in wound dressing. *J Mater Chem B*. 2013;1-8.
- Long H, Ma K, Xiao Z, Ren X. Preparation and characteristics of gelatin sponges crosslinked by microbial transglutaminase. *PeerJ*. 2017;1-18.
- Khalid P, Suman VB, Shameema S, Vinodini NA, Shela RP, Chatterjee PK. Analytical methods for nanomaterial characterization. *Int J Nano Dimens Anal methods Nanomater Charact*. 2014;5:309-20.
- Octarina, Munadzirah E, Razak FA. Physical Modification of Bovine Amniotic Membrane for Dental Application. *Journal of International Dental and Medical Research*. 2021;14(4):1425-8.
- Steiner GG, Francis W, Burrell R, Kallet MP, Steiner DM, Macias R. The healing socket and socket regeneration. *Compend Contin Educ Dent*. 2008;29:1-14.
- Maryam S, Effendi N, Kasmah K. Produksi dan Karakterisasi Gelatin dari Limbah Tulang Ayam dengan Menggunakan Spektrofotometer Ftir (Fourier Transform Infra Red). *Maj Farm*. 2019;15:96.
- Muyonga JH, Cole CGB, Duodu KG. Fourier transform infrared (FTIR) spectroscopic study of acid soluble collagen and gelatin from skins and bones of young and adult Nile perch (*Lates niloticus*). *Food Chem*. 2004;86:325-32.
- Kabiri M, Emami SH, Rafinia M, Tahriri M. Preparation and characterization of absorbable hemostat crosslinked gelatin sponges for surgical applications. *Curr Appl Phys*. 2011;11:457-61.
- uarez-Arnedo A, Figueroa FT, Clavijo C, Arbeláez P, Cruz JC, Muñoz-Camargo C. An image J plugin for the high throughput image analysis of in vitro scratch wound healing assays Alejandra. *bioRxiv*. 2020;1-21.
- Murphy CM, Haugh MG, Brien FJO. Biomaterials The effect of mean pore size on cell attachment , proliferation and migration in collagen – glycosaminoglycan scaffolds for bone tissue engineering. *Biomaterials*. 2010;31:461-6.
- León-Mancilla BH, Araiza-Téllez MA, Flores-Flores JO, Piña-Barba MC. Physico-chemical characterization of collagen scaffolds for tissue engineering. *J Appl Res Technol*. 2016;14:77-85.
- Tohamy KM, Mabrouk M, Soliman IE, Beherei HH, Aboelnasr MA. Novel alginate/hydroxyethyl cellulose/hydroxyapatite composite scaffold for bone regeneration: In vitro cell viability and proliferation of human mesenchymal stem cells. *Int J Biol Macromol*. 2018;112:448-60.
- Oikonomopoulou VP, Krokida MK, Karathanos VT. The influence of freeze drying conditions on microstructural changes of food products. *Procedia Food Sci*. 2011;1:647-54.
- Ahumada M, Jacques E, Calderon C, Martínez-Gómez F. Porosity in biomaterials: A key factor in the development of applied materials in biomedicine. *Handb Ecomater*. 2019;5:3503-22.
- Nair M, Shepherd JH, Best SM, Cameron RE. MicroCT analysis of connectivity in porous structures: Optimizing data acquisition and analytical methods in the context of tissue engineering. *J R Soc Interface*. 2020;17:1-16.
- Maji K, Dasgupta S, Pramanik K, Bissoyi A. Preparation and Evaluation of Gelatin-Chitosan-Nanobioglass 3D Porous Scaffold for Bone Tissue Engineering. *Int J Biomater*. 2016;2016:1-14.
- López Angulo DE, do Amaral Sobral PJ. Characterization of gelatin/chitosan scaffold blended with aloe vera and snail mucus for biomedical purpose. *Int J Biol Macromol*. 2016;92:645-53.
- Peng L, Xiang RC, Jia WW, Dong XX, Wang GE. Preparation and evaluation of porous chitosan/collagen scaffolds for periodontal tissue engineering. *J Bioact Compat Polym*. 2006;21:207-20.
- Cheng X, Shao Z, Li C, Yu L, Raja MA, Liu C. Isolation, characterization and evaluation of collagen from jellyfish Rhopilema esculentum kishinouye for use in hemostatic applications. *PLoS One*. 2017;12:1-21.
- Guclu ZA, Hurt AP, Ohia L, Coleman NJ. Composite Sponges for in Situ Alveolar Bone Regeneration Following Tooth Extraction. *Acta Phys Pol A*. 2017;131:580-3.
- Munadzirah E, Soekartono RH, Rafli RBV, Yuliati A, Kartikasari N. Effect of rSLPI Amnion Membrane Application on Incision Wound of Rattus Norvegicus in Collagen and VEGF Expression. *Journal of International Dental and Medical Research*. 2017;10(2):239-43.

Characterization of Bovine Sponge Amnion (BSA) by a Novel Process for Dental Treatment

by Octarina FKG

Submission date: 26-Jan-2024 09:07AM (UTC+0700)

Submission ID: 2278641574

File name: haracterization-D22_1783_Dian_Agustin_Wahjuningrum_Indonesia.pdf (1.4M)

Word count: 4501

Character count: 24056

10

Characterization of Bovine Sponge Amnion (BSA) by a Novel Process for Dental Treatment

Octarina¹, Elly Munadzir^{2*}, Fathilah Abdul Razak³, Basril Abbas⁴, Helal Soekartono²,
Titien Hary Agustantina², Meircurius Dwi Condro Surboyo⁵

1. Doctoral Program, Faculty of Dental Medicine, Universitas Airlangga, Surabaya 60132, Indonesia.

2. Department of Dental Material, Faculty of Dental Medicine, Universitas Airlangga, Surabaya 60132, Indonesia.

3. Department of Oral and Craniofacial Sciences, Faculty of Dentistry, Universitas of Malaya, Kuala Lumpur 50603, Malaysia.

4. Centre for Application of Isotopes and Radiation, Jl. Lebak Bulus Raya no. 49, Jakarta 12070, Indonesia.

5. Department of Oral Medicine, Faculty of Dental Medicine, Universitas Airlangga, Surabaya 60132, Indonesia.

Abstract

One of dental treatments in dentistry is tooth extraction. This treatment will cause injury after extraction and biomaterial is needed to accelerate wound healing. Bovine amniotic membrane is a biomaterial with collagen content that can accelerate wound healing. However, in the process of application, the membrane will tear easily and is difficult to apply.

The purpose of this study was to fabricate the bovine amniotic membrane into bovine sponge amnion (BSA) and characterize it. BSA was derived from bovine amniotic membrane with the addition of gelatin which then underwent freeze-drying process and lyophilization. BSA was then characterized using Fourier Transform Infrared Spectroscopy (FTIR), scanning electron microscopy (SEM), and swelling test.

The result of characterization using FTIR indicated that amide A, amide B, amide I, amide II, and triple helix of collagen groups were shown in the FTIR graphs. The surface characterization using SEM showed a mean porosity size of 304.866 μm with a porosity proportion of 64%. The swelling analysis showed the proportion of swelling was 1320 times.

The BSA has an ideal pore size, functional group and swelling ability for biomaterial candidate in dentistry.

Experimental article (J Int Dent Med Res 2022; 15(2): 479-484)

Keywords: Biomaterials; biomedical; bovine sponge amnion; bovine amniotic membrane; human & health.

Received date: 15 January 2022

Accept date: 02 March 2022

Introduction

Post extraction complications was ranked second for dental problems in Indonesia.¹ Complications that occur after tooth extraction will cause large wounds that have the potential to cause critically size bone defects which causes bone healing is not optimal.² The wound healing process is a complex process that involves bio-cellular and biochemical activities that run continuously.³ The wound healing process will involve activation and synchronization of intracellular, intercellular, and extracellular

mechanisms, including coagulation and inflammatory events, accumulation of fibrous tissue, collagen deposition, epithelialization, wound contraction, tissue granulation, and remodelling.⁴ Wounds caused by extraction require materials that can accelerate wound healing.⁵ One of the requirements for wound healing materials is that they can interact well with biological systems.⁶

Amniotic membrane is a biomaterial that has anti-inflammatory and antimicrobial properties that play a role in repairing and regenerating damaged tissue. Amniotic membrane has growth factors such as epidermal growth factor (EGF), transforming growth factor alpha (TGF- α), keratinocyte growth factor (KGF), hepatocyte growth factor (HGF), basic fibroblast growth factor (bFGF), transforming growth factor beta (TGF- β), keratinocyte growth factor receptor (KGR), and hepatocyte growth factor receptor (HGFR).⁷ The extracellular matrix of the amniotic membrane consists of collagen types I, II, III, IV,

*Corresponding author:

Elly Munadzir
Department of Dental Material
Faculty of Dental Medicine, Universitas Airlangga,
St. Prof. Dr. Moestopo 47 Surabaya, Indonesia 60132.
Telp. 8231-5030255.
E-mail: elly-m@fkg.unair.ac.id

laminin, nidogen, fibronectin, and proteoglycans.⁸ One source of the amniotic membrane is bovine. Some of the advantages of bovine amniotic membrane are the ease of legality, ethics, and that it is not affected by ritual beliefs as in the human amniotic membrane.⁹ Bovine amniotic membrane can be produced in large quantities on farms, with good food quality so as to produce an amniotic membrane with good elasticity and thickness. The bovine amniotic membrane is rich in growth factors and collagen, which can accelerate the re-epithelialization process in wound healing.¹⁰

The limitation of the amniotic membrane is its low mechanical properties so that it is easily degraded.¹¹ This property also causes the amniotic membrane to tear easily upon application.¹² Ease of degradation of the amniotic membrane, will affect the integrity of the structure which is important when healing the extraction wounds with long-term treatment.¹³ To solve this problem, modification of the amniotic membrane is required.¹⁴ In dental treatment, the amniotic membrane can be combined with gelatin. Gelatin acts as a binder to form a structure and improve mechanical properties without losing the characteristics of the amniotic membrane.¹⁵ Fabrication the amniotic membrane in the form of a sponge is carried out to provide ease of use. The amniotic membrane with freeze-drying treatment will form a porous structure that resembles a sponge.¹⁶ The porous structure of the sponge helps the gas exchange and absorption of wound exudate. This structure also plays a role as a place for cell growth and vascularization as well as carrying medicaments to accelerate the wound healing process.^{17,18}

Material characterization will provide information about the physical and chemical properties of the material. The physical and chemical properties of a material can be influenced by the size and components of the material.¹⁹ The functional group, surface morphology, and swelling are important characteristics of a biomaterial to be used for biomedical applications. The bovine amniotic membrane contains collagen which plays a role in wound regeneration. For ease of application, it is fabricated into a bovine sponge amnion. Based on this, this study aimed to fabricate a bovine amniotic membrane in the form of a bovine sponge amnion (BSA) then analyze its characteristics using Fourier Transform Infrared

Spectroscopy (FTIR), scanning electron microscopy (SEM), and the percentage of swelling as candidates for dental treatment.²⁰

Materials and methods

Fabrication of bovine amnion sponge specimen

This research is a laboratory experimental in vitro study. Bovine amniotic membrane was obtained from the Installation Center for Biomaterials Bank Network Dr. Soetomo Surabaya. The fresh bovine amniotic membrane was cleaned in order to wash away the blood clots that stuck on the membrane and rinsed with saline solution containing antibiotics for four times.




Figure 1. Bovine sponge amnion (BSA) fabrication scheme.

The inner amniotic membrane was separated from the chorion (Figure 1A) and put into deep freezer at -80°C for 24 hours (Figure 1B). The membrane was then freeze-dried for eight hours (Figure 1C). After the freeze-drying process was carried out, the amniotic membrane would become a dry thin sheet. The next step was cutting the dry thin sheet of bovine amniotic membrane into small pieces (Figure 1D), then crushed with a mortar and pestle until it became smaller pieces of particles (Figure 1E). The particles were sieved with a sieving machine until it was $250\text{ }\mu\text{m}$ in size (Figure 1F). The bovine amniotic membrane was then added with water

(H₂O) in a ratio of 1:1 to form the amniotic slurry (Figure 1G). The amniotic slurry was then mixed with bovine gelatin in 1:1 ratio and processed by freeze-drying method for 2×24 hours (Figure 1H and 1I). When the amniotic sponge was obtained, the sponge was cut into a size of 1.5×5 mm (Figure 1J).

Fourier transform infrared spectroscopy (FTIR) analysis

FTIR analysis was conducted at the Center for Isotopes and Radiation at the Nuclear Power Agency (PAIR BATAN). The sample of bovine sponge amnion was added with KBr powder (*Sigma Aldrich, Germany*) in a ratio of 1:50, crushed using a mortar and pestle. After it was mixed homogeneously, the sample was put into a sample compartment and then tested using an FTIR device for characterization. Functional group BSA analysis was using IR Prestige-21 (*Shimadzu, Japan*). The analysis in this study used the measurement of the percentage of transmittance with a resolution of 4.0 and at a wavelength of 4000 to 400 cm⁻¹.

Scanning electron microscope (SEM) analysis

SEM analysis was conducted in Faculty of Mathematics and Natural Sciences, Bandung Institute of Technology (FMIPA ITB). BSA samples were prepared with a gold coating before testing. The prepared samples were then put into chamber and vacuumed. The analysis of sample surface, morphology, and components were characterized using SEM (*JEOL, JSM 6510-LA*). The pore size in this study was measured 30 times in different places with a 50x SEM magnification with Image-J software. The percentage of porosity was analyzed by using SEM images with a magnification of 100x with Origin Pro 2021 software.

Swelling analysis

Swelling analysis was conducted at DMTCORE FKG USAKTI. In the preliminary test, the swelling presentation on the sample was carried out for 1, 10, 15, 20 minutes, and 24 hours. However, the results of the swelling presentation were not significantly different. In this study, the 10-minute immersion showed the best results and used for swelling analysis calculation.

The initial weight (Ww) of the BSA, was weighed before immersion using an analytical balance (*Fujitsu FSAR-210, Japan*). BSA samples were immersed in 5 mL phosphate buffer solution

(PBS). After immersion, the samples were gently placed on filter paper with a diameter of 9 mm (*Whatman, US*) for three seconds to absorb excess water to measure the final weight (Wd). Swelling analysis was calculated using following equation:

$$\text{Swelling} = \frac{Ww - Wd}{Wd} \times 100\%$$

Results

FTIR analysis

As shown in Figure 2 that OH, Amide A, Amide B, Amide I, Amide II, Amide III, and triple helix of collagen functional group peaks can be observed in the graphs (OH = 3307.92 cm⁻¹; Amide Peak A = 3074.53 cm⁻¹; Amide peak B = 2929.87 cm⁻¹; Amide peak I = 1674.21 cm⁻¹; Amide Peak II = 1584.84 cm⁻¹; Amide Peak III = 1240.23 cm⁻¹; Triple helix of collagen: 1454.53 cm⁻¹).




Figure 2. Analysis of bovine sponge amnion using FTIR.

The amide A absorption band occurred in the presence of an N-H strain bond from the amide group associated with a hydrogen bond and OH group. The absorption of the amide-I protein absorption band is associated with the C = O stretching mode. Amide-II protein absorption band is associated with the N – H bending mode and the C – N stretching mode. The amide band III is produced from a combination of the C – N stretching phase and the N – H plane bending, with some contribution from the C – C stretching and the C = O bending vibration.

Discussion

In this study, a bovine amniotic membrane was fabricated with the addition of gelatin to form a bovine sponge amnion. The gelatin used was a ready-made bovine gelatin. The addition of gelatin in this study was carried out with the aim of bonding the biomaterial so that it became solid, producing a firm sponge that did not tear easily during biomedical applications. This study also aimed to analyze the interaction between collagen and gelatin at the time of manufacture. This material characterization can be analyzed using FTIR, SEM, and swelling procedures.

Collagen is a protein substance in the form of fiber which is the main part of connective tissue needed for wound healing, scar tissue formation, and formation of bone matrices. Collagen is a key component in wound healing. Initially, type III collagen is more dominant, which then is replaced by type I collagen. After 20 days, the granulation tissue is replaced by collagen and bone begins to form at the base and margins of the extraction socket.²⁰ The bovine amniotic membrane has the main composition of collagen which plays a role in wound healing. This study was conducted to analyze what components were contained during the fabrication of BSA. The organic components contained in BSA are indicated by the peak of the functional group which can be analyzed using FTIR.

FTIR analysis was carried out to ensure that the compound produced was gelatin by comparing the results of the sample spectrum and the gelatin standard. The structure of gelatin, like most proteins, has carbonyl, amine, and hydroxyl group. Gelatin gives rise to the typical IR absorption of amide A at wavenumbers 3600–2300 cm^{-1} , amide I at 1636–1661 cm^{-1} , amide II at 1560–1335 cm^{-1} , and amide III at 1300–1200 cm^{-1} (Maryam *et al.* 2019).²¹ This FTIR image supports the findings of this study and shows that the gelatin and collagen in the bovine amniotic membrane bind well in the presence of the amide III functional group.²² Collagen band characteristics shown in the graphs imply that collagen is retained at each decellularization process.⁴

Triple helix of collagen is type I collagen. In particular, among the various types of collagens, type I collagen is the most abundant extracellular matrix component and can be used as a scaffold,

12

cell migration promotion, wound healing, and tissue regeneration. Resorbable collagen membranes have been used in an integrated manner in tissue regeneration and guided bone regeneration procedures due to its biocompatibility and ability to promote wound healing. The composition of collagen in the fabrication of BSA shows that this biomaterial has potential as a material used for dental applications.

SEM analysis

The SEM analysis results showed irregular pore of BSA samples with magnification was 50x and 100x (Figure 3A and 3B). The pore is a combination of macro and micro pore sizes. The results of this study showed the smallest pore was 133.101 μm and the largest was 605.803 μm with an average pore of 304.866 μm . The porosity percentage in BSA showed 64% porosity (Figure 3C). Pore structure in biomaterial is needed and important in wound healing process where gas permeability, absorption of wound exudates, cell development, exchange of nutrients, and metabolites will occur inside the pores.¹⁸ The microstructure, pore size, and porosity percentage of a biomaterial can be analyzed using SEM. The combination of macro and micro pores sizes in this study can be caused or influenced by the fabrication method. Freeze-drying method creates gelatin sponges with large and clearly visible pores.²³




Figure 3. SEM overview of the morphology and microstructure of bovine sponge amnion. (A) 50x magnification, (B) 100x magnification; (C) Image of 3D color fill surface overview of bovine sponge amnion porosity percentage.

Image-J is a software for analyzing digital images that has been widely used in medicine and biology.²⁴ In this study, the magnification was at 50x SEM image to calculate pore size. These results are in line with the study results of Murphy *et al.* (2010) that the ideal mean size of the pores for cell proliferation and migration in bone tissue formation is 325 μm . Minimum pore size for bone growth is in the optimal range of 100–135 μm . The pore size exceeding 300 μm will play a role in the process of vascularization.²⁵ Combined pores larger than 300 μm lead to direct osteogenesis while pores smaller than 300 μm can promote osteochondral ossification. As a scaffold, the pores on sponges have an oval structure with an ideal diameter ranging from 100 to 500 μm .²⁶

To allow cell growth to occur, the scaffold must have a porous structure with a porosity value of 80-90% to support regeneration. The greater the percentage of scaffold porosity, the faster the cell proliferation and differentiation.²⁷ In this study, the percentage porosity analysis was less than 80%. This can be affected by the freeze temperature and the freeze-drying method used.²⁸ This porosity percentage value can be influenced by the location where SEM images were taken. The definition of the size of the porosity is limited to the pixel size of the SEM images. Image pixel size affects the smallest measurable scale value and causes large deviations in the mean of porosity percentage. Biomaterials used for biomedicine generally have soft pores that formed a three-dimensional network.²⁹ Analysis with the use of Micro-computed X-ray tomography (Micro-CT) can improve the accuracy of the calculation of the percentage of porosity in biomaterials with porous structures. This analysis can also provide three-dimensional characterization of complex multi-porous microarchitectures.³⁰ After all, SEM results in this study indicate that this material has a combination of macro and micro size pore structures that can act as a material for wound healing in dental treatment.

	Before Immersion (gr)	After Immersion (gr)	Swelling percentage (%)
Specimen 1	0.0019	0.0270	1321
Specimen 2	0.0017	0.0223	1211
Specimen 3	0.0017	0.0260	1429
Mean \pm SD			1320 \pm 108

Table 1. The percentage of swelling after immersion in PBS solution.

The swelling ability plays an important role in tissue healing process, where the transfer of nutrients and body fluids into the biomaterial will be maximized if a biomaterial has good swelling

capability.³¹ In this study, when the sample was put into the PBS solution, the sample immediately expanded and after ten minutes, the mean swelling percentage reached 1320%. This result is in line with that conducted by Angulo and Sobral (2016) who used a incorporation of gelatin to fabricate scaffolds for tissue engineering. Gelatin has the ability to increase the swelling ability of the biomaterial.³² BSA contains collagen which causes this biomaterial to have high hydrophilicity properties. Hydrophilicity is the ability of collagen to bind water through hydrogen bonds, so that collagen is more easily dissolved in water. If the hydrogen bonds meet water (H_2O), then there is an intermolecular attraction at the H atom which is attached to the very electronegative atoms (N, O, and F).³³ A study by Cheng *et al.* (2017) shows that collagen sponges show a higher water absorption rate this can be caused with the presence of a porous collagen sponge structure and water retention of the collagen fibers.³⁴

Research conducted by Guclu *et al* (2017) increases the swelling capacity of a biomaterial that can occur up to 1200% to 1900% as a candidate biomaterial for bone healing.³⁵⁻³⁷ This shows that swelling is also an important feature for developing suitable tissue engineering constructs for bone defect regeneration.²⁷ The study by Indrawati *et al.* (2019) shows that BSA increase the bone morphogenetic protein-2 (BMP-2), as a marker of osteogenesis from the first day to the seventh day after the application in post-extraction of rat teeth.¹² This swelling presentation has been able to make BSA as a material for wound healing not only of soft tissue but also of hard tissue.

Conclusions

Based on the results of this study, it shows the potential characteristics of BSA as a biomaterial that can improve wound healing in dental treatment. Further studies are recommended for in vivo animal studies to analyze the effect of biomaterials on wound healing and regeneration after tooth extraction.

Acknowledgments

The author would like to thank the Installation of Tissue Bank and Cells of RSUD Dr. Soetomo Surabaya for the bovine sponge amnion (BSA). The author also thanks all laboratories that have supported this study, especially the DMTCORE Laboratory of Trisakti University. Last but not least, the author would like to thank Ms. Agnes Victoria, S.K.G for the lovely illustrations in this paper.

3 Declaration of Interest

The authors declare that there are no conflicts of interest.

Ethical policy and institutional review board statement

Ethical clearance had been obtained from the Ethics Commission of the Faculty of Dental Medicine, Airlangga University, Surabaya (No. 293/KKEPK. FKG/XII/2016).

Swelling analysis

In this study, the 10-minute immersion showed the results of the swelling percentage of 1321%, 1211%, and 1423% which are shown in Table 1.

References

1. Riskesdas. Laporan Riskesdas 2018. Badan Penelitian dan Pengembangan Kesehatan. Kementerian Kesehatan Republik Indonesia. 2018;181-222
2. Walsh DP, Raftery RM, Chen G, Heise A, O'Brien FJ, Cryan SA. Rapid healing of a critical-sized bone defect using a collagen-hydroxyapatite scaffold to facilitate low dose, combinatorial growth factor delivery. *J Tissue Eng Regen Med*. 2019;13:1843-53.
3. Eming S, Paul M, Tomic-Canic M. Wound repair and regeneration: Mechanisms, signaling, and translation. *Sci Transl Med*. 2014;6:1-36.
4. Ballesteros ACV, Puella HRS, Lopez-Garcia JA, Bernal-Ballen A, Mosquera DLN, Forero DMM, et al. Bovine decellularized amniotic membrane: Extracellular matrix as scaffold for mammalian skin. *Polymers (Basel)*. 2020;12.
5. Pei Y, Ye D, Zhao Q, Wang X, Zhang C, Huang W, et al. Effectively promoting wound healing with cellulose/gelatin sponges constructed directly from a cellulose solution. *J Mater Chem B*. 2015;3:7518-28.
6. O'Brien FJ. Biomaterials & scaffolds for tissue engineering. *Mater Today*. 2011;14:88-95.
7. Koizumi N, Inatomi T, Sotozono C, Fullwood NJ, Quantock AJ, Kinoshita S. Growth factor mRNA and protein in preserved human amniotic membrane. *Curr Eye Res*. 2000;20:173-7.
8. Niknejad H, Peirovi H, Jorjani M, Ahmadiani A, Ghanavi J, Seifalian M, et al. Properties Of The Amniotic Membrane For Potential Use In Tissue Engineering. *Eur Cells Mater*. 2008;88-99.
9. Putra NHD, Matulutan F, Wibowo MD, Danardono E. Effects of dried bovine amniotic membrane as prosthetics of abdominal fascial defect closure observed by the expression of platelet-derived growth factor in Rattus norvegicus wistar strain. *Syst Rev Pharm*. 2020;11:987-91.
10. Yang Z, Cui Q, An R, Wang J, Song X, Shen Y, et al. Comparison of microbiomes in ulcerative and normal mucosa of recurrent aphthous stomatitis (RAS)-affected patients. *BMC Oral Health*. 2020;20:1-8.
11. Cai M, Zhang J, Guan L, Zhao M. Novel implantable composite biomaterial by fibrin glue and amniotic membrane for ocular surface reconstruction. *J Mater Sci Mater Med*. 2015;26.
12. Indrawati DW, Munadzir E, Sulistyawati TIB, El Fadhlallah PM. Sponge amnion potential in post tooth extraction wound healing by interleukin-6 and bone morphogenetic protein-2 expression analysis: An animal study. *Dent Res J (Isfahan)*. 2019;16:283-8.
13. Dimitriou R, Mataliotakis GI, Calori GM, Giannoudis P V. The role of barrier membranes for guided bone regeneration and restoration of large bone defects: Current experimental and clinical evidence. *BMC Med*. 2012;10:1-24.
14. Dadkhah Tehrani F, Firouzeh A, Shabani I, Shabani A. A Review on Modifications of Amniotic Membrane for Biomedical Applications. *Front Bioeng Biotechnol*. 2021;8:1-25.
15. Zhang Q, Qian C, Xiao W, Zhu H, Guo J, Ge Z, et al. Development of a visible light, cross-linked GelMA hydrogel containing decellularized human amniotic particles as a soft tissue replacement for oral mucosa repair. *RSC Adv*. 2019;9:18344-52.
16. Kang M, Cho S, Cho Lee AR. Effect of freeze dried bovine amniotic membrane extract on full thickness wound healing. *Arch Pharm Res*. 2013;36:472-8.
17. Cheng H, Li C, Jiang Y, Wang B, Wang F, Mao Z, et al. Facile preparation of poly saccharide-based sponges and its potential application in wound dressing. *J Mater Chem B*. 2013;1-8.
18. Long H, Ma K, Xiao Z, Ren X. Preparation and characteristics of gelatin sponges crosslinked by microbial transglutaminase. *PeerJ*. 2017;1-18.
19. Khalid P, Suman VB, Shameema S, Vinodini NA, Shela RP, Chatterjee PK. Analytical methods for nanomaterial characterization. *Int J Nano Dimens Anal methods Nanomater Charact*. 2014;5:309-20.
20. Octarina, Munadzir E, Razak FA. Physical Modification of Bovine Amniotic Membrane for Dental Application. *Journal of International Dental and Medical Research*. 2021;14(4):1425-8.
21. Steiner GG, Francis W, Burrell R, Kallet MP, Steiner DM, Macias R. The healing socket and socket regeneration. *Compend Contin Educ Dent*. 2008;29:1-14.
22. Maryam S, Effendi N, Kasmah K. Produksi dan Karakterisasi Gelatin dari Limbah Tulang Ayam dengan Menggunakan Spektrofotometer Ftir (Fourier Transform Infra Red). *Maj Farm*. 2019;15:96.
23. Muyonga JH, Cole CGB, Duodu KG. Fourier transform infrared (FTIR) spectroscopic study of acid soluble collagen and gelatin from skins and bones of young and adult Nile perch (*Lates niloticus*). *Food Chem*. 2004;86:325-32.
24. Kabiri M, Emami SH, Rafinia M, Tahriri M. Preparation and characterization of absorbable hemostat crosslinked gelatin sponges for surgical applications. *Curr Appl Phys*. 2011;11:457-61.
25. uarez-Armedo A, Figueroa FT, Clavijo C, Arbeláez P, Cruz JC, Muñoz-Camargo C. An image J plugin for the high throughput image analysis of in vitro scratch wound healing assays Alejandra. *bioRxiv*. 2020:1-21.
26. Murphy CM, Haugh MG, Brien FJO. Biomaterials The effect of mean pore size on cell attachment , proliferation and migration in collagen – glycosaminoglycan scaffolds for bone tissue engineering. *Biomaterials*. 2010;31461-6.
27. León-Mandilla BH, Araiza-Téllez MA, Flores-Flores JO, Piña-Barba MC. Physico-chemical characterization of collagen scaffolds for tissue engineering. *J Appl Res Technol*. 2016;14:77-85.
28. Tohamy KM, Mabrouk M, Soliman IE, Beherei HH, Aboelnasr MA. Novel alginate/hydroxyethyl cellulose/hydroxyapatite composite scaffold for bone regeneration: In vitro cell viability and proliferation of human mesenchymal stem cells. *Int J Biol Macromol*. 2018;112:448-60.
29. Oikonomopoulou VP, Krokida MK, Karathanos VT. The influence of freeze drying conditions on microstructural changes of food products. *Procedia Food Sci*. 2011;1:647-54.
30. Ahumada M, Jacques E, Calderon C, Martinez-Gómez F. Porosity in biomaterials: A key factor in the development of applied materials in biomedicine. *Handb Ecomater*. 2019;5:3503-22.
31. Nair M, Shepherd JH, Best SM, Cameron RE. MicroCT analysis of connectivity in porous structures: Optimizing data acquisition and analytical methods in the context of tissue engineering. *J R Soc Interface*. 2020;17:1-16.
32. Maji K, Dasgupta S, Pramanik K, Bissoyi A. Preparation and Evaluation of Gelatin-Chitosan-Nanobioglass 3D Porous Scaffold for Bone Tissue Engineering. *Int J Biomater*. 2016;2016:1-14.
33. López Angulo DE, do Amaral Sobral PJ. Characterization of gelatin/chitosan scaffold blended with aloe vera and snail mucus for biomedical purpose. *Int J Biol Macromol*. 2016;92:645-53.
34. Peng L, Xiang RC, Jia WW, Dong XX, Wang GE. Preparation and evaluation of porous chitosan/collagen scaffolds for periodontal tissue engineering. *J Bioact Compat Polym*. 2006;21:207-20.
35. Cheng X, Shao Z, Li C, Yu L, Raja MA, Liu C. Isolation, characterization and evaluation of collagen from jellyfish *Rhopilema esculentum* kishinouye for use in hemostatic applications. *PLoS One*. 2017;12:1-21.
36. Guclu ZA, Hurt AP, Ohia L, Coleman NJ. Composite Sponges for in Situ Alveolar Bone Regeneration Following Tooth Extraction. *Acta Phys Pol A*. 2017;131:580-3.
37. Munadzir E, Soekartono RH, Rafli RBV, Yulianti A, Kartikasari N. Effect of rSLPI Amnion Membrane Application on Incision Wound of Rattus Norvegicus in Collagen and VEGF Expression. *Journal of International Dental and Medical Research*. 2017;10(2):239-43.

Characterization of Bovine Sponge Amnion (BSA) by a Novel Process for Dental Treatment

ORIGINALITY REPORT

17%	16%	10%	10%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universitas Airlangga Student Paper	3%
2	www.mdpi.com Internet Source	2%
3	repository.unair.ac.id Internet Source	2%
4	journal.unpad.ac.id Internet Source	1%
5	Submitted to Pennsylvania College of Technology Student Paper	1%
6	www.ukm.my Internet Source	1%
7	Submitted to University of Auckland Student Paper	1%
8	worldwidescience.org Internet Source	1%

e-journal.unair.ac.id

9	Internet Source	1 %
10	Submitted to Universitas Jember Student Paper	1 %
11	Submitted to Cardiff University Student Paper	1 %
12	Zhang, Yaya. "A Design, Development, and Evaluation of Bio-Patch for Myocardial Tissue Repair", The University of Tennessee Health Science Center, 2023 Publication	1 %
13	hrcak.srce.hr Internet Source	1 %
14	"Novel Biomaterials for Regenerative Medicine", Springer Science and Business Media LLC, 2018 Publication	1 %
15	hdl.handle.net Internet Source	<1 %
16	www.artificial-organs.com Internet Source	<1 %
17	aip.scitation.org Internet Source	<1 %
18	www.sysrevpharm.org Internet Source	<1 %

Exclude quotes On

Exclude matches

< 15 words

Exclude bibliography On

Characterization of Bovine Sponge Amnion (BSA) by a Novel Process for Dental Treatment

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6