



SCOG

SCIENTIFIC CONTRIBUTION OIL AND GAS

JOURNAL

ISSN : 2089-3361

e-ISSN : 2541-0520



VOLUME 47 NO. 1, APRIL 2024

ACCREDITED RISTEK DIKTI
79/E/KPT/2023



HOME ABOUT LOGIN SEARCH CURRENT ARCHIVES CONTACTS JOURNAL LEMIGAS

Home > About the Journal > **Editorial Team**

Editorial Team

Ensured Editor

Ir Mustafid Gunawan, M.E., Testing Center for Oil and Gas LEMIGAS

Chief Editor

Prof. Dr. Bambang Widarsono, National Research and Innovation Agency, Indonesia

Chief Deputy Editor

Basuki Basuki, Testing Center for Oil and Gas LEMIGAS, Indonesia
Dr. Setyo Widodo, Testing Center for Oil and Gas LEMIGAS, Indonesia

Managing Editor

Purwo Subagiyo, Testing Center for Oil and Gas LEMIGAS, Indonesia
Nurhadi Setiawan, Testing Center for Oil and Gas LEMIGAS, Indonesia

Associate Editor

Dr. Junita Trivianty, Testing Center for Oil and Gas LEMIGAS, Indonesia
Dr. Humbang Purba, Testing Center for Oil and Gas LEMIGAS, Indonesia
Dr. Tri Muji Susantoro, National Research and Innovation Agency, Indonesia
Dr. Cut Nanda, Testing Center for Oil and Gas LEMIGAS, Indonesia
Dr. Rachmi Kartini, Testing Center for Oil and Gas LEMIGAS, Indonesia
Dr. Christina Ani Setyaningsing, Testing Center for Oil and Gas LEMIGAS, Indonesia
Dr. Riesta Anggarani, Testing Center for Oil and Gas LEMIGAS, Indonesia
Dr. Usman Usman, National Research and Innovation Agency, Indonesia
Himawanto Himawanto, Testing Center for Oil and Gas LEMIGAS, Indonesia

Editorial Board

Peer Reviewer

Peer Review Process/Policy

Aim & Scope

Copyright Policy/ Author Guidelines

Online Submission

Presence of Substance

Publication Ethics

USER

Username

Password

☐ Remember me

Login

AUTHOR GUIDELINE



International Editorial Boards

Prof. Dr. Bambang Widarsono, National Research and Innovation Agency, Indonesia
Prof. Dr. Istadi M.T., University of Diponegoro, Indonesia
Prof. Dr. Maizar Rahman, Testing Center for Oil and Gas LEMIGAS, Indonesia
Prof. Dr. M. Kholil, Bogor Agricultural University/USAHID, Indonesia
Prof. Septoratto Siregar, Ph.D, Bandung Institute of Technology, Indonesia
Prof. M. Udiarto, Testing Center for Oil and Gas LEMIGAS, Indonesia
Prof. Dr. R. P. Koesoemadinata, Bandung Institute of Technology, Indonesia
Prof Pudji Permadi, Ph.D, Bandung Institute of Technology, Indonesia
Prof. Renanto Ph.D., Technology Institute of Surabaya, Indonesia
Dr. Anda Lucia, Testing Center for Oil and Gas LEMIGAS, Indonesia
Dr. Ulrike Schacht, Geo Marine Chemist, University of Adelaide, Australia
Dr. Robert John Morley, Palynologist and Stratigrapher, Palynova, United Kingdom
Dr. Nur Suhascaryo, University of National Development "Veteran" Yogyakarta, Indonesia
Dr. Hyundon Shin, Inha University, Korea, Republic of
Dr. Suhendra Suhendra, Sunan Kalijaga State Islamic University, Indonesia
Dr. Usman Usman, National Research and Innovation Agency, Indonesia
Dr. Restu Juniah, University of Lampung, Indonesia
Dr. Dedy Kristanto, University of National Development "Veteran" Yogyakarta, Indonesia
Dr. Eko Budi Lelono, Palynologist, Agency for Geology, Indonesia
Dr. Mudjito Mudjito, Testing Center for Oil and Gas LEMIGAS, Indonesia
Dr. Adiwar Adiwar, Testing Center for Oil and Gas LEMIGAS, Indonesia
Dr Oberlin Sidjabat, Testing Center for Oil and Gas LEMIGAS, Indonesia
Valentina Alvarez, Ph. D., Centro de Investigación en Ecosistemas de la Patagonia (CIEP), Chile
Ryozo Noguchi, Ph. D., University of Kyoto, Japan
Bagas Pudji Laksono, Ph.D, Gadjah Mada University, Indonesia
Jonathan Setyoko Hadimuljono, Testing Center for Oil and Gas LEMIGAS, Indonesia
John G Kaldi, Ph.D, University of Adelaide, Australia
Dr. Nuning Vita Hidayati, University of Soedirman, Indonesia
Dr. Noor Cahyo D. Aryanto, Exploration Geology and Petrology Geochemistry, Indonesia

Language Editor

Ferry Imanuddin Sadikin, Testing Center for Oil and Gas LEMIGAS, Indonesia
Windy Permata Luvian, Testing Center for Oil and Gas LEMIGAS, Indonesia

Design Layout and OJS

Andry Hidayat, Testing Center for Oil and Gas LEMIGAS, Indonesia

Circulation

Wulandari Dianningtyas, Testing Center for Oil and Gas LEMIGAS, Indonesia, Indonesia

OJS Website Administrator

Yasdi Pramesti, Testing Center for Oil and Gas LEMIGAS, Indonesia

Photographer

TEMPLATE

ANUS
TEMP

TOOLS



JOURNAL CONTENT

Search

Search Scope

All

Search

- Browse
- By Issue
 - By Author
 - By Title
 - Other Journals

Journal Help

Visitor Counter:

Visits	23 974
Pages	237 172

Dicky Christian Fransiscus, Testing Center for Oil and Gas LEMIGAS, Indonesia



Copyright 2023 | Scientific Contributions Oil and Gas (SCOG). Powered By OJS

LEMIGAS



Indexed On :





HOME ABOUT LOGIN SEARCH CURRENT ARCHIVES CONTACTS JOURNAL LEMIGAS

Home > Archives > **Vol 47, No 2 (2024)**

Vol 47, No 2 (2024)

Table of Contents

Articles

- Bioprospecting of Halophilic Bacteria Staphylococcus Haemolyticus Strain Stp-Griv-024 as Biosurfactant Producer and its Potential Application for Microbial Enhanced Oil Recovery** 75-88

Izzul Islam, Cut Nanda Sari, Ali Budhi Kusuma, Ika Kurniasari, Lisnawati Lisnawati, Rini Surya Ningsih, Elisa Rosani

- Quantifying the Impact of Siderite Composition and Reservoir Resistivity (Rt) on Water Saturation Estimation in Low-Resistivity Sideritic Sandstone Reservoirs Using the Graphic Plot Method** 89-100

Sarju Winardi, Sugeng Sapto Surjono, Donatus Hendra Amijaya, Wiwit Suryanto

- Updated Geochemical Characterization of Hydrocarbon Potential of Surface Sample in Lariang-Karama Sub-Basin, West Sulawesi, Indonesia** 101-115

Yarra Sutadiwiria, Muhammad Taufiq Fathaddin, Imam Setiaji Ronoatmojo, Dewi Syavitri, Cahyaningratri P R, Barona Belladesta, Yeftamikhya Yeftamikhya, Andy Livsey

- Pectin Extraction of Jackfruit Peel as a Biopolymer Potential with Microwave Assisted Extraction Method** 117-128

Muhammad Khairul Afdhol, Fiki Hidayat, Tomi Erfando, Dita Putri Purnama

- Seismic Data Processing and Seismic Inversion in The Ray Parameter Domain: Common Reflection Point (CRP) Stack and Ray Impedance** 129-139

Wahyu Triyoso, Edycakra Immanuel Sinaga, Madaniya Oktariena

- The Influence of Side Slip Velocity on Subsurface Displacement** 141-150

Editorial Board

Peer Reviewer

Peer Review Process/Policy

Aim & Scope

Copyright Policy/ Author Guidelines

Online Submission

Presence of Substance

Publication Ethics

USER

Username

Password

☐ Remember me

Login

AUTHOR GUIDELINE




The Influence of Side-Slip Velocity on Subsurface Displacement

Imam Setiaji Ronoatmojo, Muhammad Burhannudinnur, Yarra Sutadiwiria, Dewi Syavitri

Modeling The Distribution of Wastewater and Determining Zone of Initial Dilution from Waste Discharge Into The Sea From Fuel Terminal Activities in Balikpapan Bay

Randi Andhika, Purwanto Purwanto, Fuad Muhammad



Copyright 2023 | Scientific Contributions Oil and Gas (SCOG). Powered By OJS



Indexed On :



TEMPLATE

ANUS
TEMP

TOOLS



JOURNAL CONTENT

Search

Search Scope

All

Search

- Browse
- By Issue
 - By Author
 - By Title
 - Other Journals

Journal Help

Visitor Counter:

Visits	23 953
Pages	236 738





Updated Geochemical Characterization of Hydrocarbon Potential of Surface Sample in Lariang-Karama Sub-Basin, West Sulawesi, Indonesia

Yarra Sutadiwiria¹, Muhammad Taufiq Fathaddin², Imam Setiaji Ronoatmojo¹, Dewi Syavitri¹,
Cahyaningratri, P.R.¹, Barona Belladesta¹, Yeftamikha³ and Andy Livsey⁴

¹Geology Department, Faculty of Earth Technology and Energy, Universitas Trisakti
Kyai Tapa Street No. 1, Jakarta 11440, Indonesia

²Master's Program of Petroleum Engineering, Faculty of Earth Technology and Energy, Universitas Trisakti
Kyai Tapa Street No. 1, Jakarta 11440, Indonesia

³Faculty of Engineering, University of Auckland, 20 Symonds Street, Auckland 1010, New Zealand
34 Princes Street, Auckland CBD, Auckland 1010, New Zealand

⁴PT Horizon Geoconsulting
Pondok Indah Office Tower 2, 15th Floor, Sultan Iskandar Muda Street Kav. VTA, Pondok Indah,
South Jakarta, 12310, Indonesia

Corresponding author: yarra.sutadiwiria@trisakti.ac.id

Manuscript received: June 25th, 2024; Revised: July 16th, 2024

Approved: July 19th, 2024; Available online: July 22th, 2024

ABSTRACT - The Lariang-Karama sub-basin is situated within the Makassar basin, located in the West Sulawesi region. It has source rocks which have the potential to produce hydrocarbons and to become a target for new field discoveries in hydrocarbon exploration activities. It is on this basis that the authors are interested in conducting research in the Makassar basin area and its surroundings, which has been carried out since 2017. This is a comprehensive literature review study from 2017 to 2022, focusing on the origin and composition of biomarkers, paleogeography of facies sources from oil seeps (2017-2018), paleofacies and biomarker characteristics of Paleogene to Neogene rocks (2019), contribution of terrestrial materials based on geochemical and biostratigraphic analysis (2020), tectonic processes occurring in the basin, and geochemical characterization of source rocks in the research area (2022). The FS-12 surface sample in shale may belong to the Kalumpang Formation, located in West Sulawesi. The results of TOC analysis were 0.62%, indicating sufficient quantity. Based on the results of the cross plot between HI values of < 50 and Tmax of 499 °C, the category was found to be overmature and could not produce hydrocarbons. The results of this research, along with those of previous research and the latest geochemical study, were obtained in the West Sulawesi region and the eastern Makassar Strait. These findings consisted of marine shale source rocks in West Sulawesi and lacustrine shale source rocks in the eastern Makassar Strait.

Keywords: source rock, West Sulawesi, total organic carbon (TOC).

© SCOG - 2024

How to cite this article:

Yarra Sutadiwiria, Muhammad Taufiq Fathaddin, Imam Setiaji Ronoatmojo, Dewi Syavitri, Cahyaningratri, P.R., Barona Belladesta, Yeftamikha and Andy Livsey, Scientific Contributions Oil and Gas, 47 (2) pp. 101-115. DOI.org/10.29017/SCOG.47.2.1624.

INTRODUCTION

Hydrocarbons play a very important role as an energy resource to support the Indonesian economy (Praptisih 2022). Exploration of hydrocarbons is a crucial activity in the oil and gas sector that involves searching for new reserves to meet production needs and developing existing oil fields. An essential component of exploration is geochemical analysis, which can be carried out on surface samples and hydrocarbon seeps. Various geochemical analyses have been conducted on surface samples and seeps in West Sulawesi. In research by (Sutadiwiria et al. 2017, 2018) on oil seeps in the Lariang and Karama areas of West Sulawesi, the oleanane saturation found in these seeps indicates that the source rocks in the West Sulawesi area were formed in delta and near-coastal environments. This oil seep is known to originate from Eocene coal and Toraja shale coal from the Kalumpang formation. The results of biomarker analysis on GC-MS show that the source of oil seep in Lariang can be traced back to terrestrial organic material, with the additional presence of organic materials from algae and herbs (Sutadiwiria 2019).

The Lariang-Karama sub-basin in the Makassar basin has the potential to produce hydrocarbons. This study aims to refine the predictions regarding the capacity of the area's source rock to produce hydrocarbons by using geochemical analyses of hydrocarbons. These analyses determine the quality of organic materials from rock-eval pyrolysis (REP), the quantity of organic materials from total organic carbon (TOC), the degree of maturity of the hydrocarbon from cross-plot Tmax vs HI, gas chromatography (GC), and gas chromatography/mass spectrometry (GC/MS).

Regional Geology

Makassar basin is located in Indonesia and is bounded by different geographical features. To the north lies the Mangkalihat High, while the Java Sea is situated in the south. In the west, it is bordered by the East Kalimantan mainland, which includes the Mahakam Delta and the SSW-NNE fold structure. On the east, it is bounded by the mainland of West Sulawesi, which has a thrust fold in the SSW-NNE direction. The Makassar basin was formed during the pre-Tertiary to late tertiary extensional stages, which occurred no earlier than the early paleocene (Sutadiwiria et al. 2022). Calvert & Hall (2003) have reported that the research area is located in the eastern part of the Makassar Basin. The basin

was formed during the early tertiary, paleocene, and middle Eocene-Late Eocene. This formation was due to its expansion. The rifting led to the formation of graben and half-graben, and the subsidence formation took place during the early period of the rifting. This process caused the island of Sulawesi to move from east to southeast and separate from the mainland of Sunda/Kalimantan. The area is subjected to extensional compressional forces, which have led to the upliftment, folding, and shearing of the pre-tertiary basement, resulting in the formation of rock depressions and elevations (Figure 1).

Regional Tectonic

The Makassar basin, where the Lariang-Karama Sub-Basin is formed, has three main fault patterns: northeast-southwest, northwest-southeast, and north-south. According to (Fraser 2003, Calvert and Hall 2007, and Raharjo et al. 2012), the formation of Makassar basin can be divided into six tectonic stages (Figure 2). These stages are the Pre-rift stage (pre- Tertiary), Syn-rift package stage (Middle-Late Eocene), Post-rift package stage (Oligocene), Post-rift package stage (Early-Middle Miocene), Post-rift package stage (Late Miocene) and Foreland basin package stage (Early-recent Pliocene).

Regional Stratigraphy

The study area is marked on a stratigraphic column (Calvert & Hall 2003; Raharjo et al. 2012) (Figure 2). The area is part of the Lariang-Karama Sub-Basin, which is situated in the eastern region of Makassar Basin. The geological formations in the area are arranged in the following order from the oldest to the youngest. Basement (Mesozoic) is composed of metamorphic rocks, black shale, and volcanic rocks. Kalumpang Formation (Middle Eocene-Late Eocene) consists of claystone, carbonate shale, coal seams, siltstone, and quartz sandstone. Budung-budung formation (Middle Eocene-Early Miocene) consists of shale, shaly mudstone, coal, siltstone, clastic limestone, and conglomerate. The lisu formation consists of early-late miocene mudstone, sandstone, and layers of volcanic product debris and tuff. The Pasang kayu formation (early pliocene pleistocene) consists mainly of conglomerate and layers of sandstone and mudstone. Based on regional geology, (Calvert & Hall 2007 and Raharjo et al. 2012) identified the source rocks in West Sulawesi into two potential formations: the eocene Kalumpang formation and the miocene Budung-Budung formation. According




Figure 1
The tectonic setting that influences the Makassar Basin (Fraser 2003)

to (Waples 1985), source rocks can be classified into three groups: effective source rock, possible source rock, and potential source rock. Effective source rock is formed by organic materials which can accumulate and generate hydrocarbons. Possible source rock has the potential to store and produce hydrocarbons, but it has not been evaluated. On the other hand, potential source rock is immature but can store and produce hydrocarbons as it matures.

METHODOLOGY

This research involved the use of geochemical analysis, including total organic carbon (TOC) measurement, rock-eval pyrolysis analysis, cross-plotting Tmax vs HI and HI vs OI for hydrocarbon maturity assessment, gas chromatography (GC), and gas chromatography/mass spectrometry (GC/MS). Additionally, Microsoft Excel software was used to generate graphs. The geochemical analysis was carried out in the geochemical laboratory of PT.

Geoservices, and the standard laboratory methods have been internationally recognized according to ISO 900:2015. The results of this study were combined with the findings of previous studies to provide an updated and comprehensive geochemical analysis for the West Sulawesi region.

Total Organic Carbon (TOC)

TOC is the quantity of organic carbon deposited in the rock which is expressed as a percentage of the weight of the dry rock. Organic carbon is carbon that comes from organic substances and does not come from carbonates, for example limestone. The total organic carbon (TOC) value is determined by analyzing the sample data and obtaining numerical results. Sedimentary rocks that contain a TOC value of 0.5% or lower are considered to have a very low potential to produce hydrocarbons. On the other hand, rocks that have the potential to be a source rock contain a TOC of >0.5% as per (Peters 1994) Table 1.

Table 1
Source rock quality parameters based on TOC (Peters 1994)

Quality	TOC (w.t%)
Bad	< 0.5
Fair	0.5 – 1
Good	1 – 2
Very Good	2 – 4
Excellent	> 4




Figure 2
Modified Mamuju Regional Stratigraphic Column, West Sulawesi (Calvert & Hall 2003 and Raharjo et al. 2012)

Kerogen Type

Kerogen is the organic material present in source rocks that produce oil and gas. It plays a significant role in determining which hydrocarbons will be formed. Kerogens can be classified into different types based on the combination of chemical elements that it contains, such as carbon (C), hydrogen (H),

oxygen (O), nitrogen (N), and sulfur (S). The method used to classify kerogen types is the use of the Van Krevelen diagram (Peters & Moldowan 1993), which involves plotting the hydrogen index (HI) against the oxygen index (OI) atoms. Additionally, a comparison is made between HI and Tmax values.

Rock–Eval Pyrolysis (REP)

The parameters used to determine the type of kerogen, oil, and gas products are formed at peak maturity (Peters & Cassa 1994). The pyrolysis parameters used to determine the type of hydrogen are the hydrogen index (HI) and the relationship between the ratio of S2 (a hydrocarbon formed from kerogen in rock-eval pyrolysis due to the decomposition of kerogen) and S3. Figure 3 illustrates an analysis of rock-eval pyrolysis in the study area based on (Soppanata 1996).

According to (Hunt 1996), Rock-Eval pyrolysis analysis is based on several parameters. S1 (free hydrocarbons) indicates the number of free hydrocarbons that can be vaporized with out kerogen degradation. It is formed at 300 °C, resulting from kerogen during deposition. S2 (pyrolisable hydrocarbon) indicates the number of hydrocarbons produced through the cracking process. It is formed at a temperature of 350–550°C. S3 shows the amount of CO₂ content in the rock. It is formed at 300–390 °C. Tmax shows the maximum temperature at the time of the formation of hydrocarbon cracking (S2). It is formed at 435–470 °C.

Maturity of Organic Material

The maturity of organic materials was assessed using two parameters: Tmax and Vitrinite Reflectance(Ro). The tmax value indicates the temperature at which hydrocarbons are formed through cracking (S2) and reach their maximum

intensity. Generally, this occurs at 435-470 °C, known as the oil window. On the otherhand, Ro is a parameter that indicates the increase in temperature and its effect on the reflectance of vitrinite. As the temperature rises, the reflectance gloss of vitrinite maceral willalsoincrease. Table 2 provides more information on Ro and its relationship with temperatures.

Gas chromatography (GC) and/or gas chromatography mass spectrometry (GC-MS) analyses were also used in this study. Saturate samples were subjected to gas chromatography (GC) to determine the distribution of isoprenoids and n-alkanes. In the meantime, compounds with low concentrations that could not be separated by gas chromatography (GC) were separated using gas chromatography mass spectrometry (GC-MS) (Waples & Curiale 1999).

The study utilized new data obtained from a surface sample called FS-12, which was taken from the Mamuju area in the Lariang-Karama Sub-Basin, West Sulawesi by (Sutadiwiria 2019). The availability of the data is presented inTable3. The lithostratigraphic analysis of the study area revealed that the samples were from the Middle Eocene Kalumpang Formation, which has an upper intertidal environment and comprises shale, as shown in Figure 4.

Table2
Maturity parameters of organic materials (Peters & Cassa1994)

Parameter	Maturity			Generation		
	Ro (%)	Tmax	TAI	Bit/TOC	Bitumen	PI
Immature	0.20 – 0.60	< 435	1.5 – 1.26	< 0.05	< 50	< 0.10
Early Mature	0.60 – 0.65	435 – 445	2.6 – 2.7	0.05 – 0.10	50 – 100	0.10 – 0.15
Peak Mature	0.65 – 0.90	445 – 450	2.7 – 2.9	0.15 – 0.25	150 – 250	0.25 – 0.40
Late Mature	0.90 – 1.35	450 – 470	2.9 – 3.3	-	-	> 0.40
Post Mature	> 1.35	> 470	> 3.3	-	-	-




Figure 3
Methodology of rock-eval pyrolysis analysis (Soppanata 1996).

Table 3
Availability of data used in the research

	Sample Identification	Lithology	TOC	S1	S2	S3	Tmax	HI	OI	PI	VR (%)	Original References
Surface sample	Name											
	FS-12	Shale	0.62	0.08	0.20	0.12	499	32	19	0.29	1.3	new data
	FS-2	Coal	30.56	0.61	33.29	21.01	423	109	69	0.02	0,46	Sutadiwiria et al. (2022)
	FS-20	Carbonaceous shale	6,92	0.28	11.94	2.58	429	189	37	0,02	0,38	Sutadiwiria et al. (2022)




Figure 4
Location map and geology of Mamuju and surrounding areas modification of (Calvert & Hall 2003 and Raharjo et al.2012)

RESULT AND DISCUSSION

The analysis of the maturity level and type of hydrocarbon kerogen led to the conclusion that the FS-12 sample from the Mamuju area, West Sulawesi, belonged to the kerogen type IV (Figures 5 and 6). The hydrogen index (HI) value of the sample was found to be 32 (mg/grrock/% TOC, table, Figures 5 and 6), which indicates that the kerogen has a lower potential to produce hydrocarbons as the HI value is less than the required 50 mg/gr rock (Waples 1985). More over, the t_{max} value of FS-12 was 499°C,

which falls into the post-mature category based on (Peters & Cassa 1994), Figure 5 Previous studies on the surface samples (FS-2 and FS-20) showed high TOC values (30.56% and 6.32%, respectively) and HI values of 109 mg/C and 189 mg/C. Based on the HI value, the FS-2 and FS-20 samples were likely to have the potential to be gas-prone for FS-2 (type III kerogen), and to be oil and gas-prone for FS-20 (Type II kerogen) (modification of (Sutadiwiria et al. 2022) as shown in Table 3 and Figures 5 and 6.

T_{MAX} VS HI WEST SULAWESI




Figure 5
Results of crossplot diagram of hydrogen index (HI) vs T_{max} of Mamuju West Sulawesi (Van Krevelen 1993 & Waples 1985)





Figure 6
Results of crossplot diagram of van krevelen hydrogen index (HI) vs Oxygen Index (OI) Mamuju West Sulawesi (Van Krevelen 1993)

A number of terrestrial components (resin compounds), including the biomarkers oleanane (OL), bicadinane, and gamaserane (Gm), were visible in the mass fragmentogram profile, particularly the m/z 191 ion of the saturated hydrocarbon fraction of GCMS FS-12 (Figure 7). The same biomarkers were also present in the samples FS-2 and FS-20 (Sutadiwiria et al. 2022) (Figures 8 and 9). The components W, T, and R originating from higher plants were identified as the bicadinane. These components were components a, b, c, and

d. Ion taraxastana (Tx) also appeared in several samples along with the presence of oleanane. The ternary diagram of samples FS-2, FS-12, and FS-20 (Figure 10) from Peters et al. (2005) and Huang and Meinschein (1979) indicates a relative abundance plot of sterane (C_{27} - C_{29}), which shows a relatively high proportion of C_{29} sterane. The percentages are 48.29%, 33.12%, and 55.27%. In comparison C_{27} shows Figures of 26.5%, 36.69% and 10.73%, while C_{28} has percentages of 25.21%, 30.19% and 34%, respectively.


Abundance



Time-->

Figure 7

m/z 191 partial mass chromatograms showing triterpane distribution of FS-12



Time-->

Figure 8

m/z 191 partial mass chromatograms showing triterpane distribution of FS-2. (Sutadiwiria et al. 2022)

Abundance

Ion 191.00 (190.70 to 191.70): YARRA 12 SAT.Ddatams




Figure 9
m/z 191 partial mass chromatograms showing triterpane distribution of FS-20 (Sutadiwiria et al. 2022)




Figure 10
Saturated fraction of the surface sample extract plotted on the ternary diagram according to (a) (Huang & Meinschein 1979) and (b) (Peters et al. 2005)

The mass fragmentogram ion m/z 217 GCMS profile from samples FS-12, FS-02, and FS-20 further corroborates the ternary diagram (Figures 11, 12 and 13). This suggests that the three samples contain both terrestrial and marine organic materials. According to this update, organic materials derived from terrestrial and marine organic materials are deposited in the upper intertidal-fluviodeltaic environment as coal (FS-2), shale (FS-12), and carbonaceous shale (FS-20) (Sutadiwiria et al. 2022). This finding also accords with the study (Sutadiwiria et al. 2017) that found a significant amount of resin biomarkers in the source rock, indicating input from the fluviodeltaic environment.

Upon examination of the findings of the biostratigraphic, petrographic, and radiometric analyses of the current well K-1 in the eastern Makassar strait (Harsanti et al. 2013; Sutadiwiria et al. 2022), it becomes evident that the Late Cretaceous volcanic materials are located at a depth of 17,340–17,360 feet, while the shales and shale mudstones from the early paleocene and eocene are marginally lacustrine paleofacies (Geoservices 2012). The finding of lacustrine shale source rock in the eastern Makassar Strait was made possible by the comparatively higher concentration of C27 sterane in the well K-1 sample, when compared to C28 and C29 (Sutadiwiria et al. 2022).




Figure 11
m/z 217 partial mass chromatograms showing sterane distribution of FS-12

Abundance




Figure 12
m/z 217 partial mass chromatograms showing sterane distribution of FS-2 (Sutadiwiria et al. 2022)




Figure 13
Modified m/z 217 partial mass chromatograms showing sterane distribution of FS-20 (Sutadiwiria et al. 2022)

CONCLUSION

After analyzing the hydrocarbons in the Lariang-Karama Sub-Basin, Mamuju, and West Sulawesi, it was found that the FS-12 sample from the Kalumpang formation had fair TOC content, but it was not a source rock in the study area due to its post-mature maturity level. The results indicate that it belongs to the Middle Eocene shale lithology and has no potential to become a source rock (Type IV), unlike FS-20 and FS-2, which displays potential for oil and gas generation (Types II and III kerogen, respectively). Following a thorough assessment of post-drilling laboratory study, the West Sulawesi area continues to hold potential to be explored, offering improved prospects and reduced risks for exploration activities in the region.

ACKNOWLEDGMENT

The authors would like to express their gratitude for the invaluable support provided by the Department of Geological Engineering, Faculty of Earth and Energy Technology, Universitas Trisakti, in carrying out and presenting this research.

GLOSSARY OF TERMS

Symbol	Definition	Unit
Tmax	Maximum Temperature	
TOC	Total Organic Carbon	
w.t%	Weight percent (of TOC)	
REP	Rock Eval Pyrolysis	
GC	Gass Chromatography	
GC-MS	Gass Chromatography and Mass Spectroscopy	
HI	Hydrogen Index	
OI	Oxygen Index	
PI	Production Index	
VR	Vitrinite Reflectance (Ro)	
S1	Free hydrocarbons	
S2	Pyrolyzable hydrocarbon	
S3	Amount of CO ₂ content present	

Symbol	Definition	Unit
HI = mg/gr rock/% TOC	the amount of hydrogen relative to the amount of organic carbon present in a sample	
C	Carbon	
H	Hydrogen	
O	Oxygen	
N	Nitrogen	
S	Sulfur	
°C	Degrees Celsius	

REFERENCES

- Calvert, S.J. & Hall, R., 2007, Cenozoic Evolution of the Lariang and Karama Regions, North Makassar Basin, Western Sulawesi, Indonesia. *Petroleum Geoscience*, 13, 353–368. doi.org/10.1144/1354-079306-757.
- Calvert, S.J. & Hall, R., 2003, The Cenozoic Geology of The Lariang and Karama Regions, Western Sulawesi: New Insight into The Evolution of The Makassar Straits Region.
- Fraser, T.H., Jackson, B.A., Barber, P.M., Baillie, P., & Myers, K., 2003, The West Sulawesi fold belt and other new plays within the North Makassar Strait—a prospectivity review. *Proceedings Indonesian Petroleum Association*, 29th Annual Convention, p. 429-450.
- Hunt, J.M., 1996, *Petroleum Geochemistry and Geology* (2nd ed.). W.H. Freeman and Company.
- Peters, K.E. & Cassa, M.R., 1994, *Applied Source Rock Geochemistry. The Petroleum System from Source to Trap*. The American Association of Petroleum Geologists Memoir, 60.
- Peters, K.E. & Moldowan, J.M., 1993, *The Biomarker Guide, Interpretating Molecular Fossils in Petroleum and Ancient Sediments*. Prentice Hall, Englewood Cliffs, New Jersey 07632.
- Praptisih 2022, *Karakteristik Geokimia Organik pada Formasi Cibulakan di Daerah Cirebon, Jawa Barat*. Lembaran Publikasi Minyak dan Gas

Bumi, Vol. 56 No. 2, 123 – 135. doi.org/10.29017/LPMGB.56.2.1176

Raharjo, S., Seago, R., Jatmiko, E. W., Hakim, F. B. & Meckel, L.D., 2012, Basin Evolution and Hydrocarbon Geochemistry of The Lariang Karama Basin: Implications for Petroleum System in on Shore West Sulawesi. Proceedings of the Indonesian Petroleum Association, 36th Annual Convention and Exhibition.

Soppanata, M.A., 1996, Buku Praktikum Geokimia Hidrokarbon. Universitas Trisakti. Buku Praktikum.

Sutadiwiria, Y., Burhannudinnur, M., Jambak, M. A., Cahyaningratri, P. R., & Yeftamikha 2022, Geochemical Characterization of Coal, Carbonaceous Shale, and Marine Shale as Source Rock in West Sulawesi, Indonesia. Indonesian Journal on Geoscience, 9(3). <https://doi.org/10.17014/ijog.9.3.303-314>

Sutadiwiria, Y., 2019, Peranan Biomarker dan Paleofasies pada Rekonstruksi Tektonik Batuan Paleogen – Neogen di Selat Makassar Bagian Timur dan Sulawesi Barat. PhD Thesis, Universitas Padjadjaran (unpublished).

Sutadiwiria, Y., Hamdani, A.H., Sendjaja, Y.A., Haryanto, I. & Yeftamikha 2018, The Biomarker Composition of Some Oil Seeps from West Sulawesi, Indonesia. Indonesian Journal on Geoscience, 5, No. 3, 211-220. doi.org/10.17014/ijog.5.3.211-220.

Sutadiwiria, Y., Hamdani, A.H., Andriana, Y., Haryanto, I. & Sunardi, E., 2017, Origin of Oil Seeps in West Sulawesi Onshore, Indonesia: Geochemical Constraints and Paleogeographic Reconstruction of the Source Facies. In Journal of Geological Sciences and Applied Geology (Vol. 2, Issue 1).

Van Krevelen, D.W., 1993, Coal: Typology, Physics, Chemistry, Constitution. 979 pp., 3rd edition, Elsevier, Amsterdam, the Netherlands.

Waples, D.W., 1985, Geochemistry in Petroleum Exploration. International Human Resources Development Corporation, Boston.

Paper SCOG 2024-1

by Yarra Sutadiwiria FTKE

Submission date: 23-Aug-2024 09:41AM (UTC+0700)

Submission ID: 2433024349

File name: Paper_SCOG_2024-1.pdf (1.29M)

Word count: 3820

Character count: 20863

**UPDATED GEOCHEMICAL CHARACTERIZATION OF HYDROCARBON
POTENTIAL OF SURFACE SAMPLE IN LARIANG-KARAMA
SUB-BASIN, WEST SULAWESI, INDONESIA**

Yarra Sutadiwiria^{1*}, Muhammad Taufiq Fathaddin², Imam Setiaji Ronoatmojo¹,
Dewi Syavitri¹, Cahyaningratri, P.R.¹, Barona Belladesta¹, Yeftamikha³, and Andy Livsey⁴

¹Geology Department, Faculty of Earth Technology and Energy, Universitas Trisakti,
Jln. Kyai Tapa No. 1, Jakarta 11440, Indonesia

²Master's Program of Petroleum Engineering,
Faculty of Earth Technology and Energy, Universitas Trisakti,
Jln. Kyai Tapa No. 1, Jakarta 11440, Indonesia

³Faculty of Engineering, University of Auckland,
20 Symonds Street, Auckland 1010, New Zealand.

⁴PT Horizon Geo Consulting, Pondok Indah Office Tower 2, 15th Floor, Jln. Sultan Iskandar
Muda Kav. VTA, Pondok Indah, Jakarta Selatan 12310

*corresponding author: yarra.sutadiwiria@trisakti.ac.id

Abstrak

Lariang-Karama merupakan salah satu subcekungan di Cekungan Makassar yang terletak di wilayah Sulawesi Barat. Diharapkan dapat menghasilkan hidrokarbon dan berpotensi menjadi sasaran penemuan lapangan baru dalam kegiatan eksplorasi hidrokarbon, selain untuk memperkirakan adanya potensi batuan induk untuk menghasilkan hidrokarbon. Atas dasar inilah maka penulis tertarik untuk melakukan studi dan penelitian di daerah Cekungan Makassar dan sekitarnya, yang dilakukan sejak tahun 2017 hingga sekarang. Kajian ini merupakan studi literatur review yang komprehensif hasil penelitian oleh penulis tahun 2017 – 2022, yang menjelaskan tentang asal usul dan komposisi biomarker, paleogeografi sumber fasies dari rembesan minyak (2017-2018), paleofasies dan karakteristik biomarker batuan Paleogen hingga Neogen (2019), kontribusi material terrestrial berdasarkan analisis geokimia dan biostratigrafi (2020), proses tektonik yang terjadi di cekungan tersebut, dan karakterisasi geokimia batuan induk di daerah penelitian (2022). Hasil review menunjukkan sampel permukaan FS-12 yang berupa serpih diduga berasal dari Formasi Kalumpang yang terletak di wilayah Sulawesi Barat. Hasil analisis menunjukkan serpih tersebut mempunyai TOC 0,62%, dengan kuantitas yang cukup. Namun, hasil cross plot antara nilai HI < 50 dan Tmax 499 °C menunjukkan bahwa kategori tersebut terlalu matang dan tidak dapat menghasilkan hidrokarbon. Berdasarkan hasil penelitian ini, dan digabungkan dengan penelitian sebelumnya diperoleh hasil studi geokimia terbaru di wilayah Sulawesi Barat, dan Selat Makassar bagian timur, berupa batuan induk marine shale yang berada di Sulawesi Barat dan batuan induk lacustrine shale di Selat Makassar bagian timur.

Kata kunci: batuan induk, Sulawesi Barat, total karbon organik.

Abstract

The Lariang-Karama Sub-Basin is situated within the Makassar Basin, located in the West Sulawesi region. It has source of rocks which has the potential to produce hydrocarbons and to become a target for new field discoveries in hydrocarbon exploration activities. It is on this basis that the authors are interested in conducting research in the Makassar Basin area and its surroundings, which has been carried out since 2017. This is a comprehensive literature review study from 2017 to 2022, focusing on the origin and composition of biomarkers, paleogeography of facies sources from oil seeps (2017-2018), paleofacies and biomarker characteristics of Paleogene to Neogene rocks (2019), contribution

of terrestrial materials based on geochemical and biostratigraphic analysis (2020), tectonic processes occurring in the basin, and geochemical characterization of source rocks in the research area (2022). The FS-12 surface sample in shale may belong to the Kalumpang Formation, located in West Sulawesi. The results of TOC analysis were 0.62%, indicating sufficient quantity. Based on the results of the cross plot between HI values of < 50 and Tmax of 499 °C, the category was found to be overmature and could not produce hydrocarbons. The results of this research, along with those of previous research and the latest geochemical study, were obtained in the West Sulawesi region and the eastern Makassar Strait. These findings consisted of marine shale source rocks in West Sulawesi and lacustrine shale source rocks in the eastern Makassar Strait.

Keywords: source rock, West Sulawesi, total organic carbon (TOC)

I. INTRODUCTION

Hydrocarbons play a very important role as an energy resource to support the Indonesian economy (Praptisih, 2022). Exploration of hydrocarbons is a crucial activity in the oil and gas sector that involves searching for new reserves to meet production needs and developing existing oil fields. An essential component of exploration is geochemical analysis, which can be carried out on surface samples and hydrocarbon seeps.

Various geochemical analyses have been conducted on surface samples and seeps in West Sulawesi. In research by Sutadiwiria et al. (2017 and 2018) on oil seeps in the Lariang and Karama areas of West Sulawesi, the oleanane saturation found in these seeps indicates that the source rocks in the West Sulawesi area were formed in delta and near-coastal environments. This oil seep is known to originate from Eocene coal and Toraja shale coal from the Kalumpang Formation. The results of biomarker analysis on GC-MS show that the source of oil seep in Lariang can be traced back to terrestrial organic material, with the additional presence of organic materials from algae and herbs (Sutadiwiria, 2019).

The Lariang-Karama Sub-Basin in the Makassar Basin has the potential to produce hydrocarbons. This study aims to refine the predictions regarding the capacity of the area's source rock to produce hydrocarbons by using geochemical analyses of hydrocarbons. These analyses determine the quality of organic materials from rock-eval pyrolysis (REP), the quantity of organic materials from total organic carbon (TOC), the degree of maturity of the hydrocarbon from cross-plot Tmax vs HI, gas chromatography (GC), and gas chromatography/mass spectrometry (GC/MS).

Makassar Basin is located in Indonesia and is bounded by different geographical features. To the north lies the Mangkalihat High, while the Java Sea is situated in the south. In the west, it is bordered by the East Kalimantan mainland, which includes the Mahakam Delta and the SSW-NNE fold structure. On the east, it is bounded by the mainland of West Sulawesi, which has a thrust fold in the SSW-NNE direction.

The Makassar Basin was formed during the pre-Tertiary to Late Tertiary extensional stages, which occurred no earlier than the Early Paleocene (Sutadiwiria et al., 2022). Calvert and Hall (2003) have reported that the research area is located in the eastern part of the Makassar Basin. The basin was formed during the Early Tertiary, Paleocene, and Middle Eocene-Late Eocene. This formation was due to its expansion. The rifting led to the formation of graben and half-graben, and the subsidence formation took place during the early period of the rifting. This process caused the island of Sulawesi to move from east to southeast and separate from the mainland of Sunda/Kalimantan. The area is subjected to extensional compressional forces, which have led to the upliftment, folding, and shearing of the pre-tertiary basement, resulting in the formation of rock depressions and elevations (Figure 1).

II. REGIONAL

2.1 Regional Geology

Table 1. Source rock quality parameters based on TOC (Peters, 1994)

Quality	TOC (%)
Bad	< 0.5
Fair	0.5 – 1
Good	1 – 2
Very Good	2 – 4
Excellent	> 4




Figure 2. Modified Mamuju Regional Stratigraphic Column, West Sulawesi (Calvert and Hall, 2003 and Raharjo et.al, 2012)

3.2 Kerogen Type

Kerogen is the organic material present in source rocks that produce oil and gas. It plays a significant role in determining which hydrocarbons will be formed. Kerogens can be classified into different types based on the combination of chemical elements that it contains, such as carbon (C), hydrogen (H), oxygen (O), nitrogen (N), and sulfur (S). The method used to classify kerogen types is the use of the Van Krevelen diagram (Peters and Moldowan, 1993), which involves plotting the hydrogen index (HI) against the oxygen index (OI) atoms. Additionally, a comparison is made between HI and Tmax values.

3.3 Rock-Eval Pyrolysis (REP)

The parameters used to determine the type of kerogen, oil, and gas products are formed at peak maturity (Peters and Cassa, 1994). The pyrolysis parameters used to determine the type of hydrocarbon are the hydrogen index (HI) and the relationship between the ratio of S2 (a hydrocarbon formed from kerogen in rock-eval pyrolysis due to the decomposition of kerogen) and S3. Figure 3 illustrates an analysis of rock-eval pyrolysis in the study area based on Soppanata (1996).

According to Hunt (1996), Rock-Eval pyrolysis analysis is based on several parameters. S1 (free hydrocarbons) indicates the number of free hydrocarbons that can be vaporized without kerogen degradation. It is

formed at 300 °C, resulting from kerogen during deposition. S2 (pyrolisable hydrocarbon) indicates the number of hydrocarbons produced through the cracking process. It is formed at a temperature of 350–

550 °C. S3 shows the amount of CO₂ content in the rock. It is formed at 300–390 °C. Tmax shows the maximum temperature at the time of the formation of hydrocarbon cracking (S2). It is formed at 435–470 °C.

11

Table 2. Maturity Parameters of Organic Materials (Peters and Cassa, 1994)

Parameter	Maturity			Generation		
	Ro (%)	Tmax	TAI	Bit/TOC	Bitumen	PI
2 Immature	0.20 – 0.60	< 435	1.5 – 1.26	< 0.05	< 50	< 0.10
Early Mature	0.60 – 0.65	435 – 445	2.6 – 2.7	0.05 – 0.10	50 – 100	0.10 – 0.15
Peak Mature	0.65 – 0.90	445 – 450	2.7 – 2.9	0.15 – 0.25	150 – 250	0.25 – 0.40
Late Mature	0.90 – 1.35	450 – 470	2.9 – 3.3	-	-	> 0.40
Post Mature	> 1.35	> 470	> 3.3	-	-	-

3.4 Maturity of Organic Material

The maturity of organic materials was assessed using two parameters: Tmax and Vitrinite Reflectance (Ro). The Tmax value indicates the temperature at which hydrocarbons are formed through cracking (S2) and reach their maximum intensity. Generally, this occurs at 435–470 °C, known as the oil window. On the other hand, Ro is a parameter that indicates the increase in temperature and its effect on the reflectance of vitrinite. As the temperature rises, the reflectance gloss of vitrinite maceral will also increase. Table 2 provides more

information on Ro and its relationship with temperatures.

Gas chromatography (GC) and/or gas chromatography mass spectrometry (GC-MS) analyses were also used in this study. Saturate samples were subjected to gas chromatography (GC) to determine the distribution of isoprenoids and n-alkanes. In the meantime, compounds with low concentrations that could not be separated by gas chromatography (GC) were separated using gas chromatography mass spectrometry (GC-MS) (Waples and Curiale, 1999).




Figure 3. Methodology of Rock-Eval Pyrolysis Analysis (Soppanata, 1996).

The study utilized new data obtained from a surface sample called FS-12, which was taken from the Mamuju area in the Lariang-Karama Sub-Basin, West Sulawesi by Sutadiwiria in 2019. The availability of the data is presented

in Table 3. The lithostratigraphic analysis of the study area revealed that the samples were from the Middle Eocene Kalumpang Formation, which has an upper intertidal environment and comprises shale, as shown in Figure 4.

Table 3. Availability of data used in the research

Surface sample	Sample Identification	6 Lithology	TOC	S1	S2	S3	Tmax	HI	OI	PI	VR (%)	Original References
	Name											
	FS-12	Shale	0.62	0.08	0.20	0.12	499	32	19	0.29	1.3	new data
	FS-2	Coal	30.56	0.61	33.29	21.01	423	109	69	0.02	0.46	Sutadiwiria et al. (2022)
	FS-20	Carbonaceous shale	6,92	0.28	11.94	2.58	429	189	37	0.02	0,38	Sutadiwiria et al. (2022)




Figure 4. Location Map and Geology of Mamuju and Surrounding Areas (Modification of Calvert and Hall (2003) and Raharjo et al. (2012))

IV. RESULTS AND DISCUSSION

The analysis of the maturity level and type of hydrocarbon kerogen led to the conclusion that the FS-12 sample from the Mamuju area, West Sulawesi, belonged to the kerogen type IV (Figures 5 and 6). The hydrogen index (HI) value of the sample was found to be 32 (mg/gr rock/% TOC, Table 3, Figures 5 and 6), which indicates that the kerogen has a lower potential to produce hydrocarbons as the HI value is less than the required 50 mg/gr rock (Waples, 1985). Moreover, the Tmax value of FS-12 was 499 °C, which falls into the post-mature category based on Peters and Cassa (1994, Figure 5).

Previous studies on the surface samples (FS-2 and FS-20) showed high TOC values (30.56% and 6.32%, respectively) and HI values of 109 mg/C and 189 mg/C. Based on the HI value, the FS-2 and FS-20 samples were likely to have the potential to be gas-prone for FS-2 (Type III kerogen), and to be oil and gas-prone for FS-20 (Type II kerogen) (modification of Sutadiwiria et al., 2022) as shown in Table 3, and Figures 5 and 6.




Figure 5. Results of Crossplot Diagram of Hydrogen Index (HI) vs Tmax of Mamuju West Sulawesi (Van Krevelen, 1993 and Waples, 1985)




Figure 6. Results of Crossplot Diagram of Van Krevelen Hydrogen Index (HI) vs Oxygen Index (OI) Mamuju West Sulawesi (Van Krevelen, 1993)

A number of terrestrial components (resin compounds), including the biomarkers oleanane (OL), bicadinane, and gamaserane (Gm), were visible in the mass fragmentogram profile, particularly the m/z 191 ion of the saturated hydrocarbon fraction of GCMS FS-12 (Figure 7). These same biomarkers were also present in the samples FS-2 and FS-20 (Sutadiwiria et al., 2022) (Figures 8 and 9). The components W, T, and R originating from higher plants were identified as the bicadinane. These components were components a, b, c, and d. Ion taraxastana (Tx) also appeared in several samples along with the presence of oleanane.

The ternary diagram of samples FS-2, FS-12, and FS-20 (Figure 10) from Peters et al. (2005) and Huang and Meinschein (1979) indicates a relative abundance plot of sterane ($C_{27} - C_{29}$), which shows a relatively high proportion of C_{29} sterane. The percentages are 48.29%, 33.12%, and 55.27%. In comparison C_{27} shows figures of 26.5%, 36.69%, and 10.73%, while C_{28} has percentages of 25.21%, 30.19%, and 34%, respectively.




Figure 7. m/z 191 partial mass chromatograms showing triterpane distribution of FS-12




Figure 8. m/z 191 partial mass chromatograms showing triterpane distribution of FS-2 (Sutadiwiria et al., 2022)




Figure 9. m/z 191 partial mass chromatograms showing triterpane distribution of FS-20 (Sutadiwiria et al., 2022)




Figure 10. Saturated fraction of the surface sample extract plotted on the ternary diagram according to (a) Huang and Meinschein, 1979 and (b) Peters et al., 2005

The mass fragmentogram ion m/z 217 GCMS profile from samples FS-12, FS-02, and FS-20 further corroborates the ternary diagram (Figures 11, 12, and 13). This suggests that the three samples contain both terrestrial and marine organic materials. According to this update, organic materials derived from terrestrial and marine organic materials are deposited in the

upper intertidal-fluviodeltaic environment as coal (FS-2), shale (FS-12), and carbonaceous shale (FS-20) (Sutadiwiria et al., 2022). This finding also accords with the study (Sutadiwiria et al., 2017) that found a significant amount of resin biomarkers in the source rock, indicating input from the fluviodeltaic environment.




Figure 11. m/z 217 partial mass chromatograms showing sterane distribution of FS-12




Figure 12. m/z 217 partial mass chromatograms showing sterane distribution of FS-2 (Sutadiwiria et al., 2022)




Figure 13. Modified m/z 217 partial mass chromatograms showing sterane distribution of FS-20 (Sutadiwiria et al., 2022)

Upon examination of the findings of the biostratigraphic, petrographic, and radiometric analyses of the current well K-1 in the eastern Makassar Strait (Harsanti et al., 2013, Sutadiwiria et al., 2022), it becomes evident that the Late Cretaceous volcanic materials are located at a depth of 17,340–17,360 feet, while the shales and shale mudstones from the Early

Paleocene and Eocene are marginally lacustrine facies (Geoservices, 2012). The finding of lacustrine shale source rock in the eastern Makassar Strait was made possible by the comparatively higher concentration of C₂₇ sterane in the well K-1 sample, when compared to C₂₈ and C₂₉ (Sutadiwiria et al., 2022).

CONCLUSION

After analyzing the hydrocarbons in the Lariang-Karama Sub-Basin, Mamuju, and West Sulawesi, it was found that the FS-12 sample from the Kalumpang Formation had fair TOC content, but it was not a source rock in the study area due to its post-mature maturity level. The results indicate that it belongs to the Middle Eocene shale lithology and has no potential to become a source rock (Type IV), unlike FS-20 and FS-2, which displays potential for oil and gas generation (Types II and III kerogen, respectively). Following a thorough assessment of post-drilling laboratory study, the West Sulawesi area continues to hold potential to be explored, offering improved prospect⁹ and reduced risks for exploration activities in the region.

ACKNOWLEDGMENTS

The authors would like to express their gratitude for the invaluable support provided by the Department of Geological Engineering, Faculty of Earth and Energy Technology, Universitas Trisakti, in carrying out and presenting this research.

GLOSSARY

Unit	Definition	Symbol
Tmax	Maximum Temperature	
⁸ TOC	Total Organic Carbon	
w.t%	Weight percent (of TOC)	
REP	Rock Eval Pyrolysis	
GC	Gas Chromatography	
GC-MS	Gas Chromatography and Mass Spectroscopy	
⁶ HI	Hydrogen Index	
OI	Oxygen Index	
PI	Production Index	

VR	Vitrinite Reflectance (Ro)	
S1	Free hydrocarbons	
S2	Pyrolysable hydrocarbon	
S3	Amount of CO ₂ content present	
HI = mg/gr rock/% TOC	¹³ the amount of hydrogen relative to the amount of organic carbon present in a sample	
C	Carbon	
H	Hydrogen	
O	Oxygen	
N	Nitrogen	
S	Sulfur	
°C	Degrees Celsius	

REFERENCES

- Calvert, S. J. and Hall, R. (2007). Cenozoic Evolution of the Lariang and Karama Regions, North Makassar Basin, Western Sulawesi, Indonesia. *Petroleum Geoscience*, 13, 353–368.
- Calvert, S. J. and Hall, R. (2003). *The Cenozoic Geology of The Lariang and Karama Regions, Western Sulawesi: New Insight into The Evolution of The Makassar Straits Region*.
- Fraser, T.H., Jackson, B.A., Barber, P.M., Baillie, P., and Myers, K. (2003). The West Sulawesi fold belt and other new plays within the North Makassar Strait—a prospectivity review. *Proceedings Indonesian Petroleum Association, 29th Annual Convention*, p. 429-450.
- Hunt, J.M. (1996). *Petroleum Geochemistry and Geology* (2nd ed.). W.H. Freeman and Company.
- Peters, K. E. and Cassa, M. R. (1994). Applied Source Rock Geochemistry. The Petroleum System from Source to Trap. *The American Association of Petroleum Geologists Memoir*, 60.
- Peters, K.E. dan Moldowan, J.M., 1993. *The*

- Biomarker Guide, Interpreting Molecular Fossils in Petroleum and Ancient Sediments*. Prentice Hall, Englewood Cliffs, New Jersey 07632.
- Praptisih (2022). Karakteristik Geokimia Organik pada Formasi Cibulakan di Daerah Cirebon, Jawa Barat. *Lembaran Publikasi Minyak dan Gas Bumi*, Vol. 56 No. 2, 123 – 135.
- Raharjo, S., Seago, R., Jatmiko, E. W., Hakim, F. B., and Meckel, L. D. (2012). Basin Evolution and Hydrocarbon Geochemistry of The Lariang Karama Basin: Implications for Petroleum System in on Shore West Sulawesi. *Proceedings of the Indonesian Petroleum Association, 36th Annual Convention and Exhibition*.
- Soppanata, M. A. (1996). Buku Praktikum Geokimia Hidrokarbon. Universitas Trisakti. *Buku Praktikum*.
- Sutadiwiria, Y., Burhannudinnur, M., Jambak, M. A., Cahyaningratri, P. R., and Yeftamikha. (2022). Geochemical Characterization of Coal, Carbonaceous Shale, and Marine Shale as Source Rock in West Sulawesi, Indonesia. *Indonesian Journal on Geoscience*, 9(3). <https://doi.org/10.17014/ijog.9.3.303-314>
- Sutadiwiria, Y. (2019). Peranan Biomarker dan Paleofasies pada Rekonstruksi Tektonik Batuan Paleogen – Neogen di Selat Makassar Bagian Timur dan Sulawesi Barat. *PhD Thesis*, Universitas Padjadjaran (*unpublished*).
- Sutadiwiria, Y., Hamdani, A.H., Sendjaja, Y.A., Haryanto, I., dan Yeftamikha (2018). The Biomarker Composition of Some Oil Seeps from West Sulawesi, Indonesia. *Indonesian Journal on Geoscience*, 5, No. 3, 211-220.
- Sutadiwiria, Y., Hamdani, A.H., Andriana, Y., Haryanto, I., and Sunardi, E. (2017). Origin of Oil Seeps in West Sulawesi Onshore, Indonesia: Geochemical Constraints and Paleogeographic Reconstruction of the Source Facies. In *Journal of Geological Sciences and Applied Geology* (Vol. 2, Issue 1).
- Van Krevelen, D. W. (1993): Coal: Typology, Physics, Chemistry, Constitution. 979 pp., 3rd edition, Elsevier, Amsterdam, the Netherlands.
- Waples, D. W. (1985). Geochemistry in Petroleum Exploration. *International Human Resources Development Corporation, Boston*.

Paper SCOG 2024-1

ORIGINALITY REPORT

14%

SIMILARITY INDEX

12%

INTERNET SOURCES

8%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1

ijog.geologi.esdm.go.id

Internet Source

4%

2

www.scbaghdad.edu.iq

Internet Source

3%

3

K.G. Herlienda Elizabeth, Yarra Sutadiwiria, Imam Setiaji Ronoatmojo, P.R. Cahyaningratri, Moehammad Ali Jambak, Andy Livsey.

"Source rock characterization based on geochemical observations using Well BO-1, Central Sumatra Basin", IOP Conference Series: Earth and Environmental Science, 2024

Publication

2%

4

Peter Surjo, Reno Pratiwi, Ratnawati, Rike Yudianti, Eniya L. Dewi, Slamet. "Highly Efficient CuO-doped Titania Nanotube Arrays in Photocatalysis-Electrocoagulation Process for Bacterial Disinfection", Case Studies in Chemical and Environmental Engineering, 2024

Publication

1%

www.karyailmiah.trisakti.ac.id

5	Internet Source	1 %
6	dggs.alaska.gov Internet Source	1 %
7	jptam.org Internet Source	1 %
8	www.mdpi.com Internet Source	1 %
9	Juliette Charbonnel, Sébastien Dubourg, Etienne Testard, Ludovic Broche et al. "Preliminary study of all-solid-state batteries: Evaluation of blast formation during the thermal runaway", iScience, 2023 Publication	<1 %
10	era.library.ualberta.ca Internet Source	<1 %
11	etd.uwc.ac.za Internet Source	<1 %
12	pr.hec.gov.pk Internet Source	<1 %
13	revistas.uis.edu.co Internet Source	<1 %

Exclude bibliography On