


Cover : <https://www.ijmbs.info/index.php/ijmbs/issue/view/48>



Link Editorial Board: <https://www.ijmbs.info/index.php/ijmbs/editorialboard>




# International Journal of Medical and Biomedical Studies

ISSN(online): 2558-8688 || ISSN(print): 2558-868X

[Register](#) [Logi](#)

[HOME](#) [ABOUT](#) [CURRENT ISSUE](#) [ARCHIVES](#) [SUBMISSIONS](#)



[HOME](#) / [Editorial Board](#)

## Editorial Board


**Editor-In-Chief**  
**Sanyam Gandhi**  
110 Denecroft Crescent Road, Hillingdon, London UB109HZ  
(Regulatory affairs strategy lead  
Shire-Takeda, Paddington, London, UK)  
E-mail: [editor@ijmbs.info](mailto:editor@ijmbs.info)

**Associate Editor**  
**Irami Araújo Filho**  
Federal University of Rio Grande do Norte, Onofre Lopes University Hospital, Department of Surgery.  
Av. Nilo Peçanha 620, Petrópolis 59012300 - Natal, RN – Brazil

**Assistant Editor**  
**Mr. Ajay Sharma**  
Sr Research Assistant, Department of Pediatric Research, M at UT M D Anderson Cancer Center, Houston. Texas  
**Chikako Yanagiba**  
Research Associate II, Jasper Therapeutics Inc.,  
Redwood City, California

**Editorial Board**  
**Dr. Bahare Salehi**  
PhD. (Medical Sciences)  
Shahid Beheshti University of Medical Sciences, Tehran, Iran

[MAKE A SUBMISSION](#)



E-mail : [editor@ijmbs.info](mailto:editor@ijmbs.info)

**CURRENT ISSUE**

Atom	1.0
ISS	2.0
ISS	1.0

Open Journal Systems

**KEYWORDS**

sympathovagal balance  
family history  
and  
ketofol  
copla  
retrospective study  
atractumum prunum  
ovarian, contrast, diffusion, mri  
reconstruction  
comparative study radiological  
ed 36  
ketofo  
copla  
mip  
physic  
sacral index  
acrophordons  
arthritis  
lute  
bip  
quill sre

Link Indexing: <https://www.ijmbs.info/index.php/ijmbs>



# International Journal of Medical and Biomedical Studies

ISSN(online): 2389-8898 | ISSN(print): 2389-8897

[Register](#) [Login](#)

[HOME](#) [ABOUT](#) [CURRENT ISSUE](#) [ARCHIVES](#) [SUBMISSIONS](#)

## International Journal of Medical and Biomedical Studies

### International Online and Print Journal

**PubMed and PubMed Central (PMC) (NLM ID: 101738825. Selected citations only)**

Index Copernicus Value 2023: 80.32

\*Impact Factor: 3.465

International Committee of Medical Journal Editors (ICMJE)

**CALL FOR PAPER - Fast Track Publication Hub**

**VOLUME - 09 - ISSUE - 6 - 2025**

If you are facing some problem in Online Submission, kindly Email us on following

E-mails- || [editor@ijmbs.info](mailto:editor@ijmbs.info) || [editorijmbs.info@gmail.com](mailto:editorijmbs.info@gmail.com) ||

MAKE A SUBMISSION



E-mail : [editor@ijmbs.info](mailto:editor@ijmbs.info)

CURRENT ISSUE

atom 1.0

pdf 0.0

pdf 1.0

Open Journal Systems

KEYWORDS



### Aims and Scope

International Journal of Medical and Biomedical Studies (IJMBS) is an international, peer-reviewed, open access, online journal dedicated to the rapid publication of full-length original research papers, short communications, invited reviews, Case studies and editorial commentary and news, Opinions & Perspectives and Book Reviews written at the invitation of the Editor in all areas of the Medical and Biomedical Studies.

International Journal of Medical and Biomedical Studies is an academic journal which aims to publish complete and reliable source of information in the field of medical research providing the fundamental knowledge for further research. IJMBS commits to publish the results of researches in medical practice, risk factors and preventive medicine, safety and effectiveness of treatments, and diagnostic tools that are used to treat any disease. Our journal aims to attract an international audience of Medicine professionals enthusiastic to the most updated and substantial medical progress through the full spectrum of medical research.

### Medical Studies| Biomedical Studies ||

Anatomy, Anthropology, Embryology And Histology, Physiology, Endocrinology, Microbiology: Bacteriology, Mycology, Parasitology And Virology, General Pathology And Pathological Anatomy, Internal Medicine, Pediatrics And Pediatric Surgery, Neurology And Neurosurgery, Surgery, Obstetrics And Gynecology, Otorhinolaryngology, Ophthalmology, Dermatology And Venereology, Radiology, Chest Diseases, Thoracic Surgery And Tuberculosis, Cancer, Public Health, Social Medicine And Epidemiology, Cardiovascular Diseases And Cardiovascular Surgery, Rehabilitation And Physical Medicine, Gerontology And Geriatrics, Developmental Biology And Teratology, Human Genetics, Nuclear Medicine, Anesthesiology, Hematology, Immunology, Serology And Transplantation, Biophysics, Bioengineering And Medical Instrumentation, Urology And Nephrology, Clinical And Experimental Biochemistry, Clinical And Experimental Pharmacology, Arthritis And Rheumatism, Psychiatry, Orthopedic Surgery, Plastic Surgery, Occupational Health And Industrial Medicine, Health Policy, Economics And Management, Drug Literature, Adverse Reaction Titles, Pharmacy, Drug Dependence, Alcohol Abuse And Alcoholism, Environmental Health And Pollution Control, Virology, Gastroenterology, Forensic Science Abstracts, Epilepsy Abstracts, Leprosy And Other Mycobacterial Diseases, Toxicology.

Articles are published as they are accepted and are freely available on the journal's website to facilitate rapid and broad dissemination of research findings to a global audience.

### Top Reasons for publication with us

**Quick Quality Review:** The journal has strong international team of editors and reviewers, Rapid Decision and Publication

**Other features:** DIDS and DOI: Assigned and Implemented the Open Review System (ORS).

**Important Notice:**

Author can now directly send their manuscript as an email attachment to [editor@ijmbs.info](mailto:editor@ijmbs.info)

Link Daftar Isi: <https://www.ijmbs.info/index.php/ijmbs/issue/view/48>

**THE EFFECT OF ATTENTION PROCESS TRAINING (APT-II) ON COGNITIVE AND DAILY LIFE FUNCTIONING IN PATIENTS WITH A MILD COGNITIVE IMPAIRMENT: A RANDOMIZED CONTROLLED TRIAL**

---

Marine Saba, Jean-Paul Rwabihama, Éric Bouvard, Pascale Mettling, Élise Sztulman, Nadège Lemarié, Pascale Piolino, Sophie Blanchet

[PDF](#)

**DETERMINANTS OF POOR SLEEP QUALITY AMONG EDUCATIONAL WORKERS**

---

Chichi Tamara, Yenny

[PDF](#)

**A CLINICAL STUDY OF THYROID DYSFUNCTION IN OVERWEIGHT AND OBESE PATIENT-A HOSPITAL BASED STUDY**

---

Ram Babu Saini, Surendra Singh, Y K Sanadhya

[PDF](#)

**ANEMIA IN SICKLE CELL TRAIT**

---

Bushra Khanam , Srajit Singh , Manish Kumar Saraf, Rudraksh Gautam

[PDF](#)

**THE EFFECT OF SOCIO-DEMOGRAPHIC WITH AUB IN PERIMENOPAUSE**

---

Ankita Jamwal, Rohini Rao

[PDF](#)

**THE MENSTRUAL PATTERN OF AUB IN PERIMENOPAUSE**

---

Ankita Jamwal, Rohini Rao

[PDF](#)

**A STUDY ON CORRELATION OF SERUM PROSTATE SPECIFIC ANTIGEN WITH CLINICAL AND PATHOLOGICAL VARIABLES IN PATIENTS OF PROSTATOMEGALY**

---

Sanjay Patidar , Jaydip Sinh Kathota , Kritika Kamal , Sudhanshu Tiwari, Prashant Nakrani

[PDF](#)

**STUDY OF 24 CASES OF NECROTISING FASCIITIS FOR PREDISPOSING FACTOR AND THEIR MANAGEMENT**

---

Prabhat Kumar Sinha, Neha Thawait, Atul Vyas

[PDF](#)

Link Artikel: <https://www.ijmbs.info/index.php/ijmbs/article/view/2228>



[HOME](#) [ABOUT](#) [CURRENT ISSUE](#) [ARCHIVES](#) [SUBMISSIONS](#)

[HOME](#) / [ARCHIVES](#) / VOL. 5 NO. 10 (2021) / Articles

## DETERMINANTS OF POOR SLEEP QUALITY AMONG EDUCATIONAL WORKERS

**Chichi Tamara**

Undergraduate Program Faculty of Medicine Universitas Trisakti, Jakarta, Indonesia

**Yenny**


Department of Pharmacology & Pharmacy Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

**DOI:** <https://doi.org/10.32553/ijmbs.v5i10.2228>

### ABSTRACT

**Background:** In educational workers, poor sleep quality has a major impact on health and productivity. Although there are many known risk factors for poor sleep quality, data on these risk factors in Indonesian educational workers are lacking. Therefore, this study aims to determine the most influential risk factors for poor sleep quality in Indonesian educational workers.

**Methods:** An observational analysis study with a cross-sectional design was conducted in October-December 2020 involving 81 participants at Yayasan Pendidikan Pelita Raya, Jambi. Sleep quality, vitamin D intake, and job stress were measured using the Pittsburgh Sleep Quality Index (PSQI), Semi-Quantitative Food Frequency Questionnaire (SQ-FFQ), and Health & Safety Executive questionnaires (HSE), respectively. Data analysis was done by multiple logistic regression, with a significant level of  $p < 0.05$ .



 **PDF**

PUBLISHED

2021-10-13

HOW TO CITE

Tamara, C., & Yenny. (2021).  
DETERMINANTS OF POOR SLEEP QUALITY  
AMONG EDUCATIONAL WORKERS.



## DETERMINANTS OF POOR SLEEP QUALITY AMONG EDUCATIONAL WORKERS

Chichi Tamara<sup>1</sup>, Yenny<sup>2</sup>

<sup>1</sup>Undergraduate Program Faculty of Medicine Universitas Trisakti, Jakarta, Indonesia

<sup>2</sup>Department of Pharmacology & Pharmacy Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

**Article Info:** Received 03 September 2021; Accepted 07 October 2021

**DOI:** <https://doi.org/10.32553/ijmbs.v5i10.2228>

**Corresponding author:** Yenny

**Conflict of interest:** No conflict of interest.

### Abstract

**Background:** In educational workers, poor sleep quality has a major impact on health and productivity. Although there are many known risk factors for poor sleep quality, data on these risk factors in Indonesian educational workers are lacking. Therefore, this study aims to determine the most influential risk factors for poor sleep quality in Indonesian educational workers.

**Methods:** An observational analysis study with a cross-sectional design was conducted in October-December 2020 involving 81 participants at Yayasan Pendidikan Pelita Raya, Jambi. Sleep quality, vitamin D intake, and job stress were measured using the Pittsburgh Sleep Quality Index (PSQI), Semi-Quantitative Food Frequency Questionnaire (SQ-FFQ), and Health & Safety Executive questionnaires (HSE), respectively. Data analysis was done by multiple logistic regression, with a significant level of  $p < 0.05$ .

**Results:** The median age was 29 (range 21-55) years, poor sleep quality was found in 30 (37%) participants, 42 (51.9%) of participants had low vitamin D intake, and 48 (59.2%) reported moderate-to-high stress levels. Age  $< 30$  years ( $p < 0.040$ ), moderate-to-high stress level ( $p < 0.013$ ), and low vitamin D intake ( $p < 0.003$ ) were significant associated risk factors for sleep quality in educational workers, with low vitamin D intake being the most influential risk factors.

**Conclusion:** Younger age, stress level, and low vitamin D intake were determinants of poor sleep quality in educational workers, with low vitamin D intake being the most influential risk factors. Management of stress and vitamin D intake are needed to improve sleep quality in educational workers.

**Keywords:** educational workers, Indonesia, poor sleep quality, risk factors, vitamin D intake

### Introduction

The main task of educators and teaching staff is to educate, teach, guide, direct, train, assess, and evaluate students.<sup>(1)</sup> The educators themselves are subject to ever-changing priorities, pressures, resources and reforms.<sup>(2)</sup> Additionally, teachers also bear the work-related social burden, such as caregiving to students. High job demands and social burden render teachers vulnerable to stress, thus affecting sleep quality.<sup>(3)</sup>

Sleep is a condition of the periodic rest of the body and nervous system which is necessary for the learning, performance, and health of workers.<sup>(4)</sup> Poor sleep quality constitutes a community health problem due to its high prevalence from 21.7% to 61% and associated health problems.<sup>(5, 6)</sup> Differences in sleep quality may be due to variations in inter-regional job characteristics and types. Poor sleep quality is also related to weak working memory,<sup>(7, 8)</sup> executive function, decision making, decreased work productivity,<sup>(9)</sup> and increased absenteeism and healthcare costs.<sup>(10)</sup> The long-term effects of poor sleep may impact on worker health and safety.<sup>(11)</sup>

Risk factors of poor sleep quality are age, educational level, income, marital status, chronic disease,<sup>(12)</sup> gender,<sup>(13)</sup> body mass index, physical activity,<sup>(14)</sup> stress level,<sup>(3)</sup> and vitamin D level.<sup>(15)</sup>

Elderly aged 65 years and over tend to have considerably poor sleep quality.<sup>(12)</sup> However, Berhanu et al.<sup>(16)</sup> found a two-fold increased risk of poor sleep quality only in younger persons aged 40 – 49 years.

It is a well-known fact that there are differences in job-stress levels between professions<sup>(17)</sup> and that job stress and poor sleep are interrelated. Kottwitz et al.<sup>(3)</sup> reported an association between job stress and poor sleep in teachers, but did not determine the magnitude of the risk. The objective of Musa et al.<sup>(6)</sup> was to determine the risk factors related to poor sleep quality in high school teachers and found that stress is the only risk factor related to poor sleep quality in multivariate analysis. Teachers undergoing stress have a 1.04-fold significantly higher risk of poor sleep quality than those not experiencing stress. The study of Li et al.<sup>(18)</sup> on oil field workers found that higher work stress score was associated with lower sleep quality. Furthermore, the

study of Visvalingam *et al.* <sup>(19)</sup> on underground workers showed that respondents under stress had a 3.14-fold greater risk of poor sleep quality.

One of the roles of vitamin D in health is sleep regulation, <sup>(20)</sup> but there are few studies on the association of vitamin D intake with sleep quality. The study of Song *et al.* <sup>(15)</sup> showed that inadequate vitamin D intake had a 2.3-fold significantly greater risk of poor sleep quality than did an adequate intake and showed that respondents aged 65 years and above may have comorbid heart failure. The National Health and Nutrition Examination Survey study for 2007–2008 by Grandner *et al.* <sup>(21)</sup> showed that low vitamin D intake was associated with difficulty in maintaining sleep.

To the best of our knowledge there are no studies on the risk factors of sleep quality in Indonesian educational workers. Differences in socio-demographic characteristics, inter-professional stress levels, dietary intakes and inconsistencies in study results emphasize the need for such a study. Therefore, our objective was to find risk factors with the most influential role for poor sleep quality in educational workers.

## Methods

### Design and subject

A cross-sectional study carried out between October and December 2020 at Pelita Raya Education Foundation (Indonesian: Yayasan Pendidikan Pelita Raya), Jambi. The total of respondents is 81 subjects who were recruited using the consecutive sampling method. Inclusion criteria were 19 – 64 year old males and females active as teacher or administrative staff. Exclusion criterion: history of chronic disease (diabetes mellitus, hypertension, gastroesophageal reflux disease, or bronchial asthma).

### Sample size estimation

For the sample size estimation, researchers used the following formulas.

Infinite population formula:

$$n = \frac{Z\alpha^2 \times p \times q}{d^2}$$

Finite population formula:

$$n = \frac{n_0}{(1 + \frac{n_0}{N})}$$

The prevalence of productive adults with poor sleep quality was obtained from the study of Jung *et al.* [5], where the prevalence was 21.7%, with  $Z\alpha$  at  $\alpha = 5\% = 1.96$ , and a measurement accuracy of 0.05. Furthermore, the number of finite populations at data collection sites was 83 persons. Therefore, the estimated minimal sample size in this study was 63 persons.

## Measurements

The collected data consisted of demographic characteristics (age, sex, education, and marital status), work stress, vitamin D intake, and sleep quality. Sleep quality was measured with the Indonesian version of the Pittsburgh Sleep Quality Index (PSQI) [22], a self-rated questionnaire measuring sleep quality in monthly intervals. The PSQI measures 7 sleep components, namely subjective quality, latency, duration, habitual efficiency, disturbance, medication, and daytime dysfunction (concentration problems during the day). The PSQI consists of 19 items, each item is rated on a four-point scale ranging from 0 (no complaints in the previous month) to 3 (three times a week). Total PSQI score is between 0 and 21. Total score  $> 5$  = poor sleep quality and total score  $\leq 5$  = good sleep quality. Indonesian PSQI has adequate reliability and validity. Reliability was supported by internal consistency with a Cronbach's alpha of 0.72. <sup>(22)</sup>

Assessment of dietary vitamin intake was carried out through interviews using a modified Semi-Quantitative Food Frequency Questionnaire (SQ-FFQ), that had been validated for Indonesia and comprised a list of foods, drinks, and vitamin D-containing supplements. <sup>(23)</sup> Vitamin D intake was determined based on household measures which was converted into grams, then analyzed using the Nutrisurvey 2005 nutrition software program and compared to the 2013 Indonesian Recommended Dietary Allowances (IRDA). Vitamin D was categorized into two groups, namely low intake ( $< 15 \mu\text{g/day}$ ) and adequate intake ( $\geq 15 \mu\text{g/day}$ ). <sup>(24)</sup>

Work stress was determined by means of a work-related stress assessment questionnaire based on the Indonesian version of the 2003 Health and Safety Executive (HSE) instrument. <sup>(25)</sup> The assessment was carried out using the scoring method with a Likert scale consisting of 5 choices (never, rarely, quite often, often, and always). There are 35 question items in this questionnaire, in which the lowest total score is 35 and the highest total score is 175. The respondent's work stress level is categorized into the low-stress level (a total score of 140 – 175), medium stress level (a total score of 105 – 139), and the high-stress level (a total score of 70 – 104).

Demographic characteristics (age, gender, education, marital status) were collected using a questionnaire. Age was categorized into two groups, namely  $< 30$  years and  $\geq 30$  years, gender was categorized into male and female, education was categorized as high (diploma or bachelor) and low (high school or below), and marital status was categorized into married and single.

## Data analysis

Descriptive analysis was performed on all collected variables. Categorical data were presented in the number

of respondents (n), percentage (%), odds ratio (OR), and 95% confidence interval (95% CI). Analysis of the association between risk factors and sleep quality was by means of Chi-squared test. Risk factors with a p-value of  $< 0.25$  in the Chi-square test were candidate variables to be analyzed with multiple logistic regressions. Statistical significance for multiple logistic regressions was set at  $p < 0.05$ .

### Ethical

This study protocol was approved by Research Ethics Commission, Faculty of Medicine, Universitas Trisakti, under number 30/KER-FK/10/2020.

### Results

#### Subject characteristics

Table 1 shows the sociodemographic characteristics, work stress levels, vitamin D intake, and sleep quality of the subjects.

**Tabel 1: Distribution of sociodemographic characteristics, work stress level, vitamin D intake, and sleep quality of respondents (n=81)**

Characteristics	Median (Min –Max)	Frequency(n)	Percentage (%)
<b>Age (years)</b>	29 (21 – 55)		
19 – <30		43	53.1
$\geq 30$ – 45		28	34.6
>45 – 55		10	12.3
<b>Gender</b>			
Female		41	50.6
Male		40	49.4
<b>Level of education<sup>a</sup></b>			
High		56	69.1
Low		25	30.9
<b>Marital status<sup>a</sup></b>			
Single		36	44.4
Married		45	55.6
<b>Work stress level<sup>a</sup></b>	138 (104 – 171)		
High		1	1.2
Moderate		47	58
Low		33	40.7
<b>Vitamin D intake<sup>a</sup></b>	12 (1 – 24.6)		
Low		42	51.9
Sufficient		39	48.1
<b>Sleep quality</b>	5 (1 – 15)		
Poor		30	37
Good		51	63

<sup>a</sup>Classification of categorical data: level of education: high (bachelor – diploma), low (senior high school or below); work stress level: high (HSE score of 70 – 104), moderate (HSE score of 105 – 139); low (HSE score of 140 – 175); vitamin D intake (RDA): low ( $< 15$   $\mu\text{g/day}$ ), sufficient ( $\geq 15$  – 20  $\mu\text{g/day}$ ); sleep quality: poor (PSQI score of  $> 5$ ), good (PSQI score of  $\leq 5$ )

*Relationship between sociodemographic characteristics, work stress, and vitamin D intake in the good and poor sleep quality group*

Table 2 shows a comparison of sociodemographic characteristics, work stress, and vitamin D intake between good and poor sleep quality groups. Most respondents with poor sleep quality were  $< 30$  years old (53.5%), unmarried (58.3%), and had low vitamin D intake (54.8%). Chi-squared test results found 4 risk factors with p-value  $< 0.25$ , namely age, marital status, work stress level, and daily vitamin D intake, so that these factors qualify for further analysis using the multiple logistic regression test.



**Table 2: The relationship between sociodemographic characteristics, work stress, and vitamin D intake in the good and poor sleep quality groups (n=81)**

Risk factors	Sleep quality		OR <sup>c</sup>	95%CI <sup>c</sup>	p- value
	Poor N (%)	Good N (%)			
<b>Age (years)</b>					
< 30	23 (53.5)	20 (46.5)	1		
≥ 30			5.09	1.84 – 14.06	0.001 <sup>d</sup>
<b>Gender</b>					
Female	17 (41.5)	24 (58.5)	1		
Male	13 (32.5)	27 (67.5)	1.47	0.59 – 3.65	0.404
<b>Level of education</b>					
Low	11 (44.0)	14 (56.0)	1		
High	19 (33.9)	37 (66.1)	0.65	0.25 – 1.71	0.386
<b>Marital status</b>					
Single	21 (58.3)	15 (41.7)	1		
Married	9 (20)	36 (80)	5.60	2.09 – 15.01	0.001 <sup>d</sup>
<b>Work stress Level<sup>b</sup></b>					
Moderate-to-high	24 (50)	24 (50)	1		
Low	6 (16.2)	27 (81.8)	4.50	1.58 – 12.86	0.004 <sup>d</sup>
<b>Vitamin D Intake<sup>b</sup></b>					
Low	23 (54.8)	19 (45.2)	1		
Sufficient	7 (17.9)	32 (82.1)	5.53	2.00 – 15.33	0.001 <sup>d</sup>

<sup>b</sup>Classification of categorical data: work stress level: moderate-to-high (HSE score of 70 – 139), low (HSE score of 140 – 175); vitamin D intake: low (RDA < 15 µg/day), high (RDA ≥ 15 µg/day)

<sup>c</sup>Abbreviation: OR = odds ratio, CI = confidence interval

<sup>d</sup>Statistical analysis with Chi-square test; p<0.25 fulfill requirement for multiple logistic regression analysis.

*Factors with the most important role in poor sleep quality among educational workers*

Multiple logistic regression test results showed that 3 risk factors were significantly associated with sleep quality, namely age (p < 0.040), work stress level (p < 0.013), and vitamin D intake (p < 0.003), with the most important risk factor for poor sleep quality being low vitamin D intake (see Table 3)

**Table 3: Results of multivariate analysis between age, marital status, work stress level, vitamin D intake, and sleep quality (n=81)**

Risk factors	Adjusted OR	95%CI	p-value
<b>Age (years)</b>			
< 30	1		
≥ 30	4.38	1.07 – 17.9	0.040 <sup>f</sup>
<b>Marital status</b>			
Single	1		
Married	1.94	0.51 – 7.3	0.329
<b>Work stress level<sup>e</sup></b>			
Moderate-to-high	1		
Low	4.96	1.40 – 17.6	0.013 <sup>f</sup>
<b>Vitamin D intake<sup>e</sup></b>			
Low	1		
Sufficient	6.01	1.80 – 20.0	0.003 <sup>f</sup>

<sup>e</sup>categorical cofactor data: work stress level: low (HSE score of 140 – 175), moderate-to-high (HSE score of 70 – 130); vitamin D intake: low (RDA < 15 µg/day), high (RDA ≥ 15 – 20 µg/day)

<sup>f</sup>Statistical analysis with multiple logistic regression test; at level of significance p<0.05.

## Discussion

The prevalence of poor sleep quality in this study was quite high, accounting for 37% (Table 1) and is greater than that in China (21.7%)<sup>(5)</sup> and Thailand (33.7%),<sup>(26)</sup> but lower than in Singapore (42.5%),<sup>(19)</sup> and Malaysia (61%).<sup>(6)</sup> The difference in prevalence between our study and the other studies could be due to differences in socioeconomics, culture, and study location (research setting). Studies conducted in rural areas report a higher prevalence of good sleep quality than do studies in urban areas, due to differences in sleep habits, sleep hygiene, lifestyle, stressors, and others.<sup>(16)</sup>

The median age of the subjects in this study is 29 years (between 21–55) (see Table 1). According to age range, the respondents of this study are of productive age and none of them are in the retirement age. Our study results are in agreement with data from the Indonesian Central Bureau of Statistics stating that the Indonesian population is dominated by the productive age group (15–64 years).<sup>(27)</sup>

Our results show that respondents < 30 years old are at higher risk of poor sleep, which is 4.38 times that of respondents aged ≥ 30 years (see Table 3). These results differ from those of Dong et al. [12] where poor sleep quality is more common in the elderly group, namely 65+ years (35.1%, 95%CI = 30.4–39.8%) compared to 45–54 years (21.4%, 95%CI= 18.0–25.4%;  $p < 0.001$ ).

The study conducted by Berhanu et al.<sup>(16)</sup> showed that the 40 - 49 year age group had twice the risk of poor sleep quality than that of other age groups (Adjusted Odds Ratio (AOR) = 2; 95%CI = 1.1–3.6;  $p < 0.03$ ). Madrid-Valero et al.<sup>(13)</sup> showed that older age is associated with a 1.05-times greater risk of poor sleep quality. The difference in the results of our study and those of other investigators could be due to differences in respondent characteristics, including the younger age range of respondents (21–55 years) and the higher percentage of young persons in our study compared to those of Berhanu et al.[16] and Madrid-Valero et al.[13]

The effects of aging on sleep quality have been well identified. The physiological changes due to aging cause normal elderly persons to take longer to sleep, reduce their ability to maintain sleep, increase the frequency of daytime naps, reduce night-time sleep, and result in slow-wave sleep.

The performance of the circadian system and sleep homeostasis is also reduced. Changes in the amount and pattern of sleep hormone secretion can cause sleep quality disturbances in the elderly.<sup>(28)</sup> The causes of poor sleep quality are not exclusively due to the aging process. The subjective decline in sleep quality starts in young adults and is linear up to the age of 60 years. Thereafter there is a transient increase in quality of life (coinciding with the retirement period due to reduced

work stress). Furthermore, after the age of 66 years, the quality of sleep deteriorates again.<sup>(29)</sup>

In the present study, most of the respondents were < 30 years old (53.1%) and none were elderly (> 60 years). Although sleep quality is definitely influenced by physiological changes due to aging, the aging process between individuals is highly variable. Therefore, it seems that age is not the only factor that plays a role and contributes to the differences between our study results and those of the other studies.

Our study showed no relationship gender and sleep quality (Table 2). In general the relationship between gender as a risk factor and sleep quality is inconsistent. The study of Madrid-Valero et al.<sup>(13)</sup> showed that women had a 1.88-times greater risk than men of a significantly poor sleep quality (OR = 1.88; 95%CI = 1.54–2.28). In contrast, the study by Songkham et al.<sup>(26)</sup> on industrial workers found that the risk of poor sleep quality was 2.74 times greater in men than in women.

It is known that gender differences in sleep quality are influenced by hormonal factors.<sup>(30)</sup> However, the absence of a relationship between gender as a risk factor and poor sleep quality could be due to other factors accounting for poor sleep in younger adults, such as body mass index<sup>(31, 32)</sup> and physical activity.<sup>(14)</sup>

In our study there was no relationship between educational level and sleep quality (Table 2), which agrees with the study results of Musa et al.<sup>(6)</sup> and Deng et al.<sup>(33)</sup> of no association between educational level and sleep quality. Our study also showed no significant relationship between educational level and work stress, which may have been due to the smaller number of respondents (30.9%) with low educational level. This is because in the recruitment of employees certain educational level criteria are applied, including the prospective workload.

The present study found a relationship between marital status and poor sleep quality that was statistically not significant (see Table 3). This is in contrast with the study of Jung et al.<sup>(5)</sup> and Dong et al.<sup>(12)</sup> who found a significant relationship between marital status and sleep quality. The study of Jung et al.<sup>(5)</sup> demonstrated that unmarried respondents experience a significantly poorer quality of life than do married respondents (26.2% vs. 17.6%;  $p < 0.001$ ). Dong et al.<sup>(12)</sup> found that unmarried respondents had a 1.51-times significantly greater risk of poor sleep quality (AOR = 1.51, 95%CI = 1.03–2.22). This agrees with 2013 -2014 US National Health Interview Survey data for single parents.<sup>(34)</sup>

The present study showed that persons with moderate-to-high work stress had a 4.96-times significantly greater risk of experiencing poor sleep quality than those with low work stress. This is higher than in the study by

Musa et al. <sup>(6)</sup> who found that school teachers under stress had a 1.04-fold significantly greater risk of poor sleep quality than did those not under stress (OR= 1.04; 95%CI = 1.01-1.05). Our study also found a greater risk of poor sleep quality as compared with the study of Visvalingam et al. <sup>(19)</sup> on workers in underground workplaces, showing that subjects under stress had a 3.14-times greater risk of poor sleep quality (OR = 3.14; 95%CI = 1.39–7.09). Our study confirms that work stress is a risk factor of poor sleep quality, with many factors that may cause variations in the magnitude of the risk, even in the same profession, such as differences in the characteristics of work requirements. In addition, the study of Li et al. <sup>(18)</sup> showed that a high glucocorticoid level in the body protects against sleep disorder.

Furthermore, our study showed that persons with an inadequate daily intake of vitamin D had a 6.01-times significantly greater risk of poor sleep than did persons with an adequate vitamin D intake (Table 3). The study of Song et al. <sup>(15)</sup> on subjects aged > 65 years with comorbid heart failure also showed that persons with inadequate vitamin D intake had a 2.3-times greater risk of poor sleep than did those with adequate intake.

The National Health and Nutrition Examination Survey for 2007 – 2008 <sup>(24)</sup> showed a relationship between low vitamin D intake (OR = 0.84) and difficulty of maintaining sleep, while the 2005-2016 survey <sup>(35)</sup> showed a relationship between short sleep time and lower intake of calcium, magnesium, and vitamins A, C, D, and E in women. On the other hand, in men, short sleep time is only the effect of insufficient vitamin D intake. Another finding of the 2005-2016 survey was that sleep is also influenced by gender and intake of vitamins and other supplements. <sup>(35)</sup>

One of the many roles of vitamin D in health is sleep regulation. <sup>(20)</sup> Vitamin D receptors and their activating and degrading enzymes are located in brain areas that regulate sleep-wake cycles, e.g. the hypothalamus, the prefrontal cortex, and the substantia nigra. <sup>(36)</sup> Vitamin D is also associated with melatonin metabolism (the hormone regulating circadian rhythms and sleep). <sup>(37)</sup> Vitamin D may also inhibit the release of the inflammatory cytokines TNF- $\alpha$ , IL-1 and prostaglandin D2 which also play a role in sleep regulation. <sup>(38)</sup>

The results of our study indicate that low vitamin D intake is an influential risk factor for poor sleep quality among educational workers. The subjects of the present study were educational workers who habitually performed indoor activities throughout the day. In addition to vitamin D intake, it seems that sun exposure also needs to be considered as a factor in maintaining the adequacy of vitamin D in the body, considering the high prevalence of vitamin D deficiency in Indonesia (43.3% in male adolescents). <sup>(39)</sup>

Existing evidence shows that there is a relationship between vitamin D adequacy and type of work, in which indoor workers are more susceptible to vitamin D deficiency due to lower vitamin D intake and sun exposure than other professions who work outdoors. <sup>(22)</sup> This is consistent with a systematic review conducted by Sowah et al. <sup>(40)</sup> which shows that there are differences in vitamin D levels related to work, in which indoor workers have lower levels of vitamin D than field workers.

The relationship between vitamin D intake and sleep quality is limited, inconsistent, and comes only from observational studies. The differences in the results of those studies could be due to variations in the applied method of measuring vitamin D intake and sleep quality and differences in population and characteristics. The results of the present study, at least, lead to the notion that low vitamin D intake is a risk factor for poor sleep quality in adults, which notion still needs to be confirmed by cohort studies and randomized control trials.

The implications of our study are that the knowledge of risk factors of poor sleep quality in teachers is of benefit to them as well as to health workers. It may also be of use to policy makers who have the task of instituting preventive and organizational measures in stress management and vitamin D intake to improve sleep quality in educational workers.

This study has several limitations, namely: (1) the sleep quality measurements are subjective and self-reported so they are prone to recall bias in which objective measurements using polysomnography would be more valid but difficult to do in the community, (2) the design of this study is cross-sectional so that it cannot confirm the causal effect of various risk factors on sleep quality, (3) vitamin D levels were assessed from food intake using a semi-quantitative food frequency questionnaire which only provided an overview of vitamin D based on intake, whereas a more accurate but more expensive method would be to determine the vitamin D level in the blood, and (4) the HSE questionnaire, although much used in Indonesia for measuring work stress, has not yet been validated in Indonesians.

The strength of this study lies in its collection of epidemiological data on poor sleep quality among educational workers and several risk factors related to sleep quality. Prospective cohort studies are still needed to analyze the causal effect of various risk factors on sleep quality.

## Conclusion

The prevalence of poor sleep quality in educational workers is high. Age of < 30 years, moderate-to-high work stress level, and low vitamin D intake are risk

factors for poor sleep quality in educational workers. Low vitamin D intake is the most influential risk factor for sleep quality in educational workers.

### Acknowledgement

The authors would like to thank the Pelita Jaya Education Foundation, Jambi, Indonesia, the Dean and Vice-Deans of the Faculty of Medicine, Universitas Trisakti, and Richard Tjan DTMH (Mahidol) as proofreader.

### References

1. Financial and Development Supervisory Agency. Governmental Regulation of the Republic of Indonesia No. 19, 2017 Pertaining To Amendments To Governmental Regulation No. 74, 2008 Pertaining To Teachers (in Indonesian)2017 [Available from: <http://www.bpkp.go.id/uu/filedownload/4/140/3508.bpkp>.]
2. Elliot T, Juliason J, Katz N, et al. In Support of educators: strategies that works: EDC Learning transform lives; 2016 [Available from: <https://www.edc.org/sites/default/files/uploads/InSupportOfEducators.pdf>.]
3. Kottwitz MU, Gerhardt C, Pereira D, et al. Teacher's sleep quality: linked to social job characteristics? *Industrial health*. 2018;56(1):53-61.
4. van Dijk DM, van Rhenen W, Murre JMJ, et al. Cognitive functioning, sleep quality, and work performance in non-clinical burnout: The role of working memory. *PloS one*. 2020;15(4):e 0231906.
5. Jung YS, Chae CH, Kim YO, et al. The relationship between serum vitamin D levels and sleep quality in fixed day indoor field workers in the electronics manufacturing industry in Korea. *Annals of occupational and environmental medicine*. 2017;29:25.
6. Musa NA, Moy FM, Wong LP. Prevalence and factors associated with poor sleep quality among secondary school teachers in a developing country. *Industrial health*. 2018;56(5):407-18.
7. Stiver J, Fusco-Gessick B, Moran E, et al. Variable objective sleep quality is related to worse spatial learning and memory in young adults. *Sleep medicine*. 2021;84:114-20.
8. Xie W, Berry A, Lustig C, et al. Poor Sleep Quality and Compromised Visual Working Memory Capacity. *Journal of the International Neuropsychological Society: JINS*. 2019;25(6):583-94.
9. Gingerich SB, Seaverson ELD, Anderson DR. Association Between Sleep and Productivity Loss Among 598 676 Employees From Multiple Industries. *American journal of health promotion: AJHP*. 2018;32(4):1091-4.
10. Firat H, Yuceege M, Kiran S, et al. Absenteeism and Delay to Work Due to Sleep Disorders in the Turkish Adult Population: A Questionnaire-Based National Survey. *Workplacehealth & safety*. 2019;67(1):27-35.
11. Gustavsson K, Wierzbicka A, Matuszczyk M, et al. Sleep among primary care physicians-Association with overtime, night duties and strategies to counteract poor sleep quality. *Journal of sleep research*. 2021;30(1):e13031.
12. Dong X, Wang Y, Chen Y, et al. Poor sleep quality and influencing factors among rural adults in Deqing, China. *Sleep & breathing = Schlaf & Atmung*. 2018;22(4):1213-20.
13. Madrid-Valero JJ, Martínez-Selva JM, Ribeiro do Couto B, et al. Age and gender effects on the prevalence of poor sleep quality in the adult population. *Gaceta sanitaria*. 2017;31(1):18-22.
14. Wang F, Boros S. The effect of physical activity on sleep quality: a systematic review. *European Journal of Physiotherapy*. 2021;23(1):11-8.
15. Song EK, Wu JR. Associations of Vitamin D Intake and Sleep Quality With Cognitive Dysfunction in Older Adults With Heart Failure. *The Journal of cardiovascular nursing*. 2018;33 (4):392-9.
16. Berhanu H, Mossie A, Tadesse S, et al. Prevalence and Associated Factors of Sleep Quality among Adults in Jimma Town, Southwest Ethiopia: A Community-Based Cross-Sectional Study. *Sleep disorders*. 2018;2018:8342328.
17. Nguyen A, Nguyen L, Hoang P, et al. A Review of Occupational Stress among Certain Jobs in Vietnam. *Research in Health Science*. 2019;4:p258.
18. Li X, Gao X, Liu J. Cross-Sectional Survey on the Relationship Between Occupational Stress, Hormone Levels, and the Sleep Quality of Oilfield Workers in Xinjiang, China. *International journal of environmental research and public health*. 2019;16(18).
19. Visvalingam N, Sathish T, Soljak M, et al. Prevalence of and factors associated with poor sleep quality and short sleep in a working population in Singapore. *Occupational Health; June; London, UK*2018. p. 277-87.
20. Muscogiuri G, Barrea L, Scannapieco M, et al. The lullaby of the sun: the role of vitamin D in sleep disturbance. *Sleep medicine*. 2019;54:262-5.
21. Grandner MA, Jackson N, Gerstner JR, et al. Sleep symptoms associated with intake of specific dietary nutrients. *Journal of sleep research*. 2014;23(1):22-34.
22. Setyowati A, Chung MH. Validity and reliability of the Indonesian version of the Pittsburgh Sleep Quality Index in adolescents. *International journal of nursing practice*. 2020:e12856.

23. Rimahardika R, Subagio H, Wijayanti H. Intake of vitamin D and exposure to sunlight in persons working indoors and outdoors, (article in Indonesian). *Journal of Nutrition College*. 2017;6:333.
24. National Institutes of Health Osteoporosis and Related Bone Disease National Resource Center. Calcium and vitamin D: important at every age 2018, Oct [Available from: <https://www.bones.nih.gov/sites/bones/files/pdfs/cal-vit-d-important-any-age-508.pdf>.]
25. Marcatto F, Colautti L, Larese Filon F, et al. The HSE Management Standards Indicator Tool: concurrent and construct validity. *Occupational medicine (Oxford, England)*. 2014;64(5):365-71.
26. Songkham W, Deeluea J, Suksatit B, et al. Sleep quality among industrial workers: related factors and impact. *Journal of Health Research*. 2018;33.
27. Indonesian Central Bureau of Statistics. Indonesian population projection for 2015-2045, SUPAS results 2015 (Revised edition) (in Indonesian) 2018 [Available from: <https://www.bps.go.id/publication/2018/10/19/78d24d9020026ad95c6b5965/proyeksi-penduduk-indonesia-2015-2045-hasil-supas-2015.html>.]
28. Li J, Vitiello MV, Gooneratne NS. Sleep in Normal Aging. *Sleep medicine clinics*. 2018;13(1):1-11.
29. Lemola S, Richter D. The course of subjective sleep quality in middle and old adulthood and its relation to physical health. *The journals of gerontology Series B, Psychological sciences and social sciences*. 2013;68(5):721-9.
30. Pengo MF, Won CH, Bourjeily G. Sleep in Women Across the Life Span. *Chest*. 2018;154(1):196-206.
31. Krističević T, Štefan L, Sporiš G. The Associations between Sleep Duration and Sleep Quality with Body-Mass Index in a Large Sample of Young Adults. *International journal of environmental research and public health*. 2018;15(4).
32. Wang J, Chen Y, Jin Y, et al. Sleep quality is inversely related to body mass index among university students. *Revista da Associacao Medica Brasileira (1992)*. 2019;65(6):845-50.
33. Deng X, Liu X, Fang R. Evaluation of the correlation between job stress and sleep quality in community nurses. *Medicine*. 2020;99(4):e18822.
34. Nugent CN, Black LI. Sleep Duration, Quality of Sleep, and Use of Sleep Medication, by Sex and Family Type, 2013-2014. *NCHS data brief*. 2016(230):1-8.
35. Ikonte CJ, Mun JG, Reider CA, et al. Micronutrient Inadequacy in Short Sleep: Analysis of the NHANES 2005-2016. *Nutrients*. 2019;11(10).
36. Bivona G, Gambino CM, Iacolino G, et al. Vitamin D and the nervous system. *Neurological research*. 2019;41(9):827-35.
37. Romano F, Muscogiuri G, Di Benedetto E, et al. Vitamin D and Sleep Regulation: Is there a Role for Vitamin D? *Current pharmaceutical design*. 2020;26(21):2492-6.
38. McCarty DE, Chesson AL, Jr., Jain SK, et al. The link between vitamin D metabolism and sleep medicine. *Sleep medicine reviews*. 2014;18(4):311-9.
39. Arjana AZ, Devita N, Nurmasitoh T, et al. High Proportion of Vitamin D Deficiency in Male Adolescents in Yogyakarta Indonesia. *Proceedings of the 4th International Conference on Sustainable Innovation 2020—Health Science and Nursing (ICoSIHSN 2020)*; January2021.
40. Sowah D, Fan X, Dennett L, et al. Vitamin D levels and deficiency with different occupations: a systematic review. *BMC public health*. 2017;17(1):519.