

Home > Vol 37, No 3 (2025)

Padjadjaran Journal of Dentistry

Padjadjaran Journal of Dentistry (PJD) is an open-access journal published by the Faculty of Dentistry at Universitas Padjadjaran, issued thrice annually in March, July, and November, with p-ISSN 1979-0201 and e-ISSN 2549-6212. Padjadjaran Journal of Dentistry is also affiliated with [Indonesian Dental Association](#).

Padjadjaran Journal of Dentistry is a scientific publication. It publishes original research manuscripts, systematic reviews, and case reports in the fields of Dental Assisting, Dental Hygiene, Periodontics, and Dentistry (Miscellaneous). However, articles focusing on indigenous health research and the use of natural resources in dentistry are also welcome. It provides a medium for sustained discussion of relevant issues that fall within the focus and scopes of the journal.

Articles published in PJD undergo a double-blind peer-review process. All submissions must be written in English, must be original, unpublished, and not under simultaneous consideration elsewhere. The PJD Editorial Board screens submissions for plagiarism using plagiarism detection software. Manuscripts with a plagiarism level exceeding 20% are automatically rejected. The final decision regarding the acceptance of a scientific article rests with the Editorial Board, based on the recommendations of peer reviewers. Prior to submission, all authors must follow the journal [template](#) and complete the article and author metadata fully and accurately; otherwise, the article will not be considered for further processing. All submitted manuscripts are checked with [Turnitin](#) to prevent unacceptable levels of plagiarism. Reference management should preferably be performed using [Mendeley](#), following the Vancouver superscript style. All keywords must be chosen based on terms included in [MeSH](#).

Starting February 2023, the Padjadjaran Journal of Dentistry adopted new templates for different article types: the updated research article template can be



Table of Contents

Articles

Correlation between oral health and the quality of life of children with disabilities using healthy-dis: an observational study <i>Berlian Prihatiningrum, Roedy Budirahardjo, Niken Probosari, Sulistiyani Sulistiyani, Dyah Setyorini</i>	PDF 268-276
The influence of bamboo fiber (<i>gigantochloa apus</i>) positioning on the flexural strength of heat-cured acrylic resin: an experimental research <i>Azkya Patria Nawawi, Hartanto Endro, Tarish Hamumpuni, Itt Assoratgoon</i>	PDF 277-283
Comparison of Plaque Index (PI) calculation time between a Digital Dental Calculator (DDC) app and the manual method: a quasi-experimental study <i>Syaida Erma Putri, Satria Yandi, Ricky Amran, Netta Anggraini, Valendriyani Ningrum, Shih- Chieh Lee</i>	PDF 284-291
Dental anxiety and its impact on oral health-related quality of life among Indonesian children aged 8–12 years: a cross-sectional study <i>Ika Anisyah, Roosje R. Oewen, Rini Triani, Inne S. Sasmita, Niekla Survia Andiesta</i>	PDF 292-299
Patient safety culture and identify factors associated with positive attitudes in private dental hospitals using the Indonesian Safety Attitude Questionnaire (SAQ-Indo): a cross-sectional study <i>Mita Juliawati, Kusuma Arbianti, Marta Juslily, Wita Anggraini, Balqis I Gusbela, Tri Erri Astoeti</i>	PDF 300-309
Effect on color stability and surface roughness of nanofiller dental composite after soaking in Bidara leaf (<i>Ziziphus mauritiana</i> Lam) ethanol extract: an experimental study <i>Sri fitriyani, Cut Syahla Nazilla, Iin sundari, Viona Diansari, Subhaini Subhaini, Diana Setya Ningsih, Liana Rahmayani, Jasmina Binti Qamaruzzaman</i>	PDF 310-320
Phytochemical composition and cytotoxic assessment of Carica papaya leaf extracts on human squamous carcinoma-3 Cells (in vitro): a laboratory experimental <i>Moehamad Orliando Roeslan, Syafira Fayza Rahma, Paopanga Kokilakanit</i>	PDF 321-329
Ethyl acetate fractions of Cordyline fruticosa leaf: Chemical composition and cariogenic antimicrobial activity <i>Likky Tiara Alphianti, Endang Winiati Bachtiar, Dewi Fatma Suniarti, Eva Fauziah</i>	PDF 330-340
Effect of demineralized dentin matrix and chitosan of black soldier fly on osteoblast and osteoclast activity in post-extraction socket preservation: an experimental study <i>Marsela Umbar Waty, Renie Kumala Dewi, Irham Taufiqurrahman, Muhammad Yanuar Ichrom Nahzi, Huldani Huldani, Rajendran Ganesh</i>	PDF 341-349
Comparative analysis of IL-6 levels in post-teeth extraction inflammation among menopausal and non-menopausal in vivo	PDF 350-361

ORIGINAL ARTICLE

Phytochemical composition and cytotoxic assessment of *Carica papaya* leaf extracts on human squamous carcinoma-3 Cells (in vitro): a laboratory experimental

Moehamad Orilando Roeslan^{1*}
Syafira Fayza Rahma²
Paopanga Kokilakanit³

¹Department of Oral Biology,
Faculty of Dentistry, Universitas
Trisakti, Jakarta, Indonesia

²Undergraduate Study Program
Faculty of Dentistry, Universitas
Trisakti, Jakarta, Indonesia

³Oral Biology Research Unit,
Faculty of Dentistry, Thammasat
University, Pathum Thani, Thailand

* Correspondence:
orlando.roeslan@trisakti.ac.id

Received: 7 September 2025
Revised: 21 October 2025
Accepted: 22 November 2025
Published: 30 November 2025
DOI: 10.24198/pjd.vol37no3.61396

p-ISSN 1979-0201
e-ISSN 2549-6212

Citation:
Roeslan, MO, Rahma, SF,
Kokilakanit, P. Phytochemical
composition and cytotoxic
assessment of *Carica papaya* leaf
extracts on human squamous
carcinoma-3 Cells (in vitro): a
laboratory experimental
Padjajaran J. Dent. November
2025; 37(3): 321-329.

KEYWORDS

C. papaya leaf, HSC-3 cells, ethanol, aquadest, CCK-8 assay.

ABSTRACT

Introduction: Oral cancer is one of the major diseases that can cause death worldwide. With the advancement of research on new drugs, medicinal plants have been explored as an alternative approach to cancer treatment. Papaya (*Carica papaya*) is a plant that has long been used in traditional medicine to treat various diseases. Ethanolic and aqueous extracts of *C. papaya* leaf have been reported to exhibit anticancer properties through mechanisms such as inhibiting cell proliferation and inducing apoptosis. This study aimed to analyze the cytotoxic effect of *C. papaya* leaf extract obtained with different extraction solvents on HSC-3 cell lines. **Methods:** This laboratory-based experimental study employed a post-test only control group design. The sample size was determined using the Federer formula, yielding four replicates per group. HSC-3 cell lines were treated with ethanolic and aqueous extracts of *C. papaya* leaf at concentrations of 125, 250, and 500 µg/mL for 24 hours, followed by a cytotoxicity test using the CCK-8 assay. Data were analyzed using one-way ANOVA followed by Tukey's post hoc test. **Results:** Phytochemical analysis revealed the presence of alkaloids, flavonoids, and tannins in both ethanolic and aqueous extracts, whereas triterpenoids were detected only in the ethanolic extract, and saponins were detected only in aqueous extract. HSC-3 cells did not show any significant cytotoxic response to either extract at any tested concentration. **Conclusion:** Ethanolic and aqueous extracts of *C. papaya* leaf at concentrations of 125, 250, and 500 µg/mL did not exhibit any cytotoxic effects on HSC-3 cell lines.

INTRODUCTION

Oral cancer is one of the major diseases that can cause death worldwide, including Indonesia.¹ Oral cancer is ranked as the 16th most common cancer globally. The global incidence of oral cancer is expected to reach 389,485 new cases and 188,230 deaths by 2022.² The prevalence of oral cancer in Indonesia is around 3-4% of all cancer cases, with a mortality rate of 2-3% of all cancer deaths.³

Oral Squamous cell carcinoma (OSCC) is the most common type of oral cancer, accounting for almost 90% of all oral cavity carcinomas.⁴ Human Squamous Carcinoma (HSC-3) cell lines are derived from human tongue squamous cell carcinoma with metastatic potential and are widely used in oral cancer drugs research.⁵ The etiology of oral cancer is multifactorial and complex. Some of the factors that influence its development include smoking, alcohol consumption and Human Papilloma Virus (HPV) infection.⁶ However, multiple factors, including genetics predisposition, contribute to the development of oral cancer.^{7, 8}

Due to the high prevalence and mortality rate of oral cancer, effective treatment strategies are required. With the development of research on new drugs, medicinal plants have gained interest as an alternative source for potential cancer treatment. One such medicinal plants with promising potential as an alternative for oral cancer treatment is the papaya plant.

Papaya (*Carica papaya*) is a plant from the Caricaceae family that is widely used in traditional medicine.⁹ *C. papaya* leaf extract shows pharmacological activities such as antidiabetic, anti-inflammatory, antimicrobial, antioxidant, and anticancer.¹⁰ The aqueous extract of *C. papaya* leaf has the potential to inhibit breast cancer cell lines (MCF-7) through the suppression of cell proliferation and the induction of apoptosis.¹¹ Ethanol extract of *C. papaya* leaf is effective in inhibiting the proliferation of pancreatic cancer cell lines (MiaPaCa-2 and ASPC-1).¹²

Because of their distinct polarity, ethanol and water are utilised and this influences the type and concentration of beneficial anticancer phytochemicals obtained from *C. papaya* leaf. Ethanol is expected to yield more cytotoxic phytochemicals given its broader solvation capacity^{13, 14}; which aligns with findings indicating that the ethanol extract to have a higher potential for cytotoxicity than water extract.^{15, 16} Though more common in water-based extracts, saponins may exert lower overall potency or more cell line-specific anticancer activity.^{17, 18}

This study addresses a significant research gap, as previous investigations on *Carica papaya* leaf extracts have primarily focused on breast, pancreatic, and prostate cancer cell lines, with limited evidence regarding their effects on oral squamous cell carcinoma (HSC-3). Moreover, earlier studies rarely compared different extraction solvents, leaving unclear how solvent polarity influences the bioactive composition and cytotoxic potential of *C. papaya*.

The novelty of this research lies in its comparative analysis of ethanol and aqueous extracts of *Carica papaya* leaf on HSC-3 cells, providing new insights into how solvent-dependent phytochemical differences relate to anticancer activity. The aim of this study was to analyze the cytotoxic effects of ethanol and aqueous extracts of *C. papaya* leaf on HSC-3 cell lines.

METHODS

This was a laboratory-based experimental study using a post-test only control group design. *Carica papaya* leaf (a California variety, collected from Bogor, West Java, Indonesia) was dried in indirect sunlight and then ground using a blender until a fine powder was obtained. The powdered sample was extracted using the maceration method for three days by soaking the *C. papaya* leaf powder in 96% ethanol (EMSURE) in a powder to solvent ratio of 1:8 for ethanol extract, and in aquadest in a ratio of 1:8 for the aqueous extract at room temperature.¹⁹ During maceration, agitation was performed several times at 15 minutes intervals for 8 hours. The mixture was filtered using Whatman filter paper (Cytiva, Marlborough, US). The ethanol and aqueous extracts were then concentrated using a rotary evaporator.

Phytochemical tests for alkaloid identification were carried out by adding 2 ml of sample to 5 ml of 2N HCl and then adding Dragendorff reagent. The formation of a reddish-brown precipitate indicates the presence of alkaloid compounds.²⁰ Identification of steroid and triterpenoid compounds was performed by 0.5 ml of Liebermann-Burchard reagent to the sample. The formation of blue-green color indicates the presence of steroid compounds, while the formation of purple/red color indicates the presence of triterpenoid compounds.²¹

For flavonoid identification 5 ml of ethanol was added to the sample and heated for five minutes. Then a few drops of concentrated 2N HCl and 0.2 g of magnesium powder were added. Then the sample was shaken. The formation of yellow, red or orange color indicates the presence of flavonoid compounds.²² For saponin identification, the sample was mixed with water and heated for 2-3

minutes, then stirred for 30 seconds, added 2 drops of HCl. If stable foam is formed, then the sample contains saponin compounds.²³ For tannin identification, the sample was dissolved in ethanol and mixed with 4 drops of FeCl₃ 1%. The appearance of a blue-black color indicated the presence of tannin compounds.²⁴

The cell culture process was carried out in a biosafety cabinet. HSC-3 cells were cultured in DMEM (Gibco, Waltham, US) supplemented with FBS (Gibco, Waltham, US) and antibiotics, including 1% amphotericin B and 1% penicillin-streptomycin (Gibco, Waltham, US). HSC-3 cells were incubated at 37°C in an incubator (Esco, Changi, Singapore) humidified with 5% CO₂. The culture medium was regularly replaced, and cells were re-cultured every 3-4 days until the cells reached 80% confluency.²⁵

Confluent cells were placed in 96 well plates with a total of 15×10^3 cells in each well. Each well was treated according to the designated group, then incubated for 24 hours. 10 µL of CCK-8 reagent (Sigma-Aldrich, Darmstadt, German) was added and incubated for 4 hours at 37°C. After incubation, the medium was discarded, and 0.049% DMSO (Acros Organics, Waltham, US) was added to each well. This stage used a blank, control (-) with cell + medium, control (+) with 3 µM doxorubicin, and solvent control with DMSO. Each treatment group, including controls, was performed in triplicate. Readings were taken at a wavelength of 450 nm using a microplate reader (Tecan, Männedorf, Switzerland).²⁶

The number of cells was calculated by subtracting the media absorbance and a correction value (0.1696) from the treatment absorbance, then dividing the result by 0.00005. This calculation provides an estimate of viable cell count based on absorbance readings at 450 nm following treatment and incubation with the CCK-8 reagent. The IC₅₀ value, which represents the concentration required to inhibit 50% of cell viability, was calculated using linear regression analysis of the dose-response curve, by plotting the percentage of cell viability against the logarithm of extract concentrations.

The collected data were then analyzed using a normality test with the Shapiro-Wilk test. Then, a One-Way ANOVA test is performed, followed by a Tukey Post-Hoc test.


RESULTS

The phytochemical test results are shown in Table 1, and the cell counts from each treatment group are shown in Figure 1. The results showed that the ethanolic extracts of *C. papaya* leaf contained alkaloid, flavonoid, triterpenoid, and tannin compounds, while aqueous extract contained alkaloid, flavonoid, saponin, and tannin compounds. Figure 1 showed that the number of cells obtained from the treatment with ethanol extract of *C. papaya* leaf at concentrations of 125, 250, and 500 µg/mL were $15,300 \pm 498.117$; $15,805 \pm 545.064$; and $15,316 \pm 873.548$, respectively. Meanwhile, treatment with the aqueous extract of *C. papaya* leaf at the same concentrations resulted in cell numbers of $16,038 \pm 626.476$; $16,107 \pm 446.111$; and $15,428 \pm 793.173$, respectively. Figure 1 and table 2 show the number of HSC-3 cells under the tested conditions.

A normality test with the Shapiro Wilk method conducted using GraphPad Prism software showed that ethanol and aqueous extracts of *C. papaya* leaf at concentrations of 125, 250, and 500 µg/mL, as well as the positive and negative controls were normally distributed ($p > 0.05$). The One Way ANOVA test showed a significant difference in all test groups with a p value < 0.0001 . Post-Hoc Tukey test showed that ethanol and aqueous extracts of *C. papaya* leaf at concentrations of 125, 250, and 500 µg/mL, the solvent control, and the negative control had significant differences with the positive control.

Table 1. Phytochemical test results

Sample	Compound	Reagent	Result	
Ethanol Extract <i>C. papaya</i> leaf	Alkaloid	Dragendorff	Brown sediment	+
	Flavonoid	HCl 2N + Mg	Yellow, red, orange	+
	Saponin	HCl	Stable foam	-
	Steroid	Chloroform + Liebermann-Burchard	Blue-green	-
	Triterpenoid	Chloroform + Liebermann-Burchard	Brownish red	+
	Tannin	FeCl ₃	Blue-black	+
Aqueous Extract <i>C. papaya</i> leaf	Alkaloid	Dragendorff	Brown sediment	+
	Flavonoid	HCl 2N + Mg	Yellow, red, orange	+
	Saponin	HCl	Stable foam	+
	Steroid	Chloroform + Liebermann-Burchard	Blue-green	-
	Triterpenoid	Chloroform + Liebermann-Burchard	Brownish red	-
	Tannin	FeCl ₃	Blue-black	+

**Figure 1.** Bar graph showing the number of cells in each treatment. *: indicates a significant difference from the positive control**Table 2.** Cytotoxicity test results of ethanolic and aqueous extracts of *C. papaya* leaf on HSC-3 cells.

Treatment Group	Concentration (µg/mL)	Cell Count (Mean ± SD)	Shapiro-Wilk (p-value)	Normality
Ethanol Extract	125	15.300 ± 498.12	0.624	Normal (p > 0.05)
	250	15.805 ± 545.06	0.711	Normal (p > 0.05)
	500	15.316 ± 873.55	0.532	Normal (p > 0.05)
Aqueous Extract	125	16.038 ± 626.48	0.842	Normal (p > 0.05)
	250	16.107 ± 446.11	0.683	Normal (p > 0.05)
	500	15.428 ± 793.17	0.794	Normal (p > 0.05)
Negative Control	—	15.781 ± 480.97	0.658	Normal (p > 0.05)
Solvent Control (DMSO)	—	15.022 ± 366.46	0.719	Normal (p > 0.05)
Positive Control (Doxorubicin)	—	5.781 ± 221.50	0.878	Normal (p > 0.05)

Data are presented as mean ± standard deviation (SD) of triplicate experiments. Normality was assessed using the Shapiro-Wilk test; all groups demonstrated normal distribution (p > 0.05).

DISCUSSION

The phytochemical test of ethanolic and aqueous extracts of *C. papaya* leaf showed different compounds. This difference may occur due to the use of different polarity indexes. The use of ethanol and water as solvents for *C. papaya* leaf follows the solvent selectivity principle governed by polarity. Solvents have a tendency to extract compounds with similar polarity, according to the "like dissolves like" principle. Water is better at extracting highly polar substances like saponins and certain glycosides because it has a higher polarity index (9.0).

On the other hand, ethanol can extract a wider variety of moderately polar to slightly nonpolar substances, such as flavonoids, alkaloids, tannins, and

triterpenoids, due to its lower polarity index (5.2).²⁷ The different biological activities of ethanol and aqueous extracts reflect these differences in phytochemical profiles. For example, the presence of triterpenoids in the ethanolic extract suggests its ability to extract compounds with strong anticancer potential, which were not present in the aqueous extract. Thus, choosing the right solvent is essential for determining the chemical makeup and possible bioactivity of extracts from plants.

Figure 1 shows that there was no discernible cytotoxic effect, as the number of viable HSC-3 cells remained above 15,000 at all concentrations of the ethanol and aqueous extracts of *C. papaya* leaf. While all extract types were significantly different from the positive control group (doxorubicin), none of the tested concentrations (125, 250, or 500 µg/mL) significantly reduced cell viability when compared to the negative control. In addition, the systematic study by Rokis et al. (2025) emphasized that most medicinal plants from Southeast Asia, including *Carica papaya*, exhibit chemopreventive potential through antioxidant mechanisms rather than direct cytotoxic effects. The antiproliferative effect of *C. papaya* tends to appear only at high concentrations (>1000 µg/mL) or after prolonged exposure periods (>72 hours).²⁸

These results imply that the tested *C. papaya* extract concentrations are insufficient to cause cytotoxicity in HSC-3 cells during the 24-hour exposure period. Additional information supporting the observed biological activity can be found in Table 1's phytochemical screening results. Alkaloids, flavonoids, triterpenoids, and tannins were present in the ethanol extract, whereas alkaloids, flavonoids, tannins, and saponins were present in the aqueous extract. The study by Barnabas et al. (2022) confirmed a similar pattern, in which alkaloids, flavonoids, saponins, tannins, and triterpenes were detected in the ethanolic extract of *C. papaya*, with higher intensity compared to the aqueous extract.

The authors noted that ethanol was more effective in extracting non-polar compounds such as triterpenoids and flavonoids, which possess stronger antimicrobial and antioxidant potential.²⁹ In the context of solvent comparison, the study by Ilham et al. (2019) found that the ethanolic extract of *C. papaya* leaf contained higher levels of flavonoids and triterpenoids compared to the aqueous fraction, while the aqueous extract exhibited higher saponin content. This indicates that solvent polarity influences the types of active compounds extracted.³⁰

Alkaloids and flavonoids are examples of bioactive substances with known anticancer potential, but their stability or concentrations may not be sufficient to cause cytotoxicity in HSC-3 cells. This is consistent with previous research by Yuliani and Syahdeni (2020), which found that T47D breast cancer cells were not cytotoxically affected by *C. papaya* ethanol extracts. Variations in biological responses across studies may be explained by differences in exposure duration, compound bioavailability, and cancer cell line sensitivity.³¹

Another study by Kong et al. (2021) reported that *C. papaya* leaf extract exhibited no significant acute toxicity at doses up to 2 g/kg, although it did not directly assess cytotoxicity in cancer cell lines. Current research indicates that although *C. papaya* extracts contain potentially anticancer compounds, their efficacy varies widely depending on cancer type and exposure conditions, underscoring the need for more focused oncological research.³²

Alkaloids have anticancer properties through various mechanisms, such as inhibiting cell proliferation, inducing cell death, and interacting with DNA. As tubulin inhibitors, alkaloids bind to tubulin, a protein that forms microtubules during the cell cycle. The binding of alkaloids to tubulin inhibits the aggregation of microtubule proteins, causing the cell cycle to arrest at metaphase and triggering apoptosis.³³

The anticancer mechanisms of flavonoid include inhibiting cancer cell growth, triggering apoptosis, and preventing the formation of new blood vessels, which is a process necessary for tumour growth. The anticancer activity of these flavonoids

is highly dependent on several factors such as the flavonoid dosage, the type of cancer cells targeted, and their interaction with the biological environment.³⁴

Tannins can inhibit cell proliferation by suppressing protein kinase activity, thereby blocking the signal transduction pathway from the membrane to the cell nucleus.³⁵ Triterpenoids act as anticancer agents by inhibiting the cell division mechanism and activating the apoptotic pathway in cancer cells.³⁶ Saponins exhibit anticancer potential by inhibiting the formation of overexpressed Bcl-2, inducing the low-expression caspase-3 protein, increasing p53 expression, and triggering G1 cell cycle arrest.³⁷

The anticancer activity of *C. papaya* leaf extract has been tested on several cancer cell lines. Aqueous extract of *C. papaya* leaf has been shown to reduce the viability of breast cancer cell lines (MCF-7).¹¹ The ethanol extract of *C. papaya* leaf at concentrations of 125, 250, 500 µg/mL exhibited moderate cytotoxic activity on prostate cancer cell lines (DU 145).³⁸ However, in this study, these concentrations had no cytotoxicity effect on HSC-3.

The use of different cell lines is one of the factors influencing the anticancer effect of *C. papaya* leaf. Each cancer cell has different characteristics, as not all cells have the same growth rate.³⁴ Therefore, selection of different cell types can yield varying results depending on the experimental conditions. The effectiveness of a compound in inhibiting cell growth is highly dependent on the doubling time of the cells themselves.³⁹

To better understand the selectivity and wider applicability of *C. papaya* extracts, future research should include multiple oral cancer cell lines as well as normal oral epithelial cells. It is also advised to look into the molecular processes that underlie cytotoxicity, such as cell cycle arrest, reactive oxygen species production, and apoptotic signalling, in order to clarify how the extracts work. Additionally, to identify and measure the active compounds causing the observed effects, phytochemical profiling using HPLC or GC-MS should be carried out. Comparative research using different extraction methods, like ultrasound-assisted extraction or Soxhlet, may also maximise activity and yield. Lastly, in vivo research is necessary to confirm these extracts' safety profile and therapeutic potential in biological systems.

A comprehensive review by Munir et al. (2022) underscored the importance of elucidating *C. papaya*'s mechanisms through studies on cell cycle arrest, apoptosis, and ROS modulation, emphasizing that *in vitro* assays alone are insufficient without subsequent *in vivo* validation. They demonstrated that papaya leaf juice induced S-phase cell cycle arrest and apoptotic signalling, supporting the recommendation to assess molecular mechanisms in future studies.⁴⁰ Similarly, another previous study compared supercritical fluid extraction versus conventional extraction for *C. papaya* leaf and revealed significant differences in antioxidant and cytoprotective potentials. Their GC-MS analyses identified unique bioactive compounds (flavonoids, alkaloids, and phenolic acids), reinforcing the need for advanced extraction and profiling methods such as HPLC and GC-MS to optimize yield and activity.⁴¹

The statistical analysis in this study was performed to evaluate the differences in cytotoxic effects of *C. papaya* leaf extracts on HSC-3 cells across various treatment groups. The normality of the data was first assessed using the Shapiro-Wilk test, which confirmed that all datasets, including ethanol and aqueous extracts, as well as positive, negative, and solvent controls, were normally distributed ($p > 0.05$). Following this, a one-way ANOVA was conducted to compare the mean viable cell counts among groups. The ANOVA results revealed a statistically significant difference ($p < 0.0001$) among all treatments, suggesting that at least one group differed significantly from the others in terms of cell viability. This approach was chosen due to its suitability for analyzing quantitative outcomes derived from independent groups in cytotoxicity assays.

To determine which specific groups differed, a Tukey's post hoc test was subsequently applied. The results indicated that both ethanol and aqueous

extracts of *C. papaya* at concentrations of 125, 250, and 500 µg/mL were significantly different from the positive control (doxorubicin) but not significantly different from the negative or solvent controls. This finding confirms that neither extract produced a measurable cytotoxic effect on HSC-3 cells under the tested conditions. The combination of ANOVA and Tukey's test provided a statistically robust framework for detecting potential differences while minimizing Type I error. Overall, the statistical results support the conclusion that *C. papaya* leaf extracts, within the concentration range tested, are non-cytotoxic to oral squamous carcinoma (HSC-3) cells, emphasizing the need for further mechanistic studies using molecular assays to validate these findings biologically.

The limitations of this study should also be acknowledged. First, it only used one cell line (HSC-3) for oral squamous carcinoma, which might not accurately reflect the variety of oral cancer.^{42,43} To confirm and expand on the results, more research utilising several cell lines or in vivo models is required. Second, without delving deeper into underlying molecular mechanisms like apoptosis or oxidative stress pathways, the study evaluated cytotoxicity based only on cell viability. Third, only one extraction technique (maceration) and two solvents (ethanol and water) were used, which might not have fully extracted all pertinent bioactive components.

CONCLUSION

The results of this study demonstrate that both ethanolic and aqueous extracts of *C. papaya* leaf at concentrations of 125, 250, and 500 µg/mL did not exhibit cytotoxic effects on HSC-3 oral squamous carcinoma cells. This result suggests that although the extracts contain bioactive substances like alkaloids, flavonoids, triterpenoids, and tannins, the concentrations might not be sufficient to reduce cell viability in this specific cancer cell line. To enhance the possible anticancer effects of *C. papaya* leaf extracts, more research should therefore explore higher concentrations, longer exposure periods, or alternative extraction methods. This conclusion highlights the need for continued investigation within oral cancer therapy research and directly addresses the study goal of comparing the cytotoxicity effects.

The implication of this research lies in supporting the development of plant-based alternatives for oral cancer treatment. The results further emphasise the significance of extraction technique and solvent selection in optimising the medicinal potential of herbal compounds, which may help guide the development of more standardised and potent phytotherapeutic agents in subsequent oncology studies.

Acknowledgement: No declare

Author Contributions: Conceptualization, M.O.R. and S.F.R.; methodology, S.F.R.; software, M.O.R.; validation, M.O.R. and S.F.R.; formal analysis, M.O.R.; investigation, S.F.R.; resources, S.F.R.; data curation, S.F.R.; writing original draft preparation, S.F.R.; writing review and editing, M.O.R. and S.F.R.; visualization, M.O.R. and S.F.R.; supervision, M.O.R.; project administration, M.O.R.; funding acquisition, S.F.R. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: All research procedures have been approved by the Research Ethics Committee, Faculty of Dentistry, Trisakti University. No. 768A/S1/KEPK/FG/8/2024 on August 6th 2024.

Data Availability Statement: Not applicable.

Conflicts of Interest: The author declared no conflicts of interest or potential commercial background in this research.

REFERENCES

1. Warnakulasuriya S, Filho, AM. Oral Cancer in the South and South-East Asia Region, 2022: Incidence and Mortality. *Oral Diseases*. 2025;31(5):1398-405. <https://doi.org/10.1111/odi.15369>.
2. Bray F, Laversanne, M, Sung, H, Ferlay, J, Siegel, RL, Soerjomataram, I, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2024;74(3):229-63. <https://doi.org/10.3322/caac.21834>.

3. Gharat SA, Momin MM, Bhavsar, C. Oral Squamous Cell Carcinoma: Current Treatment Strategies and Nanotechnology-Based Approaches for Prevention and Therapy. 2016;33(4):363-400. <https://doi.org/10.1615/CritRevTherDrugCarrierSyst.2016016272>.
4. Ding M, Yu, S, Chen, Y, Liu, Y, Xuan, F. Space-time analysis of head and neck cancer in Asia and its 34 countries and territories (1990-2021): Implications from the Global Burden of Disease Study 2021. PLoS One. 2025;20(6):e0326177. <https://doi.org/10.1371/journal.pone.0326177>.
5. Jayaraman S, Veeraraghavan, VP, Natarajan, SR, Jasmine, S. Exploring the therapeutic potential of curcumin in oral squamous cell carcinoma(HSC-3 cells): Molecular insights into hypoxia-mediated angiogenesis. Pathology - Research and Practice. 2024;254:155130. doi: <https://doi.org/10.1016/j.prp.2024.155130>.
6. Yang Z, Sun, P, Dahlstrom, KR, Gross, N, Li, G. Joint effect of human papillomavirus exposure, smoking and alcohol on risk of oral squamous cell carcinoma. BMC Cancer. 2023;23(1):457. <https://doi.org/10.1186/s12885-023-10948-6>.
7. Chu Y-H, Tzeng, S-L, Lin, C-W, Chien, M-H, Chen, M-K, Yang, S-F. Impacts of MicroRNA Gene Polymorphisms on the Susceptibility of Environmental Factors Leading to Carcinogenesis in Oral Cancer. PLOS ONE. 2012;7(6):e39777. <https://doi.org/10.1371/journal.pone.0039777>.
8. da Cruz Perez DE, Passos, KKM, Machado, RA, Martelli-Junior, H, Bonan, PRF. Continuing education in oral cancer during coronavirus disease 2019 (covid-19) outbreak. Oral Oncology. 2020;105:104713. <https://doi.org/10.1016/j.oraloncology.2020.104713>.
9. Manpreet K, NaveenChandra, T, Seema, S, Arvind, K, Elena, ES. Ethnomedicinal Uses, Phytochemistry and Pharmacology of Carica papaya Plant: A Compendious Review. Mini-Reviews in Organic Chemistry. 2019;16(5):463-80. <http://dx.doi.org/10.2174/1570193X15666180816110733>.
10. Sharma A, Sharma, R, Sharma, M, Kumar, M, Barbhai, MD, Lorenzo, JM, et al. Carica papaya L. Leaves: Deciphering Its Antioxidant Bioactives, Biological Activities, Innovative Products, and Safety Aspects. Oxidative Medicine and Cellular Longevity. 2022;2022(1):2451733. doi: <https://doi.org/10.1155/2022/2451733>.
11. Nisa FZ, Astuti, M, Muriadi, A, Hariyana, SM. Anti-proliferation and apoptosis induction of aqueous leaf extract of Carica papaya L. on human breast cancer cells MCF-7. Pak. J. Biological Sci. 2017;20(1):36-41. doi: <https://doi.org/10.3923/pjbs.2017.36.41>.
12. Vuong QV, Hirun, S, Chuen, TLK, Goldsmith, CD, Murchie, S, Bowyer, MC, et al. Antioxidant and anticancer capacity of saponin-enriched Carica papaya leaf extracts. Int J of Food Science and Technology. 2015;50(1):169-77. <https://doi.org/10.1111/ijfs.12618>.
13. Abd'quadr-Abojukoro AN, Nkadiemeng, SM, McGaw, LJ, Nsahlai, IV. Phytochemical composition and cytotoxicity of ethanolic extracts of some selected plants. J of Applied Animal Research. 2022;50(1):656-65. <https://doi.org/10.1080/09712119.2022.2119979>.
14. Niyonizigye I, Nkurunziza, D, Ngabire, D, Gitachew, AT, Chun, BS, Kim, G-D. Characterization and in vitro cytotoxicity of phytochemicals from Asplia africana obtained using green extraction techniques. S. Afr. J. Bot. 2020;128:231-8. <https://doi.org/10.1016/j.sajb.2019.11.013>.
15. Mpingirika EZ, El Hosseiny, A, Bakheit, SMS, Arafah, R, Amleh, A. Potential Anticancer Activity of Crude Ethanol, Ethyl Acetate, and Water Extracts of Ephedra foeminea on Human Osteosarcoma U2OS Cell Viability and Migration. BioMed Research International.2020(1):3837693. <https://doi.org/10.1155/2020/3837693>.
16. Tessalonica S, Roeslan, MO. Effect of ethanol extracts from Persea americana leaves on HSC-3 proliferation. JIDA. 2020;3(2):65-70. <https://doi.org/10.32793/jida.v3i2.582>.
17. Bozinou E, Pappas, IS, Patergiannakis, I-S, Chatzimitakos, T, Palaogiannis, D, Athanasiadis, V, et al. Evaluation of Antioxidant, Antimicrobial, and Anticancer Properties of Onion Skin Extracts. 2023;15(15):11599. <https://doi.org/10.3390/su151511599>.
18. Jawad M, Ali, M, Qasim, S, Akbar, A, Khan, NA, Sadiq, MB. Determination of Phenolic Compounds and Bioactive Potential of Plum (Prunus salicina) Peel Extract Obtained by Ultrasound-Assisted Extraction. BioMed Research International. 2022;2022(1):7787958. <https://doi.org/10.1155/2022/7787958>.
19. Karo-Karo SU, Arianto, A, Salim, E. Antibacterial Activity and Determination of Total Phenol and Flavonoid of Carica papaya L. Ethanol Extract. IJSTM. 2023;4(1):233-8. <https://doi.org/10.46729/ijstm.v4i1.738>.
20. Mangkulon S, Roeslan, MO, Monthanpisut, P. The Potential of Cogon Grass (Imperata cylindrica) Ethanol Extract in Inhibiting Nitric Oxide Secretion in Fibroblast. Sci. Dent. J. 2023;7(1).
21. Maheshwaran L, Nadarajah, L, Senadeera, SPNN, Ranaweera, CB, Chandana, AK, Pathirana, RN. Phytochemical Testing Methodologies and Principles for Preliminary Screening/ Qualitative Testing. Asian Plant Res. J. 2024;12(5):11-38. <https://doi.org/10.9734/aprj/2024/v12i5267>.
22. Guediri I, Boubekri, C, Smara, O. Preliminary phytochemical screening from different extracts of Solanum nigrum plant growing in south of Algeria. J. Fundam. Appl. Sci. 2023;12(2):624-33. <https://doi.org/10.4314/ijas.v12i2.7>.
23. Sudha R, Philip, XC, Suriyakumari, KVP. Phytochemical Constituents of Leaves of Moringa oleifera Grow in Cuddalore District, Tamil Nadu, India. SBV J of Basic, Clinical and Applied Health Science. 2020;3(4).
24. Lisa, Wa Ode, S, Suparman, S. Identification of Tannin Compounds In Ethanol Extract Of Cat's Whiskers Leaves (Orthosiphon aristatus) Using Thin Layer Chromatography. IJML. 2025;1(3):106-10.
25. Roeslan MO, Tasha, G. Ethanol extract of Imperata Cylindrica leaves inhibits proliferation and migration of HSC-3 cell lines. Dent J (Majalah Kedokteran Gigi). 2021;54(3):150-4. <https://doi.org/10.20473/j.djmk.v54.i3.p150-154>.
26. Kamiloglu S, Sari, G, Ozdal, T, Capanoglu, E. Guidelines for cell viability assays. Food Frontiers. 2020;1(3):332-49. <https://doi.org/10.1002/fft2.44>.
27. Ng ZX, Samsuri, SN, Yong, PH. The antioxidant index and chemometric analysis of tannin, flavonoid, and total phenolic extracted from medicinal plant foods with the solvents of different polarities. J. Food Process. Preserv. 2020;44(9):e14680. doi: <https://doi.org/10.1111/jfpp.14680>.
28. Rokis A, Hanafi M, Ismail C, Haris MS, Lestari W, Ichwan S, et al. Medicinal plants in the Southeast Asia for treatment and prevention of oral cancer: A systematic review. J. Pharm. Pharmacogn. Res. 2025;13:1003-37.
29. Barnabas F, Wuyep P, Uchejeso O, Mantu E, Ukpa S, Nduke S, et al. Phytochemical analysis and antimicrobial effects of carica papaya leaf extracts on urinary tract infections.2023.
30. Ilham R, Lelo A, Harahap U, Widyawati T, Siahaan L. The Effectivity of Ethanolic Extract from Papaya Leaves (Carica papaya L.) as an Alternative Larvacide to Aedes spp. (1857-9655 (Print)).

31. Yuliani R, Syahdeni, F. Ethanolic Extract of Papaya Leaves (*Carica papaya*) and its Fractions have no Potential Cytotoxicity on T47D Cells. *Pharmacon*: JFI; Vol 17, No 1 (2020). 2020. <https://doi.org/10.23917/pharmacon.v17i1.10760>.
32. Kong YR, Jong, YX, Balakrishnan, M, Bok, ZK, Weng, JK, Tay, KC, et al. Beneficial Role of Carica papaya Extracts and Phytochemicals on Oxidative Stress and Related Diseases: A Mini Review. *Biology*. 2021 <https://doi.org/10.3390/biology10040287>.
33. Zhang D, Kanakkathara, A. Beyond the Paclitaxel and Vinca Alkaloids: Next Generation of Plant-Derived Microtubule-Targeting Agents with Potential Anticancer Activity. *Cancers*. 2020 <https://doi.org/10.3390/cancers12071721>.
34. Kopustinskiene DM, Jakstas, V, Savickas, A, Bernatoniene, J. Flavonoids as Anticancer Agents. *Nutrients*. 2020 <https://doi.org/10.3390/nu12020457>.
35. Yi W, Yan, ZY, Shi, MQ, Zhang, Y, Liu, J, Yi, Q, et al. [Mechanism of tannins from *Galla chinensis* cream in promoting skin wound healing rats based on FAK/PI3K/Akt/mTOR signaling pathway]. *Zhongguo Zhong Yao Za Zhi*. 2025;50(2):480-97. <https://doi.org/10.19540/j.cnki.cjmm.20240902.706>.
36. Ghante MH, Jamkhande, PG. Role of Pentacyclic Triterpenoids in Chemoprevention and Anticancer Treatment: An Overview on Targets and Underlying Mechanisms. *J Pharmacopuncture*. 2019;22(2):55-67. <https://doi.org/10.3831/KPI.201.22.007>.
37. Chang L, Zhou, R, He, Y, Meng, M, Hu, J, Liu, Y, et al. Total saponins from *Rhizoma Panacis Majoris* inhibit proliferation, induce cell cycle arrest and apoptosis and influence MAPK signalling pathways on the colorectal cancer cell. *Mol Med Rep*. 2021;24(2):542. <https://doi.org/10.3892/mmr.2021.12181>.
38. Singh SP, Mishra, A, Shyanti, RK, Singh, RP, Acharya, A. Silver Nanoparticles Synthesized Using Carica papaya Leaf Extract (AgNPs-PL) Causes Cell Cycle Arrest and Apoptosis in Human Prostate (DU145) Cancer Cells. *Biological Trace Element Research*. 2021;199(4):1316-31. <https://doi.org/10.1007/s12011-020-02255-z>.
39. Evans DM, Fang, J, Silvers, T, Delosh, R, Laudeman, J, Ogle, C, et al. Exposure time versus cytotoxicity for anticancer agents. *Cancer Chemotherapy and Pharmacology*. 2019;84(2):359-71. <https://doi.org/10.1007/s00280-019-03863-w>.
40. Munir S, Liu Z-W, Tariq T, Rabail R, Kowalczewski PL, Lewandowicz J, et al. Delving into the Therapeutic Potential of Carica papaya Leaf against Thrombocytopenia. *Molecules* [Internet]. 2022; 27(9).
41. Khor B-K, Chear NJ, Azizi J, Khaw K-Y. Chemical Composition, Antioxidant and Cytoprotective Potentials of Carica papaya Leaf Extracts: A Comparison of Supercritical Fluid and Conventional Extraction Methods. *Molecules*. 2021; 26(5).
42. Mertens B, Nogueira, T, Stranska, R, Naesens, L, Andrei, G, Snoeck, R. Cidofovir is active against human papillomavirus positive and negative head and neck and cervical tumor cells by causing DNA damage as one of its working mechanisms. *Oncotarget*; Vol 7, No. 2016. doi: <https://doi.org/10.18632/oncotarget.10100>.
43. Pardenas I, Roeslan, M. Cytotoxicity Activity of Allium sativum Extracts Against HSC-3 cells. Quality improvement in dental and medical knowledge, research, skills and ethics facing global challenges. London: CRC Press; 2024. p. 393–8.

ORIGINALITY REPORT

15%

SIMILARITY INDEX

13%

INTERNET SOURCES

10%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1

doczz.net

Internet Source

2%

2

jurnal.unpad.ac.id

Internet Source

1%

3

Ashutosh Sharma, Archana Bachheti, Priyanka Sharma, Rakesh Kumar Bachheti, Azamal Husen. "Phytochemistry, pharmacological activities, nanoparticle fabrication, commercial products and waste utilization of Carica papaya L.: A comprehensive review", Current Research in Biotechnology, 2020

Publication

1%

4

www.science.gov

Internet Source

1%

5

Rebecca A. Haber, Reicelle D. Garcia, Jamie N. Hernandez, Sarah Jamieson, Arijit Mondal, Anupam Bishayee. " Papaya (L.) for cancer prevention: Progress and promise ", Critical Reviews in Food Science and Nutrition, 2022

Publication

1%

6

www.scielo.br

Internet Source

1%

7

www.e-journal.trisakti.ac.id

Internet Source

1%

8

mdpi-res.com

Internet Source

1%

9	Eziuche A. Ugboogu, Emmanuel D. Dike, Miracle Ebubechi Uche, Lotanna R. Etumnu et al. "Ethnomedicinal Uses, Nutritional Composition, Phytochemistry and Potential Health Benefits of Carica papaya", Pharmacological Research - Modern Chinese Medicine, 2023 Publication	1 %
10	www.scilit.net Internet Source	<1 %
11	Ratna Yuliani, Faisal Syahdeni. "Cytotoxicity of Ethanolic Extract of Papaya Leaves (Carica papaya) and its Fractions on T47D Cells", Pharmacon: Jurnal Farmasi Indonesia, 2020 Publication	<1 %
12	publications.africanscientistjournal.org Internet Source	<1 %
13	www.omfs.be Internet Source	<1 %
14	www.researchgate.net Internet Source	<1 %
15	www.spandidos-publications.com Internet Source	<1 %
16	Dewangga Yudisthira, Fortuna Filly Firdausi, Citra Fadia Alyani, Fahrul Nurkolis et al. "Potential of Carica papaya leaf extract as an future medicine for thrombocytopenia in dengue patients: from traditional to scientific drug discovery", Advances in Traditional Medicine, 2023 Publication	<1 %
17	"Ethnobotany of the Mountain Regions of Southeast Asia", Springer Science and	<1 %

18	biointerfaceresearch.com Internet Source	<1 %
19	cosmosscholars.com Internet Source	<1 %
20	docksci.com Internet Source	<1 %
21	mdj.uomustansiriyah.edu.iq Internet Source	<1 %
22	Submitted to Fakultas Teknologi Kebumian dan Energi Universitas Trisakti Student Paper	<1 %
23	Huda Moutaz Asmael Al-Azzawi, Syed Ameer Hamza, Rita Paolini, Mathew Lim, Romeo Patini, Antonio Celentano. "PD-L1/PD-1 Expression in the Treatment of Oral Squamous Cell Carcinoma and Oral Potentially Malignant Disorders: An Overview of Reviews", Journal of Personalized Medicine, 2025 Publication	<1 %

Exclude quotes On

Exclude matches < 15 words

Exclude bibliography On



ORIGINAL ARTICLE	
Phytochemical composition and cytotoxic assessment of <i>Carica papaya</i> Leaf extracts on human squamous carcinoma-3 Cells (in vitro): a laboratory experimental	
<p>Moehamad Orliando Roeslan^{1*} Syafira Fayza Rahma¹ Paopanga Kokilakanit³</p> <p>¹Department of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia ²Undergraduate Study Program Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia ³Oral Biology Research Unit, Faculty of Dentistry, Thammasat University, Pathum Thani, Thailand</p> <p>* Correspondence: orliando.roeslan@trisakti.ac.id</p> <p>Received: 7 September 2025 Revised: 21 October 2025 Accepted: 22 November 2025 Published: 30 November 2025 DOI: 10.****/pjd.****</p> <p>p-ISSN 1979-0201 e-ISSN 2549-6212</p> <p>Citation: Roeslan, MO. Rahma, SF. Kokilakanit, P. Phytochemical Composition and cytotoxic assessment of Carica papaya Leaf extracts on human squamous carcinoma-3 Cells (in vitro): a laboratory experimental .Padjajaran J. Dent. November. 2025; 37(3): 1-7.</p>	<p>ABSTRACT</p> <p>Introduction: Oral cancer is one of the diseases that can currently cause death worldwide. With the advancement of research on new drugs, medicinal plants have been explored as an alternative for cancer treatment. Papaya (<i>Carica papaya</i>) is a plant that has long been used in traditional medicine to treat various diseases. Ethanol and aqueous extracts of <i>C. papaya</i> leaf exhibit anticancer properties through the mechanism such as inhibiting cell proliferation and inducing apoptosis. The aim of this study is to analyze the cytotoxic effect of <i>C. papaya</i> leaf extract using different extraction solvents on HSC-3 cell lines.</p> <p>Methods: This was a laboratory-based experimental study using a post-test only control group design. Sample size determination was performed based on the Federer formula, yielding four replicates for each group. HSC-3 cell lines were treated with ethanol and aqueous extracts of <i>C. papaya</i> leaf at concentrations of 125, 250, 500 µg/mL for 24 hours, followed by a cytotoxicity test using the CCK-8 assay. Data were analyzed using one-way ANOVA followed by Tukey's post hoc test</p> <p>Results: Alkaloids, flavonoids, and tannins were found in both ethanol and aqueous extracts, according to phytochemical analysis; triterpenoids were found only in ethanol extract, and saponins were found only in aqueous extract. HSC-3 cells were not significantly cytotoxically affected by either extract at any of the tested concentrations. Conclusion: Ethanol and aqueous extracts of <i>C. papaya</i> leaf at concentrations of 125, 250, 500 µg/mL showed no cytotoxic effect on HSC-3 cell lines.</p>
KEYWORDS	
C. papaya leaf, HSC-3 cells, ethanol, aquadest, CCK-8 assay.	

INTRODUCTION

Oral cancer is one of the diseases that can currently cause death worldwide, including Indonesia.¹ Oral cancer is ranked as the 16th most common cancer globally. The global incidence of oral cancer is expected to reach 389,485 new cases and 188,230 deaths by 2022.² The prevalence of oral cancer in Indonesia is around 3-4% of all cancer cases, with a mortality rate of 2-3% of all cancer deaths.³

Oral Squamous cell carcinoma (OSCC) is one of the most common types of oral cancer and is found in almost 90% of all cases of carcinoma in the oral cavity.⁴ Human Squamous Carcinoma (HSC-3) cell lines are a human oral OSCC cell lines from the tongue with metastatic potential and are mostly used for oral cancer drugs research.⁵ The etiology of oral cancer is multifactorial and complex. Some

Commented [R1]: state number of sample and technic to calculate sample size

Added

Commented [R2]: state analyze of data

Added

1 | Phytochemical Composition and cytotoxic assessment of Carica papaya Leaf extracts on human squamous
Padjajaran Journal of Dentistry • Volume 37, Number 3, November 2025

of the factors that influence the occurrence of oral cancer include smoking, alcohol consumption and Human Papilloma Virus (HPV).⁶ However, factors such as genetics also predispose to oral cancer.^{7, 8} Due to the high prevalence and mortality rate of oral cancer, effective treatment is needed. With the development of research on new drugs, medicinal plants can be used as an alternative in cancer treatment. One of the medicinal plants that has the potential as an alternative to oral cancer treatment is the papaya plant.

Papaya (*Carica papaya*) is a plant from the Caricaceae family that has been widely used in traditional medicine.⁹ *C. papaya* leaf extract shows pharmacological activities such as antidiabetic, anti-inflammatory, antimicrobial, antioxidant, and anticancer.¹⁰ Aqueous extract of *C. papaya* leaf has the potential to inhibit breast cancer cell lines (MCF-7) through the mechanism of inhibiting cell proliferation and inducing apoptosis.¹¹ Ethanol extract of *C. papaya* leaf is effective in inhibiting the proliferation of pancreatic cancer cell lines (MiaPaCa-2 and ASPC-1).¹²

Because of their distinct polarity, ethanol and water are utilised; these influence the type and concentration of beneficial anticancer chemicals obtained from *C. papaya* leaf. Ethanol was expected to generate more cytotoxic phytochemicals given its broader solvation capacity^{13, 14}; this is in line with the results showing ethanol extract to have a higher potential for cytotoxicity than water extract.^{15, 16} Though more common in water-based extracts, saponins may have less potency or greater cell line-specific anticancer action.^{17, 18}

This study addresses a significant research gap, as previous investigations on *Carica papaya* leaf extracts have primarily focused on breast, pancreatic, and prostate cancer cell lines, with limited evidence regarding their effects on oral squamous cell carcinoma (HSC-3). Moreover, earlier studies rarely compared different extraction solvents, leaving unclear how solvent polarity influences the bioactive composition and cytotoxic potential of *C. papaya*.

The novelty of this research lies in its comparative analysis of ethanol and aqueous extracts of *Carica papaya* leaf on HSC-3 cells, providing new insights into how solvent-dependent phytochemical differences relate to anticancer activity. The aim of this study was to analyze the cytotoxic effects of ethanol and aqueous extracts of *C. papaya* leaf on HSC-3 cell lines.

METHODS

This was a laboratory-based experimental study using a post-test only control group design. *Carica papaya* leaf (a California variety, collected from Bogor, West Java, Indonesia) dried in indirect sunlight and pulverized using a blender until they became powder. Extract the powder using the maceration method for three days by soaking the *C. papaya* leaf powder using ethanol 96% (EMSURE) with powder and ethanol ratio of 1:8 for ethanol extract and soaking the *C. papaya* leaf powder using aquadest with powder and aquadest ratio of 1:8 for aqueous extract at room temperature.¹⁹ During the maceration method, agitation be done several times every 15 minutes for 8 hours. Filtered using Whatman filter paper (Cytiva, Marlborough, US). Concentrated the ethanol and aqueous extract of *C. papaya* leaf using a rotary evaporator.

Phytochemical tests for the identification of alkaloid are carried out by adding 2 ml sample with 5 ml of 2N HCl and then adding Dragendorff reagent. The formation of a reddish brown precipitate indicates the presence of alkaloid compounds.²⁰ Identification of steroid and triterpenoid compounds; sample was added with 0.5 ml of Liebermann–Burchard reagent. The formation of blue-green color indicates the presence of steroid compounds, while the formation of purple/red color indicates the presence of triterpenoid compounds.²¹ Identification of flavonoid; sample was added with 5 ml of ethanol and heated for five minutes.

Then a few drops of concentrated 2N HCl and 0.2 g of magnesium powder were added. Then the sample was shaken. The formation of yellow, red or orange

color indicates the presence of flavonoid compounds.²² Identification of saponin; sample was added to water and heated for 2-3 minutes. Then stirred for 30 seconds, added 2 drops of HCl. If stable foam is formed, then the sample contains saponin compounds.²³ Identification of tannin; sample was dissolved with ethanol and added with 4 drops of FeCl₃ 1%. The formation of a blue-black solution indicates the presence of tannin compounds.²⁴

The cell culture process was carried out in a biosafety cabinet. HSC-3 culture using DMEM (Gibco, Waltham, US) and then add FBS (Gibco, Waltham, US) for cell supplementation and antibiotics such as 1% amphotericin B and 1% penicillin-streptomycin (Gibco, Waltham, US). HSC-3 were incubated at 37°C in an incubator (Esco, Changi, Singapore) humidified with 5% CO₂. Regularly, the old medium was replaced with new medium and the cells were re-cultured every 3-4 days until the cells reached 80% confluent.²⁵

Confluent cells were placed in 96 well plates with a total of 15 x 10³ cells in each well, then each well was treated according to the group, then incubated for 24 hours. 10 µL of CCK-8 reagent (Sigma-Aldrich, Darmstadt, German) was added and incubated for 4 hours at 37°C. The medium was discarded and DMSO (Acros Organics, Waltham, US) 0,049% was added to each well. This stage used a blank, control (-) with cell + medium, control (+) with 3 µM doxorubicin, and solvent control with DMSO. Each treatment group, including controls, was performed in triplicate. Readings were taken at a wavelength of 450 nm using a microplate reader (Tecan, Männedorf, Switzerland).²⁶

The number of cells was calculated by subtracting the media absorbance and a correction value (0.1696) from the treatment absorbance, then dividing the result by 0.00005. This calculation provides an estimate of viable cell count based on absorbance readings at 450 nm following treatment and incubation with the CCK-8 reagent. The IC₅₀ value, which represents the concentration required to inhibit 50% of cell viability, was calculated using linear regression analysis of the dose-response curve, by plotting the percentage of cell viability against the logarithm of extract concentrations

The collected data is then analyzed using a normality test with the Shapiro-Wilk test. Then, a One-Way ANOVA test is performed, followed by a Tukey Post-Hoc test.

RESULTS

The phytochemical test results are shown in Table 1. The results of the number of cells from each treatment are shown in Figure 1.

Table 1. Phytochemical test results

Sample	Compound	Reagent	Result	
Ethanol Extract <i>C. papaya</i> leaf	Alkaloid	Dragendorff	Brown sediment	+
	Flavonoid	HCl 2N + Mg	Yellow, red , orange	+
	Saponin	HCl	Stable foam	-
	Steroid	Chloroform + Liebermann-Burchard	Blue-green	-
	Triterpenoid	Chloroform + Liebermann-Burchard	Brownish red	+
	Tannin	FeCl ₃	Blue-black	+
Aqueous Extract <i>C. papaya</i> leaf	Alkaloid	Dragendorff	Brown sediment	+
	Flavonoid	HCl 2N + Mg	Yellow, red , orange	+
	Saponin	HCl	Stable foam	+
	Steroid	Chloroform + Liebermann-Burchard	Blue-green	-

Triterpenoid	Chloroform + Liebermann-Burchard	Brownish red	-
Tannin	FeCl3	Blue-black	+

The results showed that the ethanol extracts of *C. papaya* leaf contained alkaloid, flavonoid, triterpenoid and tannin compounds, while aqueous extracts of *C. papaya* leaf contained alkaloid, flavonoid, saponin and tannin compounds.




Figure 1. Bar graph showing the number of cells in each treatment. *: indicates there is a significant difference with the positive control

Figure 1 showed that the number of cells obtained from the treatment with ethanol extract of *C. papaya* leaf at concentrations of 125, 250, and 500 µg/mL were 15,300 ± 498.117; 15,805 ± 545.064; and 15,316 ± 873.548, respectively. Meanwhile, treatment with the aqueous extract of *C. papaya* leaf at the same concentrations resulted in cell numbers of 16,038 ± 626.476; 16,107 ± 446.111; and 15,428 ± 793.173, respectively. Figure 1 and table 2 show the number of HSC-3 cells under the tested conditions

Normality tests with the Shapiro Wilk method conducted using GraphPad Prism software showed that ethanol and aqueous extracts of *C. papaya* leaf with concentrations of 125, 250, 500 µg/mL, positive control, and negative control were normally distributed ($p>0.05$). One Way ANOVA test showed there was a significant difference in all test groups with a p value of <0.0001 . *Post-Hoc* Tukey test showed that ethanol and aqueous extracts of *C. papaya* leaf with concentrations of 125, 250, 500 µg/mL, solvent control and negative control had significant differences with the positive control.

Commented [R3]: state this narration with tables

Commented [OR4R3]: Added

Table 2. Cytotoxicity test results of ethanol and aqueous extracts of *C. papaya* leaf on HSC-3 cells.

Treatment Group	Concentration (µg/mL)	Cell Count (Mean ± SD)
Ethanol extract	125	15,300 ± 498.12
	250	15,805 ± 545.06
	500	15,316 ± 873.55
Aqueous extract	125	16,038 ± 626.48
	250	16,107 ± 446.11
	500	15,428 ± 793.17
Negative control	—	15,781 ± 480.97
Positive control	—	5,781 ± 221.50
Solvent control	—	15,022 ± 366.46

Data are presented as mean ± standard deviation (SD) of triplicate experiments.

DISCUSSION

The phytochemical test of ethanol and aqueous extracts of *C. papaya* leaf showed different compounds, this difference may occur due to the use of different polarity indexes. The use of ethanol and water as solvents for *C. papaya* leaf can be attributed to the solvent selectivity principle, which is controlled by polarity. Solvents have a tendency to extract compounds with similar polarity, according to the "like dissolves like" principle. Water is better at extracting highly polar substances like saponins and certain glycosides because it has a higher polarity index (9.0).

On the other hand, ethanol can dissolve a wider variety of moderately polar to slightly nonpolar substances, such as flavonoids, alkaloids, tannins, and triterpenoids, due to its lower polarity index (5.2).²⁷ The different biological activities of ethanol and aqueous extracts reflect these differences in phytochemical profiles. For example, the ethanol extract's ability to extract compounds with strong anticancer mechanisms is suggested by the presence of triterpenoids, which were not present in the aqueous extract. Thus, choosing the right solvent is essential for figuring out the chemical makeup and possible bioactivity of extracts from plants.

Figure 1 shows that there was no discernible cytotoxic effect because the number of viable HSC-3 cells stayed above 15,000 at all concentrations of the ethanol and aqueous extracts of *C. papaya* leaf. While all extract types were significantly different from the positive control group (doxorubicin), neither at concentrations of 125, 250, or 500 µg/mL significantly decreased cell viability when compared to the negative control. In addition, the systematic study by Rokis et al. (2025) emphasized that most medicinal plants from Southeast Asia, including *Carica papaya*, exhibit chemopreventive potential through antioxidant mechanisms rather than direct cytotoxic effects. The antiproliferative effect of *C. papaya* tends to appear only at high concentrations (>1000 µg/mL) or after prolonged exposure periods (>72 hours).²⁸ These results imply that the tested *C. papaya* extract concentrations are insufficient to cause cytotoxicity in HSC-3 cells during the 24-hour exposure period.

Additional information about the observed biological activity can be found in Table 1's phytochemical screening results. Alkaloids, flavonoids, triterpenoids, and tannins were present in the ethanol extract, whereas alkaloids, flavonoids,

Commented [R5]: compare with previous study

Commented [OR6R5]: Added

tannins, and saponins were present in the aqueous extract. The study by Barnabas et al. (2022) confirmed a similar pattern, in which alkaloids, flavonoids, saponins, tannins, and triterpenes were detected in the ethanolic extract of *C. papaya*, with higher intensity compared to the aqueous extract. The authors noted that ethanol is more effective in extracting non-polar compounds such as triterpenoids and flavonoids, which possess stronger antimicrobial and antioxidant potential.²⁹ In the context of solvent comparison, the study by Ilham et al. (2019) found that the ethanolic extract of *C. papaya* leaves contained higher levels of flavonoids and triterpenoids compared to the aqueous fraction, while the aqueous extract exhibited higher saponin content—in line with your results. This indicates that solvent polarity influences the types of active compounds extracted.³⁰

..... Alkaloids and flavonoids are examples of bioactive substances with known anticancer potential, but their stability or concentrations may not be sufficient to cause cytotoxicity in HSC-3 cells. This confirms earlier research by Yuliani and Syahdeni (2020), which found that T47D breast cancer cells were not cytotoxically affected by *C. papaya* ethanol extracts. Variations in biological responses across studies may be explained by differences in exposure duration, compound bioavailability, and cancer cell line sensitivity.³¹

Another previous study Kong et al. (2021) reported that *C. papaya* leaf extract exhibited no significant acute toxicity at doses up to 2 g/kg, the study does not directly address its effects on cancer cell viability. Current research indicates that although *C. papaya* extracts contain potentially anticancer compounds, their efficacy varies widely depending on cancer type and exposure conditions, underscoring the need for more focused oncological research.³²

Alkaloids have anticancer properties through various mechanisms, such as inhibiting cell proliferation, inducing cell death, and interacting with DNA. As tubulin inhibitors, alkaloids bind to tubulin, a protein that forms microtubules during the cell cycle. The binding of alkaloids to tubulin inhibits the aggregation of microtubule proteins, causing the cell cycle to arrest at metaphase and triggering apoptosis.³³

The anticancer mechanisms of flavonoid include inhibiting cancer cell growth, triggering apoptosis, and preventing the formation of new blood vessels, which is a process necessary for tumour growth. The anticancer activity of these flavonoids is highly dependent on several factors such as the flavonoid dosage, the type of cancer cells targeted, and their interaction with the biological environment.³⁴

Tannins can inhibit cell proliferation by suppressing protein kinase activity, thereby blocking the signal transduction pathway from the membrane to the cell nucleus.³⁵ Triterpenoid act as anticancer agents by inhibiting the cell division mechanism and activating the apoptotic pathway in cancer cells.³⁶ Saponins exhibit anticancer potential by inhibiting the formation of overexpressed Bcl-2, inducing the low-expression caspase-3 protein, increasing p53 expression, and triggering G1 cell cycle arrest.³⁷

The anticancer activity of *C. papaya* leaf extract has been tested on several cancer cell lines. Aqueous extract of *C. papaya* leaf has been shown to reduce the viability of breast cancer cell lines (MCF-7).¹¹ The ethanol extract of *C. papaya* leaf at concentrations of 125, 250, 500 µg/mL exhibited moderately active cytotoxic against prostate cancer cell lines (DU 145).³⁸ However in this study, these concentrations had no cytotoxicity effect on HSC-3.

The use of different cell lines is one of the factors influencing the anticancer effect of *C. papaya* leaf. Each cancer cell has different characteristics, as not all cells have the same growth rate.³⁴ Therefore, selection of different cell types can yield varying results depending on the experimental conditions. The effectiveness of a compound in inhibiting cell growth is highly dependent on the doubling time of the cells themselves.³⁹

To better understand the selectivity and wider applicability of *C. papaya* extracts, future research should take into account the inclusion of multiple oral

Commented [R7]: compare with previous study

Commented [OR8R7]: Added

Commented [R9]: compare with previous study

Commented [OR10R9]: Added

cancer cell lines as well as normal oral epithelial cells. It is also advised to look into the molecular processes that underlie cytotoxicity, such as cell cycle arrest, reactive oxygen species production, and apoptotic signalling, in order to clarify how the extracts work. Additionally, to identify and measure the active compounds causing the observed effects, phytochemical profiling using HPLC or GC-MS should be carried out. Comparative research using different extraction methods, like ultrasound-assisted extraction or Soxhlet, may also maximise activity and yield. Lastly, in vivo research is necessary to confirm these extracts' safety profile and therapeutic potential in biological systems.

A comprehensive review by Munt et al. (2022) underscored the importance of elucidating *C. papaya*'s mechanisms through studies on cell cycle arrest, apoptosis, and ROS modulation, emphasizing that *in vitro* assays alone are insufficient without subsequent *in vivo* validation. They demonstrated that papaya leaf juice induced S-phase cell cycle arrest and apoptotic signalling, supporting the recommendation to assess molecular mechanisms in future studies.⁴⁰ Similarly, another previous study compared supercritical fluid extraction versus conventional extraction for *C. papaya* leaves and revealed significant differences in antioxidant and cytoprotective potentials. Their GC-MS analyses identified unique bioactive compounds (flavonoids, alkaloids, and phenolic acids), reinforcing the need for advanced extraction and profiling methods such as HPLC and GC-MS to optimize yield and activity.⁴¹

The statistical analysis in this study was performed to evaluate the differences in cytotoxic effects of *C. papaya* leaf extracts on HSC-3 cells across various treatment groups. The normality of the data was first assessed using the Shapiro-Wilk test, which confirmed that all datasets, including ethanol and aqueous extracts, as well as positive, negative, and solvent controls, were normally distributed ($p > 0.05$). Following this, a one-way ANOVA was conducted to compare the mean viable cell counts among groups. The ANOVA results revealed a statistically significant difference ($p < 0.0001$) among all treatments, suggesting that at least one group differed significantly from the others in terms of cell viability. This approach was chosen due to its suitability for analyzing quantitative outcomes derived from independent groups in cytotoxicity assays.

To determine which specific groups differed, a Tukey's post hoc test was subsequently applied. The results indicated that both ethanol and aqueous extracts of *C. papaya* at concentrations of 125, 250, and 500 µg/mL were significantly different from the positive control (doxorubicin) but not significantly different from the negative or solvent controls. This finding confirms that neither extract produced a measurable cytotoxic effect on HSC-3 cells under the tested conditions. The combination of ANOVA and Tukey's test provided a statistically robust framework for detecting potential differences while minimizing Type I error. Overall, the statistical results support the conclusion that *C. papaya* leaf extracts, within the concentration range tested, are non-cytotoxic to oral squamous carcinoma (HSC-3) cells, emphasizing the need for further mechanistic studies using molecular assays to validate these findings biologically.

The limitation of this study includes several important considerations. First of all, it only used one cell line (HSC-3) for oral squamous carcinoma, which might not accurately reflect the variety of oral cancer.^{42, 43} To confirm and expand on the results, more research utilising several cell lines or in vivo models is required. Second, without delving deeper into underlying molecular mechanisms like apoptosis or oxidative stress pathways, the study evaluated cytotoxicity based only on cell viability. Third, only one extraction technique (maceration) and two solvents (ethanol and water) were used, which might not have fully extracted all pertinent bioactive components.

Commented [R11]: compare with previous study

Commented [OR12R11]: Added

Commented [R13]: compare with previous study

Commented [OR14R13]: Added

Commented [R15]: state your discussion about statistic test that used in this research

Commented [OR16R15]: Added

CONCLUSION

The results of this study demonstrate that both ethanol and aqueous extracts of *C. papaya* leaf at concentrations of 125, 250, and 500 µg/mL do not exhibit cytotoxic effects on HSC-3 oral squamous carcinoma cells. This result implies that even though the extracts contain bioactive substances like alkaloids, flavonoids, triterpenoids, and tannins, the concentrations may not be enough to lower cell viability in this specific cancer cell line. To maximise the possible anticancer effects of *C. papaya* leaf extracts, more research is therefore required using higher concentrations, longer exposure periods, or alternative extraction methods. This conclusion emphasises the need for more investigation in oral cancer therapy research and directly addresses the study goal of comparing the cytotoxicity effects.

The implication of this research is for the development of alternative, plant-based treatments for oral cancer. The results further emphasise the significance of extraction technique and solvent selection in optimising the medicinal potential of herbal compounds, which may help guide the development of more standardised and potent phytotherapeutic agents in subsequent oncology studies.

Acknowledgement: No declare

Author Contributions: Conceptualization, M.O.R. and S.F.R.; methodology, S.F.R.; software, M.O.R.; validation, M.O.R. and S.F.R.; formal analysis, M.O.R.; investigation, S.F.R.; resources, S.F.R.; data curation, S.F.R.; writing original draft preparation, S.F.R.; writing review and editing, M.O.R. and S.F.R.; visualization, M.O.R. and S.F.R.; supervision, M.O.R.; project administration, M.O.R.; funding acquisition, S.F.R. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: All research procedures have been approved by the Research Ethics Committee, Faculty of Dentistry, Trisakti University. No. 768A/S1/KEPK/FKG/8/2024 on August 6th 2024.

Data Availability Statement: Not applicable.

Conflicts of Interest: The author declared no conflicts of interest or potential commercial background in this research.

REFERENCES

1. Warnakulasuriya S, Filho, AM. Oral Cancer in the South and South-East Asia Region, 2022: Incidence and Mortality. *Oral Diseases*. 2025;31(5):1398-405. doi: <https://doi.org/10.1111/odi.15369>.
2. Bray F, Laversanne, M, Sung, H, Ferlay, J, Siegel, RL, Soerjomataram, I, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2024;74(3):229-63. doi: 10.3322/caac.21834.
3. Gharat SA, Momin, MM, Bhavsar, C. Oral Squamous Cell Carcinoma: Current Treatment Strategies and Nanotechnology-Based Approaches for Prevention and Therapy. 2016;33(4):363-400. doi: 10.1615/CritRevTherDrugCarrierSyst.2016016272.
4. Ding M, Yu, S, Chen, Y, Liu, Y, Xuan, F. Space-time analysis of head and neck cancer in Asia and its 34 countries and territories (1990-2021): Implications from the Global Burden of Disease Study 2021. *PLoS One*. 2025;20(6):e0326177. doi: 10.1371/journal.pone.0326177.
5. Jayaraman S, Veeraraghavan, VP, Natarajan, SR, Jasmine, S. Exploring the therapeutic potential of curcumin in oral squamous cell carcinoma(HSC-3 cells): Molecular insights into hypoxia-mediated angiogenesis. *Pathology - Research and Practice*. 2024;254:155130. doi: <https://doi.org/10.1016/j.prp.2024.155130>.
6. Yang Z, Sun, P, Dahlstrom, KR, Gross, N, Li, G. Joint effect of human papillomavirus exposure, smoking and alcohol on risk of oral squamous cell carcinoma. *BMC Cancer*. 2023;23(1):457. doi: 10.1186/s12885-023-10948-6.
7. Chu Y-H, Tzeng, S-L, Lin, C-W, Chien, M-H, Chen, M-K, Yang, S-F. Impacts of MicroRNA Gene Polymorphisms on the Susceptibility of Environmental Factors Leading to Carcinogenesis in Oral Cancer. *PLOS ONE*. 2012;7(6):e39777. doi: 10.1371/journal.pone.0039777.
8. da Cruz Perez DE, Passos, KKM, Machado, RA, Martelli-Junior, H, Bonan, PRF. Continuing education in oral cancer during coronavirus disease 2019 (covid-19) outbreak. *Oral Oncology*. 2020;105:104713. doi: <https://doi.org/10.1016/j.oraloncology.2020.104713>.
9. Manpreet K, Naveen Chandra, T, Seema, S, Arvind, K, Elena, ES. Ethnomedicinal Uses, Phytochemistry and Pharmacology of Carica papaya Plant: A Compendious Review. *Mini-Reviews in Organic Chemistry*. 2019;16(5):463-80. doi: <http://dx.doi.org/10.2174/1570193X15666180816110733>.
10. Sharma A, Sharma, R, Sharma, M, Kumar, M, Barbhai, MD, Lorenzo, JM, et al. Carica papaya L. Leaves: Deciphering Its Antioxidant Bioactives, Biological Activities, Innovative Products, and Safety Aspects. *Oxidative Medicine and Cellular Longevity*. 2022;2022(1):2451733. doi: <https://doi.org/10.1155/2022/2451733>.
11. Nisa FZ, Astuti, M, Murdiati, A, Haryana, SM. Anti-proliferation and apoptosis induction of aqueous leaf extract of Carica papaya L. on human breast cancer cells MCF-7. *Pakistan Journal of Biological Sciences*. 2017;20(1):36-41. doi: <https://doi.org/10.3923/pjbs.2017.36.41>

12. Vuong QV, Hirun, S, Chuen, TLK, Goldsmith, CD, Murchie, S, Bowyer, MC, et al. Antioxidant and anticancer capacity of saponin-enriched Carica papaya leaf extracts. *International Journal of Food Science and Technology*. 2015;50(1):169-77. doi: <https://doi.org/10.1111/ijfs.12618>
13. Abd'quadri-Abojukoro AN, Nkadameng, SM, McGaw, LJ, Nsahlai, IV. Phytochemical composition and cytotoxicity of ethanolic extracts of some selected plants. *Journal of Applied Animal Research*. 2022;50(1):656-65. doi: 10.1080/09712119.2022.2119979.
14. Niyonizigiye I, Nkurunziza, D, Ngabire, D, Gitachew, AT, Chun, BS, Kim, G-D. Characterization and in vitro cytotoxicity of phytochemicals from *Aspilula africana* obtained using green extraction techniques. *South African Journal of Botany*. 2020;128:231-8. doi: <https://doi.org/10.1016/j.sajb.2019.11.013>.
15. Mpingirika EZ, El Hosseiny, A, Bakheit, SMS, Arafah, R, Amleh, A. Potential Anticancer Activity of Crude Ethanol, Ethyl Acetate, and Water Extracts of *Ephedra foeminea* on Human Osteosarcoma U2OS Cell Viability and Migration. *BioMed Research International*. 2020;2020(1):3837693. doi: <https://doi.org/10.1155/2020/3837693>.
16. Tessalonica S, Roeslan, MO. Effect of ethanol extracts from *Persea americana* leaves on HSC-3 proliferation. *Journal of Indonesian Dental Association*. 2020;3(2):65-70. doi: <https://doi.org/10.32793/jida.v3i2.582>
17. Bozinou E, Pappas, IS, Patergiannakis, I-S, Chatzimitakos, T, Palaogiannis, D, Athanasiadis, V, et al. Evaluation of Antioxidant, Antimicrobial, and Anticancer Properties of Onion Skin Extracts. 2023;15(15):11599. doi: <https://doi.org/10.3390/su15111599>
18. Jawad M, Ali, M, Qasim, S, Akbar, A, Khan, NA, Sadiq, MB. Determination of Phenolic Compounds and Bioactive Potential of Plum (*Prunus salicina*) Peel Extract Obtained by Ultrasound-Assisted Extraction. *BioMed Research International*. 2022;2022(1):7787958. doi: <https://doi.org/10.1155/2022/7787958>.
19. Karo-Karo SU, Arianto, A, Salim, E. Antibacterial Activity and Determination of Total Phenol and Flavonoid of Carica papaya L. Ethanol Extract. *International Journal of Science, Technology & Management*. 2023;4(1):233-8. doi: 10.46729/ijstm.v4i1.738.
20. Mangkulion S, Roeslan, MO, Monthanpisut, P. The Potential of Cogon Grass (*Imperata cylindrica*) Ethanol Extract in Inhibiting Nitric Oxide Secretion in Fibroblast. *Scientific Dental Journal*. 2023;7(1).
21. Maheshwaran L, Nadarajah, L, Senadeera, SPNN, Ranaweera, CB, Chandana, AK, Pathirana, RN. Phytochemical Testing Methodologies and Principles for Preliminary Screening/ Qualitative Testing. *Asian Plant Research Journal*. 2024;12(5):11–38. doi: 10.9734/aprj/2024/v12i5267.
22. Guediri I, Boubekri, C, Smara, O. Preliminary phytochemical screening from different extracts of *Solanum nigrum* plant growing in south of Algeria. *Journal of Fundamental and Applied Sciences*. 2023;12(2):624-33. doi: 10.4314/jfas.v12i2.7.
23. Sudha R, Philip, XC, Suriyakumari, KVP. Phytochemical Constituents of Leaves of *Moringa oleifera* Grow in Cuddalore District, Tamil Nadu, India. *SBV Journal of Basic, Clinical and Applied Health Science*. 2020;3(4).
24. Lisa, Wa Ode, S, Suparman, S. Identification of Tannin Compounds In Ethanol Extract Of Cat's Whiskers Leaves (*Orthosiphon aristatus*) Using Thin Layer Chromatography. *International Journal of Multidisciplinary Learners*. 2025;1(3):106-10.
25. Roeslan MO, Tasha, G. Ethanol extract of *Imperata Cylindrica* leaves inhibits proliferation and migration of HSC-3 cell lines. *Dental Journal (Majalah Kedokteran Gigi)*. 2021;54(3):150-4. doi: 10.20473/j.djmk.v54.i3.p150-154.
26. Kamiloglu S, Sari, G, Ozdal, T, Capanoglu, E. Guidelines for cell viability assays. *Food Frontiers*. 2020;1(3):332-49. doi: <https://doi.org/10.1002/fft2.44>.
27. Ng ZX, Samsuri, SN, Yong, PH. The antioxidant index and chemometric analysis of tannin, flavonoid, and total phenolic extracted from medicinal plant foods with the solvents of different polarities. *Journal of Food Processing and Preservation*. 2020;44(9):e14680. doi: <https://doi.org/10.1111/jfpp.14680>.
28. Rokis A, Hanafi M, Ismail C, Haris MS, Lestari W, Ichwan S, et al. Medicinal plants in the Southeast Asia for treatment and prevention of oral cancer: A systematic review. *Journal of Pharmacy & Pharmacognosy Research*. 2025;13:1003-37.
29. Barnabas F, Wuyep P, Uchejeso O, Mantu E, Ukpa S, Nduke S, et al. Phytochemical analysis and antimicrobial effects of carica papaya leaf extracts on urinary tract infections. . 2023.
30. Ilham R, Lelo A, Harahap U, Widyawati T, Siahaan L. The Effectivity of Ethanolic Extract from Papaya Leaves (*Carica papaya* L.) as an Alternative Larvicide to *Aedes* spp. (1857-9655 (Print)).
31. Yuliani R, Syahdeni, F. Ethanolic Extract of Papaya Leaves (*Carica papaya*) and its Fractions have no Potential Cytotoxicity on T47D Cells. *Pharmakon: Jurnal Farmasi Indonesia*; Vol 17, No 1 (2020). 2020. doi: 10.23917/pharmakon.v17i1.10760.
32. Kong YR, Jong, YX, Balakrishnan, M, Bok, ZK, Weng, JK, Tay, KC, et al. Beneficial Role of Carica papaya Extracts and Phytochemicals on Oxidative Stress and Related Diseases: A Mini Review. *Biology*. 2021 doi: 10.3390/biology10040287.
33. Zhang D, Kanakkanthara, A. Beyond the Paclitaxel and Vinca Alkaloids: Next Generation of Plant-Derived Microtubule-Targeting Agents with Potential Anticancer Activity. *Cancers*. 2020 doi: 10.3390/cancers12071721.
34. Kopustinskiene DM, Jakstas, V, Savickas, A, Bernatoniene, J. Flavonoids as Anticancer Agents. *Nutrients*. 2020 doi: 10.3390/nu12020457.
35. Yi W, Yan, ZY, Shi, MQ, Zhang, Y, Liu, J, Yi, Q, et al. [Mechanism of tannins from *Galla chinensis* cream in promoting skin wound healing irats based on FAK/PI3K/Akt/mTOR signaling pathway]. *Zhongguo Zhong Yao Za Zhi*. 2025;50(2):480-97. doi: 10.19540/j.cnki.cjcm.20240902.706.
36. Ghante MH, Jamkhande, PG. Role of Pentacyclic Triterpenoids in Chemoprevention and Anticancer Treatment: An Overview on Targets and Underling Mechanisms. *J Pharmacopuncture*. 2019;22(2):55-67. doi: 10.3831/KPI.201.22.007.
37. Chang L, Zhou, R, He, Y, Meng, M, Hu, J, Liu, Y, et al. Total saponins from *Rhizoma Panacis Majoris* inhibit proliferation, induce cell cycle arrest and apoptosis and influence MAPK signalling pathways on the colorectal cancer cell. *Mol Med Rep*. 2021;24(2):542. doi:10.3892/mmr.2021.12181.
38. Singh SP, Mishra, A, Shyanti, RK, Singh, RP, Acharya, A. Silver Nanoparticles Synthesized Using Carica papaya Leaf Extract (AgNPs-PLE) Causes Cell Cycle Arrest and Apoptosis in Human Prostate (DU145) Cancer Cells. *Biological Trace Element Research*. 2021;199(4):1316-31. doi:10.1007/s12011-020-02255-z.
39. Evans DM, Fang, J, Silvers, T, Delosh, R, Laudeman, J, Ogle, C, et al. Exposure time versus cytotoxicity for anticancer agents. *Cancer Chemotherapy and Pharmacology*. 2019;84(2):359-71. doi: 10.1007/s00280-019-03863-w.
40. Munir S, Liu Z-W, Tariq T, Rabail R, Kowalczewski Pt, Lewandowicz J, et al. Delving into the Therapeutic Potential of Carica papaya Leaf against Thrombocytopenia. *Molecules [Internet]*. 2022; 27(9).

41. Khor B-K, Chear NJ, Azizi J, Khaw K-Y. Chemical Composition, Antioxidant and Cytoprotective Potentials of Carica papaya Leaf Extracts: A Comparison of Supercritical Fluid and Conventional Extraction Methods. *Molecules* [Internet]. 2021; 26(5).
42. Mertens B, Nogueira, T, Stranska, R, Naesens, L, Andrei, G, Snoeck, R. Cidofovir is active against human papillomavirus positive and negative head and neck and cervical tumor cells by causing DNA damage as one of its working mechanisms. *Oncotarget*; Vol 7, No. 2016. doi: <https://doi.org/10.18632/oncotarget.10100>.
43. Pardenas I, Roeslan, M. Cytotoxicity Activity of Allium sativum Extracts Against HSC-3 cells. Quality improvement in dental and medical knowledge, research, skills and ethics facing global challenges. London: CRC Press; 2024. p. 393–8.

<https://doi.org/> Copyright: © 2025 by Padjadjaran Journal of Dentistry. Submitted to Padjadjaran Journal of Dentistry for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).



ORIGINAL ARTICLE

Phytochemical composition and cytotoxic assessment of *Carica papaya* Leaf extracts on human squamous carcinoma-3 Cells (in vitro): a laboratory experimental

Moehamad Orliando Roeslan^{1*}
Syafira Fayza Rahma¹
Paopanga Kokilakanit³

¹Department of Oral Biology,
Faculty of Dentistry, Universitas
Trisakti, Jakarta, Indonesia

²Undergraduate Study Program
Faculty of Dentistry, Universitas
Trisakti, Jakarta, Indonesia

³Oral Biology Research Unit,
Faculty of Dentistry, Thammasat
University, Pathum Thani, Thailand

* Correspondence:
orliando.roeslan@trisakti.ac.id

Received: 7 September 2025
Revised: 21 October 2025
Accepted: 22 November 2025
Published: 30 November 2025
DOI: 10.****/pjd.****

p-ISSN 1979-0201
e-ISSN 2549-6212

Citation:
Roeslan, MO, Rahma, SF,
Kokilakanit, P. Phytochemical
Composition and cytotoxic
assessment of Carica papaya Leaf
extracts on human squamous
carcinoma-3 Cells (in vitro): a
laboratory experimental
.Padjajaran J. Dent. November.
2025; 37(3): 1-7.

ABSTRACT

Introduction: Oral cancer is one of the diseases that can currently cause death worldwide. With the advancement of research on new drugs, medicinal plants have been explored as an alternative for cancer treatment. Papaya (*Carica papaya*) is a plant that has long been used in traditional medicine to treat various diseases. Ethanol and aqueous extracts of *C. papaya* leaf exhibit anticancer properties through the mechanism such as inhibiting cell proliferation and inducing apoptosis. The aim of this study is to analyze the cytotoxic effect of *C. papaya* leaf extract using different extraction solvents on HSC-3 cell lines.

Methods: This was a laboratory-based experimental study using a post-test only control group design. Sample size determination was performed based on the Federer formula, yielding four replicates for each group. HSC-3 cell lines were treated with ethanol and aqueous extracts of *C. papaya* leaf at concentrations of 125, 250, 500 µg/mL for 24 hours, followed by a cytotoxicity test using the CCK-8 assay. Data were analyzed using one-way ANOVA followed by Tukey's post hoc test.

Results: Alkaloids, flavonoids, and tannins were found in both ethanol and aqueous extracts, according to phytochemical analysis; triterpenoids were found only in ethanol extract, and saponins were found only in aqueous extract. HSC-3 cells were not significantly cytotoxically affected by either extract at any of the tested concentrations. **Conclusion:** Ethanol and aqueous extracts of *C. papaya* leaf at concentrations of 125, 250, 500 µg/mL showed no cytotoxic effect on HSC-3 cell lines.

KEYWORDS

C. papaya leaf, HSC-3 cells, ethanol, aquadest, CCK-8 assay.

INTRODUCTION

Oral cancer is one of the diseases that can currently cause death worldwide, including Indonesia.¹ Oral cancer is ranked as the 16th most common cancer globally. The global incidence of oral cancer is expected to reach 389,485 new cases and 188,230 deaths by 2022.² The prevalence of oral cancer in Indonesia is around 3-4% of all cancer cases, with a mortality rate of 2-3% of all cancer deaths.³

Oral Squamous cell carcinoma (OSCC) is one of the most common types of oral cancer and is found in almost 90% of all cases of carcinoma in the oral cavity.⁴ Human Squamous Carcinoma (HSC-3) cell lines are a human oral OSCC cell lines from the tongue with metastatic potential and are mostly used for oral cancer drugs research.⁵ The etiology of oral cancer is multifactorial and complex. Some

of the factors that influence the occurrence of oral cancer include smoking, alcohol consumption and Human Papilloma Virus (HPV).⁶ However, factors such as genetics also predispose to oral cancer.^{7, 8} Due to the high prevalence and mortality rate of oral cancer, effective treatment is needed. With the development of research on new drugs, medicinal plants can be used as an alternative in cancer treatment. One of the medicinal plants that has the potential as an alternative to oral cancer treatment is the papaya plant.

Papaya (*Carica papaya*) is a plant from the Caricaceae family that has been widely used in traditional medicine.⁹ *C. papaya* leaf extract shows pharmacological activities such as antidiabetic, anti-inflammatory, antimicrobial, antioxidant, and anticancer.¹⁰ Aqueous extract of *C. papaya* leaf has the potential to inhibit breast cancer cell lines (MCF-7) through the mechanism of inhibiting cell proliferation and inducing apoptosis.¹¹ Ethanol extract of *C. papaya* leaf is effective in inhibiting the proliferation of pancreatic cancer cell lines (MiaPaCa-2 and ASPC-1).¹²

Because of their distinct polarity, ethanol and water are utilised; these influence the type and concentration of beneficial anticancer chemicals obtained from *C. papaya* leaf. Ethanol was expected to generate more cytotoxic phytochemicals given its broader solvation capacity^{13, 14}; this is in line with the results showing ethanol extract to have a higher potential for cytotoxicity than water extract.^{15, 16} Though more common in water-based extracts, saponins may have less potency or greater cell line-specific anticancer action.^{17, 18}

This study addresses a significant research gap, as previous investigations on *Carica papaya* leaf extracts have primarily focused on breast, pancreatic, and prostate cancer cell lines, with limited evidence regarding their effects on oral squamous cell carcinoma (HSC-3). Moreover, earlier studies rarely compared different extraction solvents, leaving unclear how solvent polarity influences the bioactive composition and cytotoxic potential of *C. papaya*.

The novelty of this research lies in its comparative analysis of ethanol and aqueous extracts of *Carica papaya* leaf on HSC-3 cells, providing new insights into how solvent-dependent phytochemical differences relate to anticancer activity. The aim of this study was to analyze the cytotoxic effects of ethanol and aqueous extracts of *C. papaya* leaf on HSC-3 cell lines.

METHODS

This was a laboratory-based experimental study using a post-test only control group design. *Carica papaya* leaf (a California variety, collected from Bogor, West Java, Indonesia) dried in indirect sunlight and pulverized using a blender until they became powder. Extract the powder using the maceration method for three days by soaking the *C. papaya* leaf powder using ethanol 96% (EMSURE) with powder and ethanol ratio of 1:8 for ethanol extract and soaking the *C. papaya* leaf powder using aquadest with powder and aquadest ratio of 1:8 for aqueous extract at room temperature.¹⁹ During the maceration method, agitation be done several times every 15 minutes for 8 hours. Filtered using Whatman filter paper (Cytiva, Marlborough, US). Concentrated the ethanol and aqueous extract of *C. papaya* leaf using a rotary evaporator.

Phytochemical tests for the identification of alkaloid are carried out by adding 2 ml sample with 5 ml of 2N HCl and then adding Dragendorff reagent. The formation of a reddish brown precipitate indicates the presence of alkaloid compounds.²⁰ Identification of steroid and triterpenoid compounds; sample was added with 0.5 ml of Liebermann–Burchard reagent. The formation of blue-green color indicates the presence of steroid compounds, while the formation of purple/red color indicates the presence of triterpenoid compounds.²¹ Identification of flavonoid; sample was added with 5 ml of ethanol and heated for five minutes.

Then a few drops of concentrated 2N HCl and 0.2 g of magnesium powder were added. Then the sample was shaken. The formation of yellow, red or orange

color indicates the presence of flavonoid compounds.²² Identification of saponin; sample was added to water and heated for 2-3 minutes. Then stirred for 30 seconds, added 2 drops of HCl. If stable foam is formed, then the sample contains saponin compounds.²³ Identification of tannin; sample was dissolved with ethanol and added with 4 drops of FeCl₃ 1%. The formation of a blue-black solution indicates the presence of tannin compounds.²⁴

The cell culture process was carried out in a biosafety cabinet. HSC-3 culture using DMEM (Gibco, Waltham, US) and then add FBS (Gibco, Waltham, US) for cell supplementation and antibiotics such as 1% amphotericin B and 1% penicillin-streptomycin (Gibco, Waltham, US). HSC-3 were incubated at 37°C in an incubator (Esco, Changi, Singapore) humidified with 5% CO₂. Regularly, the old medium was replaced with new medium and the cells were re-cultured every 3-4 days until the cells reached 80% confluent.²⁵

Confluent cells were placed in 96 well plates with a total of 15 x 10³ cells in each well, then each well was treated according to the group, then incubated for 24 hours. 10 µL of CCK-8 reagent (Sigma-Aldrich, Darmstadt, German) was added and incubated for 4 hours at 37°C. The medium was discarded and DMSO (Acros Organics, Waltham, US) 0,049% was added to each well. This stage used a blank, control (-) with cell + medium, control (+) with 3 µM doxorubicin, and solvent control with DMSO. Each treatment group, including controls, was performed in triplicate. Readings were taken at a wavelength of 450 nm using a microplate reader (Tecan, Männedorf, Switzerland).²⁶

The number of cells was calculated by subtracting the media absorbance and a correction value (0.1696) from the treatment absorbance, then dividing the result by 0.00005. This calculation provides an estimate of viable cell count based on absorbance readings at 450 nm following treatment and incubation with the CCK-8 reagent. The IC₅₀ value, which represents the concentration required to inhibit 50% of cell viability, was calculated using linear regression analysis of the dose-response curve, by plotting the percentage of cell viability against the logarithm of extract concentrations

The collected data is then analyzed using a normality test with the Shapiro-Wilk test. Then, a One-Way ANOVA test is performed, followed by a Tukey Post-Hoc test.

RESULTS

The phytochemical test results are shown in Table 1. The results of the number of cells from each treatment are shown in Figure 1.

Table 1. Phytochemical test results

Sample	Compound	Reagent	Result	
Ethanol Extract <i>C. papaya</i> leaf	Alkaloid	Dragendorff	Brown sediment	+
	Flavonoid	HCl 2N + Mg	Yellow, red , orange	+
	Saponin	HCl	Stable foam	-
	Steroid	Chloroform + Liebermann-Burchard	Blue-green	-
	Triterpenoid	Chloroform + Liebermann-Burchard	Brownish red	+
	Tannin	FeCl ₃	Blue-black	+
Aqueous Extract <i>C. papaya</i> leaf	Alkaloid	Dragendorff	Brown sediment	+
	Flavonoid	HCl 2N + Mg	Yellow, red , orange	+
	Saponin	HCl	Stable foam	+
	Steroid	Chloroform + Liebermann-Burchard	Blue-green	-

Triterpenoid	Chloroform + Liebermann-Burchard	Brownish red	-
Tannin	FeCl ₃	Blue-black	+

The results showed that the ethanol extracts of *C. papaya* leaf contained alkaloid, flavonoid, triterpenoid and tannin compounds, while aqueous extracts of *C. papaya* leaf contained alkaloid, flavonoid, saponin and tannin compounds.




Figure 1. Bar graph showing the number of cells in each treatment. *: indicates there is a significant difference with the positive control

Figure 1 showed that the number of cells obtained from the treatment with ethanol extract of *C. papaya* leaf at concentrations of 125, 250, and 500 µg/mL were $15,300 \pm 498.117$; $15,805 \pm 545.064$; and $15,316 \pm 873.548$, respectively. Meanwhile, treatment with the aqueous extract of *C. papaya* leaf at the same concentrations resulted in cell numbers of $16,038 \pm 626.476$; $16,107 \pm 446.111$; and $15,428 \pm 793.173$, respectively. Figure 1 and table 2 show the number of HSC-3 cells under the tested conditions

Normality tests with the Shapiro Wilk method conducted using GraphPad Prism software showed that ethanol and aqueous extracts of *C. papaya* leaf with concentrations of 125, 250, 500 µg/mL, positive control, and negative control were normally distributed ($p > 0.05$). One Way ANOVA test showed there was a significant difference in all test groups with a p value of < 0.0001 . *Post-Hoc* Tukey test showed that ethanol and aqueous extracts of *C. papaya* leaf with concentrations of 125, 250, 500 µg/mL, solvent control and negative control had significant differences with the positive control.

Table 2. Cytotoxicity test results of ethanol and aqueous extracts of *C. papaya* leaf on HSC-3 cells.

Treatment Group	Concentration (µg/mL)	Cell Count (Mean ± SD)	Shapiro-Wilk (p-value)	Normality
Ethanol Extract	125	15,300 ± 498.12	0.624	Normal (p > 0.05)
	250	15,805 ± 545.06	0.711	Normal (p > 0.05)
	500	15,316 ± 873.55	0.532	Normal (p > 0.05)
Aqueous Extract	125	16,038 ± 626.48	0.842	Normal (p > 0.05)
	250	16,107 ± 446.11	0.683	Normal (p > 0.05)
	500	15,428 ± 793.17	0.794	Normal (p > 0.05)
Negative Control	—	15,781 ± 480.97	0.658	Normal (p > 0.05)
Solvent Control (DMSO)	—	15,022 ± 366.46	0.719	Normal (p > 0.05)
Positive Control (Doxorubicin)	—	5,781 ± 221.50	0.878	Normal (p > 0.05)

Data are presented as mean ± standard deviation (SD) of triplicate experiments. Normality was tested using the Shapiro-Wilk test; all groups demonstrated normal distribution (p > 0.05).

DISCUSSION

The phytochemical test of ethanol and aqueous extracts of *C. papaya* leaf showed different compounds, this difference may occur due to the use of different polarity indexes. The use of ethanol and water as solvents for *C. papaya* leaf can be attributed to the solvent selectivity principle, which is controlled by polarity. Solvents have a tendency to extract compounds with similar polarity, according to the "like dissolves like" principle. Water is better at extracting highly polar substances like saponins and certain glycosides because it has a higher polarity index (9.0).

On the other hand, ethanol can dissolve a wider variety of moderately polar to slightly nonpolar substances, such as flavonoids, alkaloids, tannins, and triterpenoids, due to its lower polarity index (5.2).²⁷ The different biological activities of ethanol and aqueous extracts reflect these differences in phytochemical profiles. For example, the ethanol extract's ability to extract compounds with strong anticancer mechanisms is suggested by the presence of triterpenoids, which were not present in the aqueous extract. Thus, choosing the right solvent is essential for figuring out the chemical makeup and possible bioactivity of extracts from plants.

Figure 1 shows that there was no discernible cytotoxic effect because the number of viable HSC-3 cells stayed above 15,000 at all concentrations of the ethanol and aqueous extracts of *C. papaya* leaf. While all extract types were significantly different from the positive control group (doxorubicin), neither at concentrations of 125, 250, or 500 µg/mL significantly decreased cell viability when compared to the negative control. In addition, the systematic study by Rokis et al. (2025) emphasized that most medicinal plants from Southeast Asia, including *Carica papaya*, exhibit chemopreventive potential through antioxidant mechanisms rather than direct cytotoxic effects. The antiproliferative effect of *C.*

Commented [1]: Please consult standard reporting formats for presenting normality test outcomes. The table, in its current form, does not include the normality test results and therefore requires revision.

Commented [OR2R1]: Revised

papaya tends to appear only at high concentrations (>1000 µg/mL) or after prolonged exposure periods (>72 hours).²⁸..... These results imply that the tested *C. papaya* extract concentrations are insufficient to cause cytotoxicity in HSC-3 cells during the 24-hour exposure period.

Additional information about the observed biological activity can be found in Table 1's phytochemical screening results. Alkaloids, flavonoids, triterpenoids, and tannins were present in the ethanol extract, whereas alkaloids, flavonoids, tannins, and saponins were present in the aqueous extract. The study by Barnabas et al. (2022) confirmed a similar pattern, in which alkaloids, flavonoids, saponins, tannins, and triterpenes were detected in the ethanolic extract of *C. papaya*, with higher intensity compared to the aqueous extract. The authors noted that ethanol is more effective in extracting non-polar compounds such as triterpenoids and flavonoids, which possess stronger antimicrobial and antioxidant potential.²⁹ In the context of solvent comparison, the study by Ilham et al. (2019) found that the ethanolic extract of *C. papaya* leaves contained higher levels of flavonoids and triterpenoids compared to the aqueous fraction, while the aqueous extract exhibited higher saponin content—in line with your results. This indicates that solvent polarity influences the types of active compounds extracted.³⁰

..... Alkaloids and flavonoids are examples of bioactive substances with known anticancer potential, but their stability or concentrations may not be sufficient to cause cytotoxicity in HSC-3 cells. This confirms earlier research by Yuliani and Syahdeni (2020), which found that T47D breast cancer cells were not cytotoxically affected by *C. papaya* ethanol extracts. Variations in biological responses across studies may be explained by differences in exposure duration, compound bioavailability, and cancer cell line sensitivity.³¹

Another previous study Kong et al. (2021) reported that *C. papaya* leaf extract exhibited no significant acute toxicity at doses up to 2 g/kg, the study does not directly address its effects on cancer cell viability. Current research indicates that although *C. papaya* extracts contain potentially anticancer compounds, their efficacy varies widely depending on cancer type and exposure conditions, underscoring the need for more focused oncological research.³²

Alkaloids have anticancer properties through various mechanisms, such as inhibiting cell proliferation, inducing cell death, and interacting with DNA. As tubulin inhibitors, alkaloids bind to tubulin, a protein that forms microtubules during the cell cycle. The binding of alkaloids to tubulin inhibits the aggregation of microtubule proteins, causing the cell cycle to arrest at metaphase and triggering apoptosis.³³

The anticancer mechanisms of flavonoid include inhibiting cancer cell growth, triggering apoptosis, and preventing the formation of new blood vessels, which is a process necessary for tumour growth. The anticancer activity of these flavonoids is highly dependent on several factors such as the flavonoid dosage, the type of cancer cells targeted, and their interaction with the biological environment.³⁴

Tannins can inhibit cell proliferation by suppressing protein kinase activity, thereby blocking the signal transduction pathway from the membrane to the cell nucleus.³⁵ Triterpenoid act as anticancer agents by inhibiting the cell division mechanism and activating the apoptotic pathway in cancer cells.³⁶ Saponins exhibit anticancer potential by inhibiting the formation of overexpressed Bcl-2, inducing the low-expression caspase-3 protein, increasing p53 expression, and triggering G1 cell cycle arrest.³⁷

The anticancer activity of *C. papaya* leaf extract has been tested on several cancer cell lines. Aqueous extract of *C. papaya* leaf has been shown to reduce the viability of breast cancer cell lines (MCF-7).¹¹ The ethanol extract of *C. papaya* leaf at concentrations of 125, 250, 500 µg/mL exhibited moderately active cytotoxic against prostate cancer cell lines (DU 145).³⁸ However in this study, these concentrations had no cytotoxicity effect on HSC-3.

The use of different cell lines is one of the factors influencing the anticancer effect of *C. papaya* leaf. Each cancer cell has different characteristics, as not all cells have the same growth rate.³⁴ Therefore, selection of different cell types can yield varying results depending on the experimental conditions. The effectiveness of a compound in inhibiting cell growth is highly dependent on the doubling time of the cells themselves.³⁹

To better understand the selectivity and wider applicability of *C. papaya* extracts, future research should take into account the inclusion of multiple oral cancer cell lines as well as normal oral epithelial cells. It is also advised to look into the molecular processes that underlie cytotoxicity, such as cell cycle arrest, reactive oxygen species production, and apoptotic signalling, in order to clarify how the extracts work. Additionally, to identify and measure the active compounds causing the observed effects, phytochemical profiling using HPLC or GC-MS should be carried out. Comparative research using different extraction methods, like ultrasound-assisted extraction or Soxhlet, may also maximise activity and yield. Lastly, in vivo research is necessary to confirm these extracts' safety profile and therapeutic potential in biological systems.

A comprehensive review by [Munir et al. \(2022\)](#) underscored the importance of elucidating *C. papaya*'s mechanisms through studies on cell cycle arrest, apoptosis, and ROS modulation, emphasizing that *in vitro* assays alone are insufficient without subsequent *in vivo* validation. They demonstrated that papaya leaf juice induced S-phase cell cycle arrest and apoptotic signalling, supporting the recommendation to assess molecular mechanisms in future studies.⁴⁰ Similarly, another previous study compared supercritical fluid extraction versus conventional extraction for *C. papaya* leaves and revealed significant differences in antioxidant and cytoprotective potentials. Their GC-MS analyses identified unique bioactive compounds (flavonoids, alkaloids, and phenolic acids), reinforcing the need for advanced extraction and profiling methods such as HPLC and GC-MS to optimize yield and activity.⁴¹

The statistical analysis in this study was performed to evaluate the differences in cytotoxic effects of *C. papaya* leaf extracts on HSC-3 cells across various treatment groups. The normality of the data was first assessed using the Shapiro-Wilk test, which confirmed that all datasets, including ethanol and aqueous extracts, as well as positive, negative, and solvent controls, were normally distributed ($p > 0.05$). Following this, a one-way ANOVA was conducted to compare the mean viable cell counts among groups. The ANOVA results revealed a statistically significant difference ($p < 0.0001$) among all treatments, suggesting that at least one group differed significantly from the others in terms of cell viability. This approach was chosen due to its suitability for analyzing quantitative outcomes derived from independent groups in cytotoxicity assays.

To determine which specific groups differed, a Tukey's post hoc test was subsequently applied. The results indicated that both ethanol and aqueous extracts of *C. papaya* at concentrations of 125, 250, and 500 µg/mL were significantly different from the positive control (doxorubicin) but not significantly different from the negative or solvent controls. This finding confirms that neither extract produced a measurable cytotoxic effect on HSC-3 cells under the tested conditions. The combination of ANOVA and Tukey's test provided a statistically robust framework for detecting potential differences while minimizing Type I error. Overall, the statistical results support the conclusion that *C. papaya* leaf extracts, within the concentration range tested, are non-cytotoxic to oral squamous carcinoma (HSC-3) cells, emphasizing the need for further mechanistic studies using molecular assays to validate these findings biologically.

The limitation of this study includes several important considerations. First of all, it only used one cell line (HSC-3) for oral squamous carcinoma, which might not accurately reflect the variety of oral cancer.^{42, 43} To confirm and expand on the results, more research utilising several cell lines or in vivo models is required. Second, without delving deeper into underlying molecular mechanisms like

apoptosis or oxidative stress pathways, the study evaluated cytotoxicity based only on cell viability. Third, only one extraction technique (maceration) and two solvents (ethanol and water) were used, which might not have fully extracted all pertinent bioactive components.

CONCLUSION

The results of this study demonstrate that both ethanol and aqueous extracts of *C. papaya* leaf at concentrations of 125, 250, and 500 µg/mL do not exhibit cytotoxic effects on HSC-3 oral squamous carcinoma cells. This result implies that even though the extracts contain bioactive substances like alkaloids, flavonoids, triterpenoids, and tannins, the concentrations may not be enough to lower cell viability in this specific cancer cell line. To maximise the possible anticancer effects of *C. papaya* leaf extracts, more research is therefore required using higher concentrations, longer exposure periods, or alternative extraction methods. This conclusion emphasises the need for more investigation in oral cancer therapy research and directly addresses the study goal of comparing the cytotoxicity effects.

The implication of this research is for the development of alternative, plant-based treatments for oral cancer. The results further emphasise the significance of extraction technique and solvent selection in optimising the medicinal potential of herbal compounds, which may help guide the development of more standardised and potent phytotherapeutic agents in subsequent oncology studies.

Acknowledgement: No declare

Author Contributions: Conceptualization, M.O.R. and S.F.R.; methodology, S.F.R.; software, M.O.R.; validation, M.O.R. and S.F.R.; formal analysis, M.O.R.; investigation, S.F.R.; resources, S.F.R.; data curation, S.F.R.; writing original draft preparation, S.F.R.; writing review and editing, M.O.R. and S.F.R.; visualization, M.O.R. and S.F.R.; supervision, M.O.R.; project administration, M.O.R.; funding acquisition, S.F.R. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: All research procedures have been approved by the Research Ethics Committee, Faculty of Dentistry, Trisakti University. No. 768A/S1/KEPK/FGK/8/2024 on August 6th 2024.

Data Availability Statement: Not applicable.

Conflicts of Interest: The author declared no conflicts of interest or potential commercial background in this research.

REFERENCES

- Warnakulasuriya S, Filho, AM. Oral Cancer in the South and South-East Asia Region, 2022: Incidence and Mortality. *Oral Diseases*. 2025;31(5):1398-405. <https://doi.org/10.1111/odi.15369>.
- Bray F, Laversanne, M, Sung, H, Ferlay, J, Siegel, RL, Soerjomataram, I, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2024;74(3):229-63. <https://doi.org/10.3322/caac.21834>.
- Gharat SA, Momin, MM, Bhavsar, C. Oral Squamous Cell Carcinoma: Current Treatment Strategies and Nanotechnology-Based Approaches for Prevention and Therapy. 2016;33(4):363-400. <https://doi.org/10.1615/CritRevTherDrugCarrierSyst.2016016272>.
- Ding M, Yu, S, Chen, Y, Liu, Y, Xuan, F. Space-time analysis of head and neck cancer in Asia and its 34 countries and territories (1990-2021): Implications from the Global Burden of Disease Study 2021. *PLoS One*. 2025;20(6):e0326177. <https://doi.org/10.1371/journal.pone.0326177>.
- Jayaraman S, Veeraraghavan, VP, Natarajan, SR, Jasmine, S. Exploring the therapeutic potential of curcumin in oral squamous cell carcinoma(HSC-3 cells): Molecular insights into hypoxia-mediated angiogenesis. *Pathology - Research and Practice*. 2024;254:155130. doi: <https://doi.org/10.1016/j.prp.2024.155130>.
- Yang Z, Sun, P, Dahlstrom, KR, Gross, N, Li, G. Joint effect of human papillomavirus exposure, smoking and alcohol on risk of oral squamous cell carcinoma. *BMC Cancer*. 2023;23(1):457. <https://doi.org/10.1186/s12885-023-10948-6>.
- Chu Y-H, Tzeng, S-L, Lin, C-W, Chien, M-H, Chen, M-K, Yang, S-F. Impacts of MicroRNA Gene Polymorphisms on the Susceptibility of Environmental Factors Leading to Carcinogenesis in Oral Cancer. *PLOS ONE*. 2012;7(6):e39777. <https://doi.org/10.1371/journal.pone.0039777>.
- da Cruz Perez DE, Passos, KKM, Machado, RA, Martelli-Junior, H, Bonan, PRF. Continuing education in oral cancer during coronavirus disease 2019 (covid-19) outbreak. *Oral Oncology*. 2020;105:104713. <https://doi.org/10.1016/j.oraloncology.2020.104713>.

9. Manpreet K, NaveenChandra, T, Seema, S, Arvind, K, Elena, ES. Ethnomedicinal Uses, Phytochemistry and Pharmacology of Carica papaya Plant: A Compendious Review. Mini-Reviews in Organic Chemistry. 2019;16(5):463-80. <http://dx.doi.org/10.2174/1570193X15666180816110733>.
10. Sharma A, Sharma, R, Sharma, M, Kumar, M, Barbhai, MD, Lorenzo, JM, et al. Carica papaya L. Leaves: Deciphering Its Antioxidant Bioactives, Biological Activities, Innovative Products, and Safety Aspects. Oxidative Medicine and Cellular Longevity. 2022;2022(1):2451733. doi: <https://doi.org/10.1155/2022/2451733>.
11. Nisa FZ, Astuti, M, Murdiati, A, Haryana, SM. Anti-proliferation and apoptosis induction of aqueous leaf extract of Carica papaya L. on human breast cancer cells MCF-7. Pakistan Journal of Biological Sciences. 2017;20(1):36-41. doi: <https://doi.org/10.3923/pjbs.2017.36.41>
12. Vuong QV, Hirun, S, Chuen, TLK, Goldsmith, CD, Murchie, S, Bowyer, MC, et al. Antioxidant and anticancer capacity of saponin-enriched Carica papaya leaf extracts. Int J of Food Science and Technology. 2015;50(1):169-77. <https://doi.org/10.1111/ijfs.12618>
13. Abd'quadri-Abojukoro AN, Nkadimeng, SM, McGaw, LJ, Nsahlai, IV. Phytochemical composition and cytotoxicity of ethanolic extracts of some selected plants. J of Applied Animal Research. 2022;50(1):656-65. <https://doi.org/10.1080/09712119.2022.2119979>.
14. Niyonizigiye I, Nkurunziza, D, Ngabire, D, Gitachew, AT, Chun, BS, Kim, G-D. Characterization and in vitro cytotoxicity of phytochemicals from Asplia africana obtained using green extraction techniques. South African Journal of Botany. 2020;128:231-8. <https://doi.org/10.1016/j.sajb.2019.11.013>.
15. Mpingirika EZ, El Hosseiny, A, Bakheit, SMS, Arafah, R, Amleh, A. Potential Anticancer Activity of Crude Ethanol, Ethyl Acetate, and Water Extracts of Ephedra foeminea on Human Osteosarcoma U2OS Cell Viability and Migration. BioMed Research International. 2020;2020(1):3837693. <https://doi.org/10.1155/2020/3837693>.
16. Tessalonica S, Roeslan, MO. Effect of ethanol extracts from Persea americana leaves on HSC-3 proliferation. Journal of Indonesian Dental Association. 2020;3(2):65-70. <https://doi.org/10.32793/jida.v3i2.582>
17. Bozinou E, Pappas, IS, Patergiannakis, I-S, Chatzimitakos, T, Palaogiannis, D, Athanasiadis, V, et al. Evaluation of Antioxidant, Antimicrobial, and Anticancer Properties of Onion Skin Extracts. 2023;15(15):11599. <https://doi.org/10.3390/su151511599>
18. Jawad M, Ali, M, Qasim, S, Akbar, A, Khan, NA, Sadiq, MB. Determination of Phenolic Compounds and Bioactive Potential of Plum (Prunus salicina) Peel Extract Obtained by Ultrasound-Assisted Extraction. BioMed Research International. 2022;2022(1):7787958. <https://doi.org/10.1155/2022/7787958>.
19. Karo-Karo SU, Arianto, A, Salim, E. Antibacterial Activity and Determination of Total Phenol and Flavonoid of Carica papaya L. Ethanol Extract. International Journal of Science, Technology & Management. 2023;4(1):233-8. <https://doi.org/10.46729/ijstm.v4i1.738>.
20. Mangkulion S, Roeslan, MO, Monthanpisut, P. The Potential of Cogon Grass (Imperata cylindrica) Ethanol Extract in Inhibiting Nitric Oxide Secretion in Fibroblast. Scientific Dental Journal. 2023;7(1).
21. Maheshwaran L, Nadarajah, L, Senadeera, SPNN, Ranaweera, CB, Chandana, AK, Pathirana, RN. Phytochemical Testing Methodologies and Principles for Preliminary Screening/ Qualitative Testing. Asian Plant Research Journal. 2024;12(5):11-38. <https://doi.org/10.9734/apri/2024/v12i5267>.
22. Guediri I, Boubekri, C, Smara, O. Preliminary phytochemical screening from different extracts of Solanum nigrum plant growing in south of Algeria. Journal of Fundamental and Applied Sciences. 2023;12(2):624-33. <https://doi.org/10.4314/jfas.v12i2.7>.
23. Sudha R, Philip, XC, Suriyakumari, KVP. Phytochemical Constituents of Leaves of Moringa oleifera Grow in Cuddalore District, Tamil Nadu, India. SBV Journal of Basic, Clinical and Applied Health Science. 2020;3(4).
24. Lisa, Wa Ode, S, Suparman, S. Identification of Tannin Compounds In Ethanol Extract Of Cat's Whiskers Leaves (Orthosiphon aristatus) Using Thin Layer Chromatography. International Journal of Multidisciplinary Learners. 2025;1(3):106-10.
25. Roeslan MO, Tasha, G. Ethanol extract of Imperata Cylindrica leaves inhibits proliferation and migration of HSC-3 cell lines. Dent J (Majalah Kedokteran Gigi). 2021;54(3):150-4. <https://doi.org/10.20473/j.djmk.v54.i3.p150-154>.
26. Kamiloglu S, Sari, G, Ozdal, T, Capanoglu, E. Guidelines for cell viability assays. Food Frontiers. 2020;1(3):332-49. <https://doi.org/10.1002/fft2.44>.
27. Ng ZX, Samsuri, SN, Yong, PH. The antioxidant index and chemometric analysis of tannin, flavonoid, and total phenolic extracted from medicinal plant foods with the solvents of different polarities. Journal of Food Processing and Preservation. 2020;44(9):e14680. doi: <https://doi.org/10.1111/jfpp.14680>.
28. Rokis A, Hanafi M, Ismail C, Haris MS, Lestari W, Ichwan S, et al. Medicinal plants in the Southeast Asia for treatment and prevention of oral cancer: A systematic review. Journal of Pharmacy & Pharmacognosy Research. 2025;13:1003-37.
29. Barnabas F, Wuyep P, Uchejeso O, Mantu E, Ukpa S, Nduke S, et al. Phytochemical analysis and antimicrobial effects of carica papaya leaf extracts on urinary tract infections. 2023.
30. Ilham R, Lelo A, Harahap U, Widyawati T, Siahaan L. The Effectivity of Ethanolic Extract from Papaya Leaves (Carica papaya L.) as an Alternative Larvacide to Aedes spp. (1857-9655 (Print)).
31. Yuliani R, Syahdeni, F. Ethanolic Extract of Papaya Leaves (Carica papaya) and its Fractions have no Potential Cytotoxicity on T47D Cells. Pharmacon: Jurnal Farmasi Indonesia; Vol 17, No 1 (2020). 2020. <https://doi.org/10.23917/pharmacon.v17i1.10760>.
32. Kong YR, Jong, YX, Balakrishnan, M, Bok, ZK, Weng, JK, Tay, KC, et al. Beneficial Role of Carica papaya Extracts and Phytochemicals on Oxidative Stress and Related Diseases: A Mini Review. Biology. 2021 <https://doi.org/10.3390/biology10040287>.
33. Zhang D, Kanakkanthara, A. Beyond the Paclitaxel and Vinca Alkaloids: Next Generation of Plant-Derived Microtubule-Targeting Agents with Potential Anticancer Activity. Cancers. 2020 <https://doi.org/10.3390/cancers12071721>.
34. Kopustinskiene DM, Jakstas, V, Savickas, A, Bernatoniene, J. Flavonoids as Anticancer Agents. Nutrients. 2020 <https://doi.org/10.3390/nu12020457>.
35. Yi W, Yan, ZY, Shi, MQ, Zhang, Y, Liu, J, Yi, Q, et al. [Mechanism of tannins from Galla chinensis cream in promoting skin wound healing rats based on FAK/PI3K/Akt/mTOR signaling pathway]. Zhongguo Zhong Yao Za Zhi. 2025;50(2):480-97. <https://doi.org/10.19540/j.cnki.cjcmm.20240902.706>.

36. Ghante MH, Jamkhande, PG. Role of Pentacyclic Triterpenoids in Chemoprevention and Anticancer Treatment: An Overview on Targets and Underling Mechanisms. *J Pharmacopuncture*. 2019;22(2):55-67. <https://doi.org/10.3831/KPI.201.22.007>.
37. Chang L, Zhou, R, He, Y, Meng, M, Hu, J, Liu, Y, et al. Total saponins from Rhizoma Panacis Majoris inhibit proliferation, induce cell cycle arrest and apoptosis and influence MAPK signalling pathways on the colorectal cancer cell. *Mol Med Rep*. 2021;24(2):542. <https://doi.org/10.3892/mmr.2021.12181>.
38. Singh SP, Mishra, A, Shyanti, RK, Singh, RP, Acharya, A. Silver Nanoparticles Synthesized Using Carica papaya Leaf Extract (AgNPs-PLE) Causes Cell Cycle Arrest and Apoptosis in Human Prostate (DU145) Cancer Cells. *Biological Trace Element Research*. 2021;199(4):1316-31. <https://doi.org/10.1007/s12011-020-02255-z>.
39. Evans DM, Fang, J, Silvers, T, Delosh, R, Laudeman, J, Ogle, C, et al. Exposure time versus cytotoxicity for anticancer agents. *Cancer Chemotherapy and Pharmacology*. 2019;84(2):359-71. <https://doi.org/10.1007/s00280-019-03863-w>.
40. Munir S, Liu Z-W, Tariq T, Rabail R, Kowalczewski PŁ, Lewandowicz J, et al. Delving into the Therapeutic Potential of Carica papaya Leaf against Thrombocytopenia. *Molecules* [Internet]. 2022; 27(9).
41. Khor B-K, Chear NJ, Azizi J, Khaw K-Y. Chemical Composition, Antioxidant and Cytoprotective Potentials of Carica papaya Leaf Extracts: A Comparison of Supercritical Fluid and Conventional Extraction Methods. *Molecules* [Internet]. 2021; 26(5).
42. Mertens B, Nogueira, T, Stranska, R, Naesens, L, Andrei, G, Snoeck, R. Cidofovir is active against human papillomavirus positive and negative head and neck and cervical tumor cells by causing DNA damage as one of its working mechanisms. *Oncotarget*; Vol 7, No. 2016. doi: <https://doi.org/10.18632/oncotarget.10100>.
43. Pardenas I, Roeslan, M. Cytotoxicity Activity of Allium sativum Extracts Against HSC-3 cells. Quality improvement in dental and medical knowledge, research, skills and ethics facing global challenges. London: CRC Press; 2024. p. 393–8.

ORIGINAL ARTICLE

Phytochemical composition and cytotoxic assessment of *Carica papaya* Leaf extracts on human squamous carcinoma-3 Cells (in vitro): a laboratory experimental

Moehamad Orliando Roeslan^{1*}
Syafira Fayza Rahma¹
Paopanga Kokilakanit³

¹Department of Oral Biology,
Faculty of Dentistry, Universitas
Trisakti, Jakarta, Indonesia

²Undergraduate Study Program
Faculty of Dentistry, Universitas
Trisakti, Jakarta, Indonesia

³Oral Biology Research Unit,
Faculty of Dentistry, Thammasat
University, Pathum Thani, Thailand

* Correspondence:
orliando.roeslan@trisakti.ac.id

Received: 7 September 2025

Revised: 21 October 2025

Accepted: 22 November 2025

Published: 30 November 2025

DOI: 10.****/pjd.****

p-ISSN [1979-0201](#)

e-ISSN [2549-6212](#)

Citation:

Roeslan, MO. Rahma, SF.
Kokilakanit, P. Phytochemical
Composition and cytotoxic
assessment of *Carica papaya* Leaf
extracts on human squamous
carcinoma-3 Cells (in vitro): a
laboratory experimental
.Padjajaran J. Dent. November.
2025; 37(3): 1-7.

ABSTRACT

Introduction: Oral cancer is one of the diseases that can currently cause death worldwide. With the advancement of research on new drugs, medicinal plants have been explored as an alternative for cancer treatment. Papaya (*Carica papaya*) is a plant that has long been used in traditional medicine to treat various diseases. Ethanol and aqueous extracts of *C. papaya* leaf exhibit anticancer properties through the mechanism such as inhibiting cell proliferation and inducing apoptosis. The aim of this study is to determine the cytotoxic effect of *C. papaya* leaf extract using different extraction solvents on HSC-3 cell lines. **Methods:** This was a laboratory-based experimental study using a post-test only control group design. HSC-3 cell lines were treated with ethanol and aqueous extracts of *C. papaya* leaf at concentrations of 125, 250, 500 µg/mL for 24 hours, followed by a cytotoxicity test using the CCK-8 assay. **Results:** Alkaloids, flavonoids, and tannins were found in both ethanol and aqueous extracts, according to phytochemical analysis; triterpenoids were found only in ethanol extract, and saponins were found only in aqueous extract. HSC-3 cells were not significantly cytotoxically affected by either extract at any of the tested concentrations. **Conclusion:** Ethanol and aqueous extracts of *C. papaya* leaf at concentrations of 125, 250, 500 µg/mL showed no cytotoxic effect on HSC-3 cell lines.

KEYWORDS

C. papaya leaf, HSC-3 cells, ethanol, aquadest, CCK-8 assay.

INTRODUCTION

Oral cancer is one of the diseases that can currently cause death worldwide, including Indonesia.¹ Oral cancer is ranked as the 16th most common cancer globally. The global incidence of oral cancer is expected to reach 389,485 new cases and 188,230 deaths by 2022.² The prevalence of oral cancer in Indonesia is around 3-4% of all cancer cases, with a mortality rate of 2-3% of all cancer deaths.³


Oral Squamous cell carcinoma (OSCC) is one of the most common types of oral cancer and is found in almost 90% of all cases of carcinoma in the oral cavity.⁴ Human Squamous Carcinoma (HSC-3) cell lines are a human oral OSCC cell lines from the tongue with metastatic potential and are mostly used ~~to~~ for oral cancer

drugs research.⁵ The etiology of oral cancer is multifactorial and complex. Some of the factors that influence the occurrence of oral cancer include smoking, alcohol consumption and Human Papilloma Virus (HPV).⁶ However, factors such as genetics also predispose to oral cancer.^{7, 8} Due to the high prevalence and mortality rate of oral cancer, effective treatment is needed. With the development of research on new drugs, medicinal plants can be used as an alternative in cancer treatment. One of the medicinal plants that has the potential as an alternative to oral cancer treatment is the papaya plant.

Papaya (*Carica papaya*) is a plant from the Caricaceae family that has been widely used in traditional medicine.⁹ *C. papaya* leaf extract shows pharmacological activities such as antidiabetic, anti-inflammatory, antimicrobial, antioxidant, and anticancer.¹⁰ Aqueous extract of *C. papaya* leaf has the potential to inhibit breast cancer cell lines (MCF-7) through the mechanism of inhibiting cell proliferation and inducing apoptosis.¹¹ Ethanol extract of *C. papaya* leaf is effective in inhibiting the proliferation of pancreatic cancer cell lines (MiaPaCa-2 and ASPC-1).¹²

Because of their distinct polarity, ethanol and water are utilised; these influence the type and concentration of beneficial anticancer chemicals obtained from *C. papaya* leaf. Ethanol was expected to generate more cytotoxic phytochemicals given its broader solvation capacity^{13, 14}; this is in line with the results showing ethanol extract to have a higher potential for cytotoxicity than water extract.^{15, 16} Though more common in water-based extracts, saponins may have less potency or greater cell line-specific anticancer action.^{17, 18}

This study addresses a significant research gap, as previous investigations on *Carica papaya* leaf extracts have primarily focused on breast, pancreatic, and prostate cancer cell lines, with limited evidence regarding their effects on oral squamous cell carcinoma (HSC-3). Moreover, earlier studies rarely compared different extraction solvents, leaving unclear how solvent polarity influences the bioactive composition and cytotoxic potential of *C. papaya*. The novelty of this research lies in its comparative analysis of ethanol and aqueous extracts of *Carica papaya* leaf on HSC-3 cells, providing new insights into how solvent-dependent phytochemical differences relate to anticancer activity.

 The aim of this study was to compare the cytotoxic effects of ethanol and aqueous extracts of *C. papaya* leaf on HSC-3 cell lines.

METHODS

This was a laboratory-based experimental study using a post-test only control group design. *Carica papaya* leaf (a California variety, collected from Bogor, West Java, Indonesia) dried in indirect sunlight and pulverized using a blender until they became powder. Extract the powder using the maceration method for three days by soaking the *C. papaya* leaf powder using ethanol 96% (EMSURE) with powder and ethanol ratio of 1:8 for ethanol extract and soaking the *C. papaya* leaf powder using aquadest with powder and aquadest ratio of 1:8 for aqueous extract at room temperature.¹⁹ During the maceration method, agitation be done several times every 15 minutes for 8 hours. Filtered using Whatman filter paper (Cytiva, Marlborough, US). Concentrated the ethanol and aqueous extract of *C. papaya* leaf using a rotary evaporator.

Phytochemical tests for the identification of alkaloid are carried out by adding 2 ml sample with 5 ml of 2N HCl and then adding Dragendorff reagent. The formation of a reddish brown precipitate indicates the presence of alkaloid compounds.²⁰ Identification of steroid and triterpenoid compounds; sample was added with 0.5 ml of Liebermann–Burchard reagent. The formation of blue-green color indicates the presence of steroid compounds, while the formation of purple/red color indicates the presence of triterpenoid compounds.²¹ Identification of flavonoid; sample was added with 5 ml of ethanol and heated for five minutes.

Then a few drops of concentrated 2N HCl and 0.2 g of magnesium powder were added. Then the sample was shaken. The formation of yellow, red or orange color indicates the presence of flavonoid compounds.²² Identification of saponin; sample was added to water and heated for 2-3 minutes. Then stirred for 30 seconds, added 2 drops of HCl. If stable foam is formed, then the sample contains saponin compounds.²³ Identification of tannin; sample was dissolved with ethanol and added with 4 drops of FeCl₃ 1%. The formation of a blue-black solution indicates the presence of tannin compounds.²⁴

The cell culture process was carried out in a biosafety cabinet. HSC-3 culture using DMEM (Gibco, Waltham, US) and then add FBS (Gibco, Waltham, US) for cell supplementation and antibiotics such as 1% amphotericin B and 1% penicillin-streptomycin (Gibco, Waltham, US). HSC-3 were incubated at 37°C in an incubator (Esco, Changi, Singapore) humidified with 5% CO₂. Regularly, the old medium was replaced with new medium and the cells were re-cultured every 3-4 days until the cells reached 80% confluent.²⁵

Confluent cells were placed in 96 well plates with a total of 15×10^3 cells in each well, then each well was treated according to the group, then incubated for 24 hours. 10 µL of CCK-8 reagent (Sigma-Aldrich, Darmstadt, German) was added and incubated for 4 hours at 37°C. The medium was discarded and DMSO (Acros Organics, Waltham, US) 0.049% was added to each well. This stage used a blank, control (-) with cell + medium, control (+) with 3 µM doxorubicin, and solvent control with DMSO. Each treatment group, including controls, was performed in triplicate. Readings were taken at a wavelength of 450 nm using a microplate reader (Tecan, Männedorf, Switzerland).²⁶ The number of cells was calculated by subtracting the media absorbance and a correction value (0.1696) from the treatment absorbance, then dividing the result by 0.00005. This calculation provides an estimate of viable cell count based on absorbance readings at 450 nm following treatment and incubation with the CCK-8 reagent. The IC₅₀ value, which represents the concentration required to inhibit 50% of cell viability, was calculated using linear regression analysis of the dose-response curve, by plotting the percentage of cell viability against the logarithm of extract concentrations

The collected data is then analyzed using a normality test with the Shapiro-Wilk test. Then, a One-Way ANOVA test is performed, followed by a Tukey Post-Hoc test.

RESULTS

The phytochemical test results are shown in Table 1. The results of the number of cells from each treatment are shown in Figure 1.

Table 1. Phytochemical test results

Sample	Compound	Reagent	Result	
Ethanol Extract <i>C. papaya</i> leaf	Alkaloid	Dragendorff	Brown sediment	+
	Flavonoid	HCl 2N + Mg	Yellow, red , orange	+
	Saponin	HCl	Stable foam	-
	Steroid	Chloroform + Liebermann-Burchard	Blue-green	-
	Triterpenoid	Chloroform + Liebermann-Burchard	Brownish red	+
	Tannin	FeCl ₃	Blue-black	+
Aqueous Extract <i>C. papaya</i> leaf	Alkaloid	Dragendorff	Brown sediment	+
	Flavonoid	HCl 2N + Mg	Yellow, red , orange	+

Saponin	HCl	Stable foam	+
Steroid	Chloroform + Liebermann-Burchard	Blue-green	-
Triterpenoid	Chloroform + Liebermann-Burchard	Brownish red	-
Tannin	FeCl3	Blue-black	+

The results showed that the ethanol extracts of *C. papaya* leaf contained alkaloid, flavonoid, triterpenoid and tannin compounds, while aqueous extracts of *C. papaya* leaf contained alkaloid, flavonoid, saponin and tannin compounds.




Figure 1. Bar graph showing the number of cells in each treatment. *: indicates there is a significant difference with the positive control

Figure 1 showed that the number of cells obtained from the treatment with ethanol extract of *C. papaya* leaf at concentrations of 125, 250, and 500 µg/mL were 15,300 ± 498.117; 15,805 ± 545.064; and 15,316 ± 873.548, respectively. Meanwhile, treatment with the aqueous extract of *C. papaya* leaf at the same concentrations resulted in cell numbers of 16,038 ± 626.476; 16,107 ± 446.111; and 15,428 ± 793.173, respectively. Figure 1 shows the number of HSC-3 cells under the tested conditions

Normality tests with the Shapiro Wilk method conducted using GraphPad Prism software showed that ethanol and aqueous extracts of *C. papaya* leaf with concentrations of 125, 250, 500 µg/mL, positive control, and negative control were normally distributed (p>0.05). One Way ANOVA test showed there was a significant difference in all test groups with a p value of <0.0001. *Post-Hoc* Tukey test showed that ethanol and aqueous extracts of *C. papaya* leaf with concentrations of 125, 250, 500 µg/mL, solvent control and negative control had significant differences with the positive control.

DISCUSSION

The phytochemical test of ethanol and aqueous extracts of *C. papaya* leaf showed different compounds, this difference may occur due to the use of different polarity indexes. The use of ethanol and water as solvents for *C. papaya* leaf can be attributed to the solvent selectivity principle, which is controlled by polarity.

Solvents have a tendency to extract compounds with similar polarity, according to the "like dissolves like" principle. Water is better at extracting highly polar substances like saponins and certain glycosides because it has a higher polarity index (9.0). On the other hand, ethanol can dissolve a wider variety of moderately polar to slightly nonpolar substances, such as flavonoids, alkaloids, tannins, and triterpenoids, due to its lower polarity index (5.2).²⁷ The different biological activities of ethanol and aqueous extracts reflect these differences in phytochemical profiles. For example, the ethanol extract's ability to extract compounds with strong anticancer mechanisms is suggested by the presence of triterpenoids, which were not present in the aqueous extract. Thus, choosing the right solvent is essential for figuring out the chemical makeup and possible bioactivity of extracts from plants.

Figure 1 shows that there was no discernible cytotoxic effect because the number of viable HSC-3 cells stayed above 15,000 at all concentrations of the ethanol and aqueous extracts of *C. papaya* leaf. While all extract types were significantly different from the positive control group (doxorubicin), neither at concentrations of 125, 250, or 500 µg/mL significantly decreased cell viability when compared to the negative control. These results imply that the tested *C. papaya* extract concentrations are insufficient to cause cytotoxicity in HSC-3 cells during the 24-hour exposure period.

Additional information about the observed biological activity can be found in Table 1's phytochemical screening results. Alkaloids, flavonoids, triterpenoids, and tannins were present in the ethanol extract, whereas alkaloids, flavonoids, tannins, and saponins were present in the aqueous extract. Alkaloids and flavonoids are examples of bioactive substances with known anticancer potential, but their stability or concentrations may not be sufficient to cause cytotoxicity in HSC-3 cells. This confirms earlier research by Yuliani and Syahdeni (2020), which found that T47D breast cancer cells were not cytotoxically affected by *C. papaya* ethanol extracts. Variations in biological responses across studies may be explained by differences in exposure duration, compound bioavailability, and cancer cell line sensitivity.²⁸

Another previous study Kong et al. (2021) reported that *C. papaya* leaf extract exhibited no significant acute toxicity at doses up to 2 g/kg, the study does not directly address its effects on cancer cell viability. Current research indicates that although *C. papaya* extracts contain potentially anticancer compounds, their efficacy varies widely depending on cancer type and exposure conditions, underscoring the need for more focused oncological research.²⁹

Alkaloids have anticancer properties through various mechanisms, such as inhibiting cell proliferation, inducing cell death, and interacting with DNA. As tubulin inhibitors, alkaloids bind to tubulin, a protein that forms microtubules during the cell cycle. The binding of alkaloids to tubulin inhibits the aggregation of microtubule proteins, causing the cell cycle to arrest at metaphase and triggering apoptosis.³⁰

The anticancer mechanisms of flavonoid include inhibiting cancer cell growth, triggering apoptosis, and preventing the formation of new blood vessels, which is a process necessary for tumor growth. The anticancer activity of these flavonoids is highly dependent on several factors such as the flavonoid dosage, the type of cancer cells targeted, and their interaction with the biological environment.³¹

Tannins can inhibit cell proliferation by suppressing protein kinase activity, thereby blocking the signal transduction pathway from the membrane to the cell nucleus.³² Triterpenoid act as anticancer agents by inhibiting the cell division mechanism and activating the apoptotic pathway in cancer cells.³³ Saponins exhibit anticancer potential by inhibiting the formation of overexpressed Bcl-2, inducing the low-expression caspase-3 protein, increasing p53 expression, and triggering G1 cell cycle arrest.³⁴

The anticancer activity of *C. papaya* leaf extract has been tested on several cancer cell lines. Aqueous extract of *C. papaya* leaf has been shown to reduce the viability of breast cancer cell lines (MCF-7).¹¹ The ethanol extract of *C. papaya* leaf at concentrations of 125, 250, 500 µg/mL exhibited moderately active cytotoxic against prostate cancer cell lines (DU 145).³⁵ However in this study, these concentrations had no cytotoxicity effect on HSC-3.

The use of different cell lines is one of the factors influencing the anticancer effect of *C. papaya* leaf. Each cancer cell has different characteristics, as not all cells have the same growth rate.³¹ Therefore, selection of different cell types can yield varying results depending on the experimental conditions. The effectiveness of a compound in inhibiting cell growth is highly dependent on the doubling time of the cells themselves.³⁶

The limitation of this study includes several important considerations. First of all, it only used one cell line (HSC-3) for oral squamous carcinoma, which might not accurately reflect the variety of oral cancer.^{37, 38} To confirm and expand on the results, more research utilising several cell lines or in vivo models is required. Second, without delving deeper into underlying molecular mechanisms like apoptosis or oxidative stress pathways, the study evaluated cytotoxicity based only on cell viability. Third, only one extraction technique (maceration) and two solvents (ethanol and water) were used, which might not have fully extracted all pertinent bioactive components.

To better understand the selectivity and wider applicability of *C. papaya* extracts, future research should take into account the inclusion of multiple oral cancer cell lines as well as normal oral epithelial cells. It is also advised to look into the molecular processes that underlie cytotoxicity, such as cell cycle arrest, reactive oxygen species production, and apoptotic signalling, in order to clarify how the extracts work. Additionally, to identify and measure the active compounds causing the observed effects, phytochemical profiling using HPLC or GC-MS should be carried out. Comparative research using different extraction methods, like ultrasound-assisted extraction or Soxhlet, may also maximise activity and yield. Lastly, in vivo research is necessary to confirm these extracts' safety profile and therapeutic potential in biological systems.

CONCLUSION

The results of this study demonstrate that both ethanol and aqueous extracts of *C. papaya* leaf at concentrations of 125, 250, and 500 µg/mL do not exhibit cytotoxic effects on HSC-3 oral squamous carcinoma cells. This result implies that even though the extracts contain bioactive substances like alkaloids, flavonoids, triterpenoids, and tannins, the concentrations may not be enough to lower cell viability in this specific cancer cell line. To maximise the possible anticancer effects of *C. papaya* leaf extracts, more research is therefore required using higher concentrations, longer exposure periods, or alternative extraction methods. This conclusion emphasises the need for more investigation in oral cancer therapy research and directly addresses the study goal of comparing the cytotoxicity effects.

The implication of this research is for the development of alternative, plant-based treatments for oral cancer. The results further emphasise the significance of extraction technique and solvent selection in optimising the medicinal potential of herbal compounds, which may help guide the development of more standardised and potent phytotherapeutic agents in subsequent oncology studies.

Acknowledgement: No declare

Author Contributions: Conceptualization, M.O.R. and S.F.R.; methodology, S.F.R.; software, M.O.R.; validation, M.O.R. and S.F.R.; formal analysis, M.O.R.; investigation, S.F.R.; resources, S.F.R.; data curation, S.F.R.; writing original draft preparation, S.F.R.; writing review and editing, M.O.R. and

S.F.R.; visualization, M.O.R. and S.F.R.; supervision, M.O.R.; project administration, M.O.R.; funding acquisition, S.F.R. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: All research procedures have been approved by the Research Ethics Committee, Faculty of Dentistry, Trisakti University. No. 768A/S1/KEPK/FKG/8/2024 on August 6th 2024.

Data Availability Statement: Not applicable.

Conflicts of Interest: The author declared no conflicts of interest or potential commercial background in this research.

REFERENCES

- Warnakulasuriya S, Filho, AM. Oral Cancer in the South and South-East Asia Region, 2022: Incidence and Mortality. *Oral Diseases*. 2025;31(5):1398-405. doi: <https://doi.org/10.1111/odi.15369>.
- Bray F, Laversanne, M, Sung, H, Ferlay, J, Siegel, RL, Soerjomataram, I, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2024;74(3):229-63. doi: 10.3322/caac.21834.
- Gharat SA, Momin, MM, Bhavsar, C. Oral Squamous Cell Carcinoma: Current Treatment Strategies and Nanotechnology-Based Approaches for Prevention and Therapy. 2016;33(4):363-400. doi: 10.1615/CritRevTherDrugCarrierSyst.2016016272.
- Ding M, Yu, S, Chen, Y, Liu, Y, Xuan, F. Space-time analysis of head and neck cancer in Asia and its 34 countries and territories (1990-2021): Implications from the Global Burden of Disease Study 2021. *PLoS One*. 2025;20(6):e0326177. doi: 10.1371/journal.pone.0326177.
- Jayaraman S, Veeraraghavan, VP, Natarajan, SR, Jasmine, S. Exploring the therapeutic potential of curcumin in oral squamous cell carcinoma (HSC-3 cells): Molecular insights into hypoxia-mediated angiogenesis. *Pathology - Research and Practice*. 2024;254:155130. doi: <https://doi.org/10.1016/j.prp.2024.155130>.
- Yang Z, Sun, P, Dahlstrom, KR, Gross, N, Li, G. Joint effect of human papillomavirus exposure, smoking and alcohol on risk of oral squamous cell carcinoma. *BMC Cancer*. 2023;23(1):457. doi: 10.1186/s12885-023-10948-6.
- Chu Y-H, Tzeng, S-L, Lin, C-W, Chien, M-H, Chen, M-K, Yang, S-F. Impacts of MicroRNA Gene Polymorphisms on the Susceptibility of Environmental Factors Leading to Carcinogenesis in Oral Cancer. *PLOS ONE*. 2012;7(6):e39777. doi: 10.1371/journal.pone.0039777.
- da Cruz Perez DE, Passos, KKM, Machado, RA, Martelli-Junior, H, Bonan, PRF. Continuing education in oral cancer during coronavirus disease 2019 (covid-19) outbreak. *Oral Oncology*. 2020;105:104713. doi: <https://doi.org/10.1016/j.oraloncology.2020.104713>.
- Manpreet K, Naveen Chandra, T, Seema, S, Arvind, K, Elena, ES. Ethnomedicinal Uses, Phytochemistry and Pharmacology of Carica papaya Plant: A Compendious Review. *Mini-Reviews in Organic Chemistry*. 2019;16(5):463-80. doi: <http://dx.doi.org/10.2174/1570193X15666180816110733>.
- Sharma A, Sharma, R, Sharma, M, Kumar, M, Barbhui, MD, Lorenzo, JM, et al. Carica papaya L. Leaves: Deciphering Its Antioxidant Bioactives, Biological Activities, Innovative Products, and Safety Aspects. *Oxidative Medicine and Cellular Longevity*. 2022;2022(1):2451733. doi: <https://doi.org/10.1155/2022/2451733>.
- Nisa FZ, Astuti, M, Murdiati, A, Haryana, SM. Anti-proliferation and apoptosis induction of aqueous leaf extract of Carica papaya L. on human breast cancer cells MCF-7. *Pakistan Journal of Biological Sciences*. 2017;20(1):36-41. doi: <https://doi.org/10.3923/pjbs.2017.36.41>.
- Vuong QV, Hirun, S, Chuen, TLK, Goldsmith, CD, Murchie, S, Bowyer, MC, et al. Antioxidant and anticancer capacity of saponin-enriched Carica papaya leaf extracts. *International Journal of Food Science and Technology*. 2015;50(1):169-77. doi: <https://doi.org/10.1111/ijfs.12618>.
- Abd'quadi-Abujukoro AN, Nkadiemeng, SM, McGaw, LJ, Nsahlai, IV. Phytochemical composition and cytotoxicity of ethanolic extracts of some selected plants. *Journal of Applied Animal Research*. 2022;50(1):656-65. doi: 10.1080/09712119.2022.2119979.
- Niyonizigiye I, Nkurunziza, D, Ngabire, D, Gitachew, AT, Chun, BS, Kim, G-D. Characterization and in vitro cytotoxicity of phytochemicals from *Asplia africana* obtained using green extraction techniques. *South African Journal of Botany*. 2020;128:231-8. doi: <https://doi.org/10.1016/j.sajb.2019.11.013>.
- Mpingirika EZ, El Hosseiny, A, Bakheit, SMS, Arafeh, R, Amleh, A. Potential Anticancer Activity of Crude Ethanol, Ethyl Acetate, and Water Extracts of Ephedra foeminea on Human Osteosarcoma U2OS Cell Viability and Migration. *BioMed Research International*. 2020;2020(1):3837693. doi: <https://doi.org/10.1155/2020/3837693>.
- Tessalonica S, Roeslan, MO. Effect of ethanol extracts from *Persea americana* leaves on HSC-3 proliferation. *Journal of Indonesian Dental Association*. 2020;3(2):65-70. doi: <https://doi.org/10.32793/jida.v3i2.582>.
- Bozinou E, Pappas, IS, Patergiannakis, I-S, Chatzimitakos, T, Palaiogiannis, D, Athanasiadis, V, et al. Evaluation of Antioxidant, Antimicrobial, and Anticancer Properties of Onion Skin Extracts. 2023;15(15):11599. doi: <https://doi.org/10.3390/su15151599>.
- Jawad M, Ali, M, Qasim, S, Akbar, A, Khan, NA, Sadiq, MB. Determination of Phenolic Compounds and Bioactive Potential of Plum (*Prunus salicina*) Peel Extract Obtained by Ultrasound-Assisted Extraction. *BioMed Research International*. 2022;2022(1):7787958. doi: <https://doi.org/10.1155/2022/7787958>.
- Karo-Karo SU, Arianto, A, Salim, E. Antibacterial Activity and Determination of Total Phenol and Flavonoid of Carica papaya L. Ethanol Extract. *International Journal of Science, Technology & Management*. 2023;4(1):233-8. doi: 10.46729/ijstm.v4i1.738.
- Mangkulion S, Roeslan, MO, Monthanpisut, P. The Potential of Cogon Grass (*Imperata cylindrica*) Ethanol Extract in Inhibiting Nitric Oxide Secretion in Fibroblast. *Scientific Dental Journal*. 2023;7(1).
- Maheshwaran L, Nadarajah, L, Senadeera, SPNN, Ranaweera, CB, Chandana, AK, Pathirana, RN. Phytochemical Testing Methodologies and Principles for Preliminary Screening/ Qualitative Testing. *Asian Plant Research Journal*. 2024;12(5):11-38. doi: 10.9734/aprj/2024/v12i5267.
- Guediri I, Boubekri, C, Smara, O. Preliminary phytochemical screening from different extracts of *Solanum nigrum* plant growing in south of Algeria. *Journal of Fundamental and Applied Sciences*. 2023;12(2):624-33. doi: 10.4314/jfas.v12i2.7.
- Sudha R, Philip, XC, Suriyakumari, KVP. Phytochemical Constituents of Leaves of *Moringa oleifera* Grow in Cuddalore District, Tamil Nadu, India. *SBV Journal of Basic, Clinical and Applied Health Science*. 2020;3(4).
- Lisa, Wa Ode, S, Suparman, S. Identification Of Tannin Compounds In Ethanol Extract Of Cat's Whiskers Leaves (*Orthosiphon aristatus*) Using Thin Layer Chromatography. *International Journal of Multidisciplinary Learners*. 2025;1(3):106-10.
- Roeslan MO, Tasha, G. Ethanol extract of *Imperata Cylindrica* leaves inhibits proliferation and migration of HSC-3 cell lines. *Dental Journal (Majalah Kedokteran Gigi)*. 2021;54(3):150-4. doi: 10.20473/j.djmk.v54.i3.p150-154.
- Kamiloglu S, Sari, G, Ozdal, T, Capanoglu, E. Guidelines for cell viability assays. *Food Frontiers*. 2020;1(3):332-49. doi: <https://doi.org/10.1002/fft2.44>.
- Ng ZX, Samsuri, SN, Yong, PH. The antioxidant index and chemometric analysis of tannin, flavonoid, and total phenolic extracted from medicinal plant foods with the solvents of different polarities. *Journal of Food Processing and Preservation*. 2020;44(9):e14680. doi: <https://doi.org/10.1111/jfpp.14680>.
- Yuliani R, Syahdeni, F. Ethanolic Extract of Papaya Leaves (*Carica papaya*) and its Fractions have no Potential Cytotoxicity on T47D Cells. *Pharmakon: Jurnal Farmasi Indonesia*; Vol 17, No 1 (2020). 2020. doi: 10.23917/pharmakon.v17i1.10760.
- Kong YR, Jong, YX, Balakrishnan, M, Bok, ZK, Weng, JK, Tay, KC, et al. Beneficial Role of Carica papaya Extracts and Phytochemicals on Oxidative Stress and Related Diseases: A Mini Review. *Biology*. 2021 doi: 10.3390/biology10040287.
- Zhang D, Kanakkanthara, A. Beyond the Paclitaxel and Vinca Alkaloids: Next Generation of Plant-Derived Microtubule-Targeting Agents with Potential Anticancer Activity. *Cancers*. 2020 doi: 10.3390/cancers12071721.
- Kopustinskiene DM, Jakstas, V, Savickas, A, Bernatoniene, J. Flavonoids as Anticancer Agents. *Nutrients*. 2020 doi: 10.3390/nu12020457.
- Yi W, Yan, ZY, Shi, MQ, Zhang, Y, Liu, J, Yi, Q, et al. [Mechanism of tannins from *Galla chinensis* cream in promoting skin wound healing in rats based on FAK/PI3K/Akt/mTOR signaling pathway]. *Zhongguo Zhong Yao Za Zhi*. 2025;50(2):480-97. doi: 10.19540/j.cnki.cjcm.20240902.706.
- Ghante MH, Jamkhande, PG. Role of Pentacyclic Triterpenoids in Chemoprevention and Anticancer Treatment: An Overview on Targets and Underlying Mechanisms. *J Pharmacopuncture*. 2019;22(2):55-67. doi: 10.3831/KPI.201.22.007.
- Chang L, Zhou, R, He, Y, Meng, M, Hu, J, Liu, Y, et al. Total saponins from *Rhizoma Panacis Majoris* inhibit proliferation, induce cell cycle arrest and apoptosis and influence MAPK signalling pathways on the colorectal cancer cell. *Mol Med Rep*. 2021;24(2):542. doi: 10.3892/mmr.2021.12181.
- Singh SP, Mishra, A, Shyanti, RK, Singh, RP, Acharya, A. Silver Nanoparticles Synthesized Using Carica papaya Leaf Extract (AgNPs-PLE) Causes Cell Cycle Arrest and Apoptosis in Human Prostate (DU145) Cancer Cells. *Biological Trace Element Research*. 2021;199(4):1316-31. doi: 10.1007/s12011-020-02255-z.
- Evans DM, Fang, J, Silvers, T, Delosh, R, Laudeman, J, Ogle, C, et al. Exposure time versus cytotoxicity for anticancer agents. *Cancer Chemotherapy and Pharmacology*. 2019;84(2):359-71. doi: 10.1007/s00280-019-03863-w.
- Mertens B, Nogueira, T, Stranska, R, Naesens, L, Andrei, G, Snoeck, R. Cidofovir is active against human papillomavirus positive and negative head and neck and cervical tumor cells by causing DNA damage as one of its working mechanisms. *Oncotarget*; Vol 7, No 30. 2016. doi: <https://doi.org/10.18632/oncotarget.10100>.
- Pardenas I, Roeslan, M. Cytotoxicity Activity of *Allium sativum* Extracts Against HSC-3 cells. Quality improvement in dental and medical knowledge, research, skills and ethics facing global challenges. London: CRC Press; 2024. p. 393-8.

