

# Dr. Ronald Irwanto Natadidjaja, SpPD, Subsp.PTI(K), FINASIM



## Formal Education

- **Universitas Indonesia**, Subspesialis / Konsultan Penyakit Tropik dan Infeksi, Lulus 2013
- **Universitas Indonesia**, Spesialis Penyakit Dalam (Internist), Lulus 2009
- **Universitas Trisakti**, Dokter Umum, Lulus 2002
- **SMP-SMA Kolese Kanisius**, Jakarta, Lulus 1994

## Organization

- **Tim Covid-19**, RSPI Puri Indah, 2020 – sekarang
- **Bendahara**, Perhimpunan Ilmu Kedokteran Tropis dan Penyakit Infeksi Indonesia (PETRI) Jakarta, sejak 2016 - 2023
- **Sekretaris Jenderal (Sekjen)**, Pengurus Pusat Perhimpunan Pengendalian Infeksi Indonesia (PERDALIN), sejak 2016 - 2022
- **Tim Ahli** Pokja Pencegahan dan Pengendalian Infeksi (PPI), Kemenkes RI, sejak 2017-2024
- **Kepala Bagian** Ilmu Penyakit Dalam Fakultas Kedokteran Universitas Trisakti, 2013-2020
- **Pendiri dan Perintis** RASPRO Indonesia Study Group, **Yayasan Pelita RASPRO Indonesia** untuk studi resistensi antimikroba dan penggunaan antimikroba bijak Indonesia
- **Ketua PPI** RSPI Bintaro Jaya
- **Internist-Konsultan**, RSPI Puri Indah, RSPI Bintaro Jaya, dan Tzu Chi Hospital – Pantai Indah Kapuk, Jakarta Utara



@rasproindonesia

Instagram

[www.new.rasproindonesia.com](http://www.new.rasproindonesia.com)

PERTEMUAN ILMIAH TAHUNAN  
PAPDI BANTEN 13<sup>TH</sup>



# Membangun PPRA / PGA di Rumah Sakit

**dr. Ronald Irwanto Natadidjaja, SpPD, Subsp.PTI(K), FINASIM**

**Faculty of Medicine Universitas Trisakti**

**Trisakti – RASPRO Indonesia Antimicrobial Stewardship  
(TRIASE) Learning Centre**



UNIVERSITAS TRISAKTI

Adapting to The Evolving Landscape of Internal  
Medicine: A Focus on Indonesian Practice

# Agenda


- I. Latar belakang : Mikrobiologi & Tekanan Seleksi
- II. Dari Regulasi Makro menuju Ekosistem
  - a. Regulasi Makro : Evolusi
  - b. Regulasi Mikro : Kesulitan dan Kebutuhan
  - c. Sistem : eRASPRO Manual menuju Digital
  - d. Ekosistem : Dampak Timbal Balik menuju Desain Mikro
- III. Infrastruktur – Implementasi – Evaluasi Pelaksanaan PPRA / PGA



# I. Mikrobiologi & Tekanan Seleksi

---



Mechanism of Antimicrobial Resistance:  
“Selective Pressure” for Antimicrobial-Resistant Strains



Campaign to Prevent Antimicrobial Resistance in Healthcare Settings, CDC 2002

# Fungsi Kultur

- **Diagnosis Infeksi Individu**
- **Diagnosis Komunitas :**
  - **Surveillance**
  - **Pertimbangan pembuatan panduan antimikroba empirik**



# Community

Pola Kepekaan dan Resistensi Mikroorganisme  
Aerob pada  
Infeksi Jaringan Lunak Komplikata dengan  
Berbagai Manifestasi Klinisnya  
di Tiga IGD Rumah Sakit di Jakarta

## GRAM Positive

OXA Sensitive *S. aureus* : **95.5%**

## GRAM NEGATIVE

*Pseudomonas* sp Sensitive to

MEM : **92.3%**

IMP : **92.3%**

TZP : **92.3%**

LVX : **69.2%**

AMK : **84.6%**

Ronald Irwanto ,Suhendro, Khie Chen,  
Yeva Rosana, 2009

# Hospital

## UNIVERSA MEDICINA

*January-April, 2013*

*Vol.32 - No.1*

**Culture-and nonculture-based antibiotics for  
complicated soft tissue infections are comparable**

Ronald Irwanto<sup>\*,\*\*</sup>, Suhendro<sup>\*\*</sup>, Khie Chen<sup>\*\*</sup>, and Murdani Abdullah<sup>\*\*\*</sup>

## GRAM Positive

OXA Sensitive *S. aureus* : **84.6 %**

## GRAM NEGATIVE

*Pseudomonas* sp Sensitive to

MEM : **68.2%**

IMP : **78.7%**

TZP : **50.0%**

LVX : **54.5%**

AMK : **68.2%**

Ronald Irwanto ,Suhendro, Khie Chen,  
et al . Universa Medicina 2013



**@rasproindonesia**

[www.new.rasproindonesia.com](http://www.new.rasproindonesia.com)





## Laporan Peningkatan Mutu

**PENINGKATAN MUTU PENGGUNAAN ANTIBIOTIK BIJAK MELALUI KESESUAIAN TEMUAN HASIL KULTUR DENGAN KAJIAN RISIKO PASIEN MENURUT MODEL REGULASI ANTIMIKROBA SISTEM PROSPEKTIF (RASPRO)**RONALD IRWANTO NATADIDJAJA<sup>1,2</sup>, HADIANTI ADLANI<sup>2</sup>, HADI SUMARSONO<sup>2,3</sup><sup>1</sup>Departemen Ilmu Penyakit Dalam FK TRISAKTI, Jakarta<sup>2</sup>RASPRO Indonesia Study Group<sup>3</sup>Ikatan Apoteker Indonesia

Tabel 3. Kesesuaian Temuan Hasil Kultur dengan Kajian Risiko Pasien Menurut Model RASPRO

	Multisensitif		MDR				Prediksi		
	n	%	ESBL		Non ESBL		Sesuai	Tidak Sesuai	
			n	%	n	%			
<b>Gram Negatif</b>									
Acinetobacter sp.	0	0,00	0	0,00	4	10,00	4	0	
Pseudomonas sp.	0	0,00	0	0,00	7	17,50	7	0	
Klebsiela pneumonia	15	26,32	2	22,22	6	15,00	21	2	
Eschecheria coli	18	31,58	7	77,78	6	15,00	28	3	
Citrobacter koseri	0	0,00	0	0,00	1	2,50	1	0	
Enterobacter sp.	1	1,75	0	0,00	1	2,50	2	0	
Proteus sp.	0	0,00	0	0,00	2	5,00	2	0	
Providencia stuartii	0	0,00	0	0,00	1	2,50	1	0	
Pantoea agglomerans	1	1,75	0	0,00	0	0,00	1	0	
Raoultella ornithinolytica	0	0,00	0	0,00	1	2,50	1	0	
Serratia fonticola	1	1,75	0	0,00	0	0,00	1	0	
<b>Total</b>	<b>36</b>	<b>63,15</b>	<b>9</b>	<b>100,00</b>	<b>29</b>	<b>72,50</b>	<b>69</b>	<b>5</b>	
<b>Gram Positif</b>									
Staphylococcus aureus	4	7,02	0	0,00	1	*	2,50	5	0
Staphylococcus epidermidis	1	1,75	0	0,00	2	**	5,00	3	0
Enterococcus faecalis	4	7,02	0	0,00	2		5,00	5	1
Enterococcus faecium	1	1,75	0	0,00	1		2,50	1	1
Streptococcus sp.	8	14,04	0	0,00	4		10,00	12	0
Staphylococcus sp.	3	5,26	0	0,00	1		2,50	3	1
<b>Total</b>	<b>21</b>	<b>36,84</b>	<b>0</b>	<b>0,00</b>	<b>11</b>		<b>27,50</b>	<b>29</b>	<b>3</b>
<b>TOTAL</b>	<b>57</b>	<b>100,00</b>	<b>9</b>	<b>100,00</b>	<b>40</b>	<b>100,00</b>	<b>98</b>	<b>8</b>	

\* MRSA \*\* MRSE

Tabel 4. Persentase Kesesuaian Hasil Kultur dengan Kajian Risiko Infeksi Multisensitif dan MDR Model RASPRO

	Sesuai		Tidak Sesuai		Total	
	n	%	n	%	n	%
<b>Multisensitif</b>	54	94,74	3	5,26	57	100,00
<b>MDR</b>	44	89,80	5	10,20	49	100,00

## II. Dari Regulasi Makro menuju Ekosistem

---



**Desain  
Mikro**

**Regulasi MAKRO**

Disposisi

**Regulasi MIKRO**

Integrasi

**SISTEM**

Terkondisi

**EKOSISTEM**

Permenkes 8/2015

Permenkes 28/2021

Buku Panduan PGA di RS Kemenkes RI 2021

KMK No HK.01.07/MENKES/1128/2022

To KMK No. HK 01.07 / MENKES / 1596 /2024

**GLOBAL - BIROKRATIF**

SOP – Job Desk Infrastruktur2 Birokratif

Alur Kerja PGA

Alur Pengambilan Sampel

Alur Pereseapan Antimikroba

Alur Pre-otorisasi & Audit Prospektif

dll

**EKSKLUSIF - OPERASIONAL**

*Integrated Guiding - Monitoring – Evaluating –  
Reporting – Action Plan*

**KOMITMEN - KONTINUITAS**

Kondisi timbal balik yang telah terjadi antara  
kebiasaan penggunaan antimikroba bijak  
dengan turunnya risiko kemunculan MDR

# II. Dari Regulasi Makro menuju Ekosistem

a. Regulasi Makro : Evolusi

---

# Regulasi MAKRO

GLOBAL - BIROKRATIF

Permenkes 8/2015

Permenkes 28/2021

Buku Panduan PGA di RS Kemenkes RI 2021

KMK No HK.01.07/MENKES/1128/2022

## Evolusi

Permenkes 8 / 2015

Buku Panduan PGA  
di RS Kemenkes RI 2021

Audit Retrospektif



Audit Prospektif



KEPUTUSAN MENTERI KESEHATAN REPUBLIK INDONESIA  
NOMOR HK.01.07/MENKES/1128/2022  
TENTANG  
STANDAR AKREDITASI RUMAH SAKIT

a. **Pengelolaan Kegiatan Peningkatan Mutu, Keselamatan Pasien, dan Manajemen Risiko**

1) **Standar PMKP 1**

Rumah sakit mempunyai Komite/Tim Penyelenggara Mutu yang kompeten untuk mengelola kegiatan Peningkatan Mutu dan Keselamatan Pasien (PMKP) sesuai dengan peraturan perundang-undangan.

2) **Maksud dan Tujuan PMKP 1**


Peningkatan mutu dan keselamatan pasien merupakan proses kegiatan yang berkesinambungan (*continuous improvement*) yang dilaksanakan dengan koordinasi dan integrasi antara unit pelayanan dan komite-komite (Komite Medik, Komite Keperawatan, Komite/Tim PPI, Komite K3 dan fasilitas, Komite Etik, Komite PPRA, dan lain-lainnya). Oleh karena itu Direktur perlu menetapkan Komite/Tim Penyelenggara Mutu yang bertugas membantu Direktur atau Kepala Rumah Sakit dalam mengelola kegiatan peningkatan mutu, keselamatan pasien, dan manajemen risiko di rumah sakit.

Dalam melaksanakan tugasnya, Komite/ Tim Penyelenggara Mutu memiliki fungsi sesuai dengan peraturan perundang-undangan yang berlaku.


# PANDUAN PENATAGUNAAN ANTIMIKROBA DI RUMAH SAKIT

Edisi I

Direktorat Pelayanan Kesehatan Rujukan  
Direktorat Jenderal Pelayanan Kesehatan  
Kementerian Kesehatan RI  
2021



# PPRA





# II. Dari Regulasi Makro menuju Ekosistem

b. Regulasi Mikro : Kesulitan dan  
kebutuhan

---

# Regulasi MIKRO

EKSKLUSIF - OPERASIONAL

**SOP – Job Desk Infrastruktur2 Birokratif**

**Alur Kerja PGA**

**Alur Pengambilan Sampel**

**Alur Pereseapan Antimikroba**

**Alur Pre-otorisasi & Audit Prospektif**

**dll**

**PPK**

**PPAB**

It is sometimes difficult to draw a direct relationship between system interventions and their effects. In the hospital sector, many of the studies of the efficacy of AMS have reported on structural and process measures (such as the presence of guidelines and reduction in antimicrobial use)

McGowan JE. Antimicrobial stewardship: the state of the art in 2011 – focus on outcome and methods. *Infect Control Hosp Epidemiol* 2012;33(4):331–7. 7.

MacDougall C, Polk R. Antimicrobial stewardship programs in health care systems. *Clin Microbiol Rev* 2005;18(4):638–56.



## Artikel Penelitian

# Survei Persepsi Kebutuhan dan Hambatan Rumah Sakit dalam Menjalankan Fungsi Panitia Pengendalian Resistensi Antibiotik

RONALD IRWANTO<sup>1,2</sup>, DJOKO WIDODO<sup>2</sup>, AZIZA ARIYANI<sup>3</sup>, HADIANTI ADLANI<sup>2</sup>

<sup>1</sup> Fakultas Kedokteran, Universitas Trisakti, Jakarta

<sup>2</sup> Perhimpunan Kedokteran Tropis dan Penyakit Infeksi Indonesia

<sup>3</sup> Pengurus Pusat Perkumpulan Pengendalian Infeksi Indonesia

**Hasil:** Pada survei ini diperoleh 26.92% dari 156 rumah sakit yang telah menjalankan program PPRA di rumah sakit. 65.38% menyatakan hanya sebagian dokter yang duduk sebagai anggota PPRA mampu melakukan tugasnya. 40.48% dari responden rumah sakit yang telah menjalankan program PPRA mengatakan bahwa tidak adanya sistem implementasi merupakan kesulitan utama dalam menjalankan program PPRA. Sementara 61.90% mengatakan anggota PPRA rumah sakitnya baru setengah mampu melakukan restriksi antibiotik. 93.86% dari 114 responden rumah sakit yang belum menjalankan program PPRA menyatakan saat ini yang paling dibutuhkan adalah konsep yang jelas untuk menjalankan program PPRA.

# II. Dari Regulasi Makro menuju Ekosistem

a. Sistem : eRASPRO Manual  
menuju Digital

---

# SISTEM

KOMITMEN - KONTINUITAS

*Integrated Guiding - Monitoring –  
Evaluating – Reporting – Action  
Plan*

# Futuristic Fashion in Antimicrobial Used - The WHO “Kick of” in 2023

- Shifting WATCH to  $\geq 60\%$  ACCESS



RASPRO Indonesia

[www.new.rasproindonesia.com](http://www.new.rasproindonesia.com)

Aztrenonam  
Ceftazidime Avibactam  
Ceftaroline Fosamil  
Ceftolozane Tazobactam

Imipenem cilastatin-  
relebactam

Fosfomycin IV  
Colistin  
Polymixin B  
Tygecycline

RESERVED

This group includes antibiotics and antibiotic classes that **should be reserved** for treatment of confirmed or suspected infections due to multi-drug-resistant organisms. Reserve group antibiotics should be treated as “last resort” options.

Quinolones  
Azithromycin

2<sup>nd</sup>, 3<sup>rd</sup> & 4<sup>th</sup> Generation  
of Cephalosporin

Piperacillin Tazobactam  
Carbapenems

WATCH

This group includes antibiotic classes that have higher resistance potential and includes most of the highest priority agents among the Critically Important Antimicrobials for Human Medicine and/or antibiotics that are at relatively high risk of selection of bacterial resistance. These medicines should be prioritized as key targets of stewardship programs and monitoring. Selected Watch group antibiotics are recommended as essential first or second choice empiric treatment options for a limited number of specific infectious syndromes and are listed as individual medicines on the WHO Model Lists of Essential Medicines.

Ampicillin Sulbactam  
Ampicillin  
Amoxicillin Clavulanate  
Amoxicillin

1<sup>st</sup> Generation of  
Cephalosporin

Amikacin  
Gentamycin

ACCESS

This group includes antibiotics that have activity against a wide range of commonly encountered susceptible pathogens while also showing lower resistance potential than antibiotics in the other groups. Selected Access group antibiotics are recommended as essential first or second choice empiric treatment options for infectious syndromes reviewed by the EML Expert Committee and are listed as individual medicines on the Model Lists of Essential Medicines to improve access and promote appropriate use.

AWARE 2021




USAID MEDICINES, TECHNOLOGIES, AND  
PHARMACEUTICAL SERVICES (MTaPS) PROGRAM  
*Improved Access. Improved Services. Better Health Outcomes.*



## A Technical Guide to Implementing the World Health Organization's AWaRe Antibiotic Classification in MTA PS Program Countries

**Goals of AWaRe Categorization:** The overall goal is to reduce the use of antibiotics in the Watch and Reserve groups (the antibiotics most crucial for human medicine and at higher risk of resistance) and to increase the use of Access antibiotics where availability is low. The first goal of AWaRe is to have all countries report antibiotic use, through the Antimicrobial Resistance Surveillance System (GLASS), by 2023, and the second is for 60% of global antibiotic consumption to come from medicines in the Access category.<sup>7</sup> Currently, 65 countries track antibiotic use but only 29 meet the 60% Access national consumption goal.<sup>8</sup> Evidence shows that meeting the 60% goal will result in not only better use of antibiotics but also reduced costs and increased access. Reaching this threshold by 2023 will contribute to countries' achievement of the health-related Sustainable Development Goals.





### Risk Stratification Type 3

Severe /HAI/ Febrile Neutropenia /  
Threatening Organ Perforation  
AND / OR  
Immunocompromized AND / OR  
Uncontrolled DM :  
+  
History of antibiotic use in the last 30 days  
AND / OR  
History of ≥ 48 hours hospitalization in  
the last 30 days  
  
AND / OR  
History medical devices use in the last 30  
days

**Empiric Antibiotic for Severe Case or  
Suspected ESBLs or Other MDRO**

**RESERVE RESERVE WATCH WATCH**

### Risk Stratification Type 2

Non Severe / Non Life Threatening  
– Non HAIs  
Immunocompromized  
AND / OR Uncontrolled DM :  
History of antibiotic use in the last 90 days  
AND / OR  
History of ≥ 48 hours hospitalization in  
the last 90 days  
  
AND / OR  
History medical devices use in the last 90  
days

**Empiric Antibiotic for Suspected (Beta  
Lactamase Producers) to ESBLs**

**WATCH WATCH WATCH**

### Risk Stratification Type 1

Non Risk Stratification Type 3 and / or 2



**Empiric Antibiotic for Multi-Sensitive  
Organism**

**ACCESS ACCESS ACCESS ACCESS ACCESS**



# RASPRO Model on AWARE Categories Hospital Setting

Digital Mode

Patient with  
bacterial  
infection /  
preoperative

Empiric

Definitive

Prophylaxis

De-escalation

Empiric  
Step Up  
Step Down

Guidelines  
Strat Risk Type I  
Strat Risk Type II  
Strat Risk Type II



ACCESS

WATCH

RESERVE

ACCESS

WATCH

RESERVE

ACCESS

If there is a  
special case,  
outside regulation

FREE by Indication

Supervision – Restricted  
by Indication  
PGA team agreement

FREE by Indication

Supervision – Restricted  
by Indication  
PGA team agreement

Supervision  
PGA team agreement

Automatic STOP  
Order if not  
reasonable



Integrated  
Assessment  
(FORKIT)

# e-RASPRO Digital Antimicrobial Stewardship Implementation

## INDONESIA

### HERMINA HOSPITAL GROUP - INDONESIA

#### 27 User Hospitals with Centralized Monitoring in Jakarta

RS. Hermina Jatinegara  
RS. Hermina Grand Wisata  
RS. Hermina Bogor  
RS. Hermina Galaxy  
RS. Hermina Bekasi  
RS. Hermina Daan Mogot  
RS. Hermina Ciputat  
RS. Hermina Kemayoran  
RS. Hermina Tangerang  
RS. Hermina Ciruas  
RS. Hermina Serpong  
RS. Hermina Depok  
RS. Hermina Banyumanik  
RS. Hermina Pandanaran  
RS. Hermina Makassar  
RS. Hermina Ubaya

RS. Hermina Jogjakarta  
RS. Hermina Solo  
RS. Hermina Tangkuban Perahu  
RS. Hermina Padang  
RS. Hermina Pasteur  
RS. Hermina Arcamanik  
RS. Hermina Sukabumi  
RS. Hermina Karawang  
RS Hermina Bitung  
RS Hermina Purwokerto  
RS Hermina Bitung

RS Tugu Ibu  
(e-RASPRO Beta Version mode)

RS Mardi Rahayu

RS. Marzoeki Mahdi

RSUD Cempaka Putih

# II. Dari Regulasi Makro menuju Ekosistem


a. EkoSistem : Dampak Timbal  
Balik menuju Desain Mikro

---

# I. Infrastruktur – Implementasi - Evaluasi Pelaksanaan PPRA / PGA

---

## Three Months Comparison of Broad Antibiotics Unit Sold: Before and After RASPRO-RASAL Criteria Implemented



Ronald Irwanto Natadidjaja\*#, Yuhana Fitra\*\*, Yudianto Budi Saroyo\*\*,  
Augustine Matatula\*\*, Rinna Wamila Sundariningrum

(MANUAL Model)



ORIGINAL ARTICLE

Antibiotic usage at a private hospital in Central Java: results of implementing the Indonesian Regulation on the Prospective Antimicrobial System (Regulasi Antimikroba Sistem Prospektif Indonesia [RASPRO])

Ronald Irwanto Natadidjaja<sup>1,2\*</sup>, Tarcisius Henry<sup>1</sup>, Hadianti Adlani<sup>1</sup>, Aziza Ariyani<sup>1</sup> and Rika Bur<sup>1</sup>

<sup>1</sup>RASPRO Indonesia Study Group, Jakarta, Indonesia; <sup>2</sup>Infectious Disease Division, Trisakti School of Medicine, Trisakti University, Jakarta, Indonesia

*Abstract*

---

**Methods:** A pre–post-descriptive study was conducted in 2019 for 3 months at a private hospital in Central Java, Indonesia, to evaluate the implementation of the Regulation on Indonesian Antimicrobial Stewardship Program (ASP), namely, the Prospective Antimicrobial System/Regulasi Antimikroba Sistem Prospektif Indonesia (RASPRO). Outcomes were measured before and after the implementation of the RASPRO in the ward including: 1) intravenous antibiotic defined daily dose (DDD) per 100 patient-days, 2) antibiotic expenditure, and 3) antibiotic expenditure per inpatient.

**Result:** The total antibiotic consumption was expressed in DDD/100 patient-days. For the levofloxacin category, the number increased intensely from 2.38 to 15.29; carbapenem escalated from 0.51 to 2.31, ceftriaxone from 32.10 to 38.03, and ampicillin sulbactam from 1.14 to 1.18. In contrast, cefuroxime significantly reduced from 17.25 to 1.38, cefotaxime decreased from 10.33 to 6.83, gentamicin decreased from 3.18 to 1.91, and amikacin decreased from 2.27 to 2.13. The overall cephalosporin usage decreased from 19.89 to 15.41. The total antibiotic expenditure had a decline of 20.28%, followed by 14.44% reduction on the percentage of antibiotic expenditure per inpatient.

**Conclusion:** Our study describes the 3-month analysis of antimicrobial usage before and after the implementation of the RASPRO by evaluating several parameters. The antibiotic consumption expressed in DDD/100 patient-days for each antibiotic category has demonstrated that there are different impacts that may be debatable and calls for further evaluation. A decrease in the total antibiotic expenditure has also been reported. However, since our study is a preliminary study, it should be continued by further studies that involve longer study duration to observe further impacts of the program.

MEETING ABSTRACTS

Open Access



# International Conference on Prevention and Infection Control 2023

## A quantitative survey of antibiotic use at a hospital in Jambi Province Indonesia in three-month before and after implementation of antimicrobial resistance control program by Raspro concept

R. I. Natadidjaja<sup>1,2,\*</sup>, R. Asmajaya<sup>2</sup>, H. Basrie<sup>2</sup>, H. Sumarsono<sup>2</sup>

<sup>1</sup>Internal Medicine, Faculty of Medicine, Universitas Trisakti, <sup>2</sup>Pelita RASPRO Indonesia Foundation, Jakarta Barat, Indonesia

**Correspondence:** R. I. Natadidjaja

*Antimicrobial Resistance & Infection Control* 2023, **12(Suppl 1):P309**

**Introduction:** Based on Decree of Minister of Health Number 8/2015 in article 11 concerning quality indicators of Antimicrobial Resistance Control Program (ARCP)/Program Pengendalian Resistensi Antimikroba (PPRA) implementation in hospitals, it has been known that reduced quantity of antimicrobial use has become one of those indicators.

**Objectives:** This survey is a descriptive study using secondary data retrieved between July and September 2019 (3 months before implementation of RASPRO concept) as well as between October and December 2019 (3 months after the implementation), which was aimed to evaluate impacts on implementing *Regulasi Antimikroba Sistem Prospektif (RASPRO)* concept at a hospital in Jambi province, Indonesia.

**Methods:** The survey was carried out by calculating the expenditure of 3 antibiotic classes, which were the most commonly used and usually given by injection in hospitals and Intensive Care Units (ICU)s, i.e. the beta-lactam, quinolones and carbapenem.

**Results:** We found reduced use of Ceftriaxone as many as 890 ampules (37.11%), for Cefotaxime the reduction was 580 ampules (67.13%); while the use of Cefoperazone reduced as many as 76 ampules (47.50%) and Ceftazidime reduced as many as 10 ampules (7.14%). The use of Ciprofloxacin reduced as many as 327 ampules (71.40%), but there was a drastic increase in the use of Levofloxacin as many as 59 ampules (>100%). The use of Carbapenems increased, which included 79 ampules (34.20%) for Meropenem; while the use of Imipenem increased as many as 9 ampules (100%). In three months after the implementation of RASPRO concept, 92.5% prophylaxis antibiotic had been given for appropriate indication and the antibiotic use of Cefazolin 71.3%. Within three months before and after the implementation of RASPRO concept, there was a total reduction of antibiotic use, which reached 1736 ampules (40.57%).

**Conclusion:** In conclusion, the implementation of RASPRO concept can be executed as an effort to reduce the quantity of antimicrobial use in hospitals. However, larger studies and longer monitoring are required in order to identify the impact of implementation of RASPRO concepts at a hospital.

**Disclosure of Interest**  
None declared.

(MANUAL Model)

# Qualitative Evaluation of Antibiotic with Gyssens Method by RASPRO Concept for Pneumonia at Pediatric Intensive Care Unit

Rinna W. Sundariningrum,<sup>1</sup> Darmawan Budi Setyanto,<sup>2</sup> Ronald Irwanto Natadidjaja<sup>3</sup>

**Background.** Pneumonia remains the commonest infective reason for admission to intensive care as well as being the most common secondary infection acquired whilst in the pediatric intensive care unit. Inappropriate use of antibiotics can increase morbidity, mortality, patient cost, and antibiotic resistance.

**Objective.** To qualitatively evaluate antibiotic use in pneumonia with The Gyssens method by RASPRO concept.


**Methods.** We performed a descriptive, retrospective study data based on medical records of patients with pneumonia who admitted to the pediatric intensive care unit in Hermina Bekasi Hospital from May to October 2019. Records were evaluation its qualitative antibiotic using the Gyssens method by RASPRO concept.

**Result.** This study discovered 51 cases (14,46%) of severe pneumonia. We found 119 antibiotics uses including 90 (75,63%) empirical therapies and 29 (24,37%) devinitive therapies. Ampicilin sulbactam was the most common antibiotic used (15,98%), followed by cefotaxime (15,12%), meropenem (13,44%), azithromycin (11,78%) and ceftriaxone (10,92%). Based on Gyssens method by RASPRO concept, appropriate antibiotic use (category 0) accounted for 63,02%, while inappropriated use accounted for 1,68% category IVa (improper; other antibiotics were more effective), 22,69% category IIIa (improper; duration too long), 9,24% category IIIb (improper; duration too short) and 3,36% category IIa (improper; incorrect dose).

**Conclusion.** Appropriate use of antibiotics showed quite good results, namely 63,03%. The RASPRO concept can be used to reduce subjectivity bias in qualitative antibiotic assessments by the Gyssens method for pneumonia treated in the pediatric intensive care unit. **Sari Pediatri** 2020;22(2):109-14

## 9 months before & after using digital ASP model

43% decline of Inpatient Antibiotic Usage



Dr. Iin Indra Pertiwi SpPD

RASPRO Indonesia - Indonesian Grass Root Meeting on Antimicrobial Stewardship (INDOGRAM)  
World Antimicrobial Awareness Week, November 2022

To do further research in 3 hospitals , In progress publication

(Digital Model)



## Trend Changing to the ACCESS Category Antibiotic Usage after Digital Antimicrobial Stewardship Tool e-RASPRO 9 Months Implementation in an Indonesian Hospital

Hadianti Adlani<sup>1,2</sup>, Aziza Ariyani<sup>2</sup>, Ronald Irwanto Natadidjaja<sup>1,2,3</sup>, Anti Dharmayanti<sup>1,2</sup>

<sup>1</sup> Indonesian Society of Infection Control (INASIC) Branch Banten

<sup>2</sup> RASPRO Indonesia *Study Group*

<sup>3</sup> Faculty of Medicine Universitas Trisakti

**Background:** Antimicrobial Stewardship Program (ASP) is a global issue. World Health Organization (WHO) stated, there are 3 categories of antimicrobial: ACCESS, WATCH, and RESERVE. e-RASPRO as a digital ASP may alter antibiotic prescribing pattern by prioritizing ACCESS category as suggested by WHO.

**Methods:** This manuscript was a ward retrospective survey data of 9 months Define Daily Dose (DDD) average before-after implementing the electronic-RASPRO (e-RASPRO) on ACCESS & WATCH antibiotic.

**Results:** Number of inpatients 9 months before-after e-RASPRO implementation were 7,754 and 6,794. Within 9 months after implementing e-RASPRO there was a trend of antibiotic prescription shifting from WATCH category antibiotic to ACCESS category antibiotic. There was a trend of reduced Define Daily Dose (DDD) average of WATCH category antibiotic. 24.82% of 3<sup>rd</sup> generation Cephalosporin, 33.20% of Quinolones, 14.76% of Carbapenems and 100% of Piperacillin Tazobactam DDD average were reduced. While, in ACCESS Category Antibiotic, there were an elevation of Penicillin and Aminoglycosides DDD average up to 528.66% and 137.66%.

**Conclusion:** There are trend changing of DDD average from WATCH to ACCESS category antibiotic following the 9 months implementation of e-RASPRO. We need further study to judge the effectiveness of e-RASPRO as a digital ASP tools.



Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

# Clinical eHealth

journal homepage: [www.keaipublishing.com/CEH](http://www.keaipublishing.com/CEH)




## A survey on define daily dose of watch- and access-category antibiotics in two Indonesian hospitals following the implementation of digital antimicrobial stewardship tool

Ronald Irwanto Natadidjaja<sup>a,b,\*</sup>, Aziza Ariyani<sup>a</sup>, Hadiani Adlani<sup>a,c</sup>, Raymond Adianto<sup>a</sup>, Iin Indah Pertiwi<sup>a</sup>, Grace Nerry Legoh<sup>a</sup>, Alvin Lekonardo Rantung<sup>a</sup>, Hadi Sumarsono<sup>a</sup>

<sup>a</sup> RASPRO Indonesia Study Group, Indonesia


<sup>b</sup> Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

<sup>c</sup> Faculty of Medicine, Syarif Hidayatullah State Islamic University, Banten, Indonesia



**Table 1**  
DDD of Prophylactic Antibiotics in Hospital 1 and Hospital 2 at 3 Months Following the Implementation of e-RASPRO Tool.

	Hospital 1			Hospital 2		
	Before Implementing e-RASPRO	After Implementing e-RASPRO	Increase / Decrease	Before Implementing e-RASPRO	After Implementing e-RASPRO	Increase / Decrease
Cefazolin	7.16	19.13	167.18 %	2.84	2.31	-18.66 %
Ceftriaxone	4.21	4.63	9.98 %	–	–	–
Cefotaxime	2.04	2.20	7.84 %	–	–	–





**@rasproindonesia**

*Instagram*

[www.new.rasproindonesia.com](http://www.new.rasproindonesia.com)

**THANK YOU!**