

Psychologic Factors in Temporomandibular Disorders and Somatization: A Multidimensional Analysis of Personality, Coping, and Distress Among Young Adults

by Carolina D Marpaung

Submission date: 06-Jan-2025 09:49AM (UTC+0700)

Submission ID: 2440171967

File name: IJP_2024.pdf (182.3K)

Word count: 7195

Character count: 38926

22

Psychologic Factors in Temporomandibular Disorders and Somatization: A Multidimensional Analysis of Personality, Coping, and Distress Among Young Adults

8

Adrian Ujin Yap, BDS, MSc, PhD

Department of Prosthodontics, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia; Department of Dentistry, Ng Teng Fong General Hospital, National University Health System; National Dental Research Institute Singapore, National Dental Centre and Duke-NUS Medical School, Singapore Health Services, Singapore.

13

Carolina Marpaung, BDS, PhD

Department of Prosthodontics, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia.

Purpose: To analyze the association of temporomandibular disorders (TMDs) and somatic symptoms with the psychologic variables of personality, coping, and distress in young adults. Physical and psychologic correlates were also explored, along with the risk factors for TMDs/somatization. **Materials and Methods:** Participants were enlisted from a local university, and the presence of TMDs and somatic symptoms was determined with the Short-form Fonseca Anamnestic Index and Patient Health Questionnaire-15. The psychologic variables of personality, coping, and distress were assessed with the Big-Five Personality Inventory-10; brief-COPE Inventory; and Depression, Anxiety, Stress Scales-21, respectively. Statistical evaluations were performed with Mann-Whitney *U* test, Spearman correlation, and logistic regression analyses ($\alpha = .05$). **Results:** Among the 455 participants (mean age: 22.7 ± 1.2 years), 18.2% and 5.7% had TMDs and medium-to-high somatization, respectively. Participants with TMDs exhibited substantially higher somatization and psychologic distress scores than those with no TMDs. Significant differences in TMDs, conscientiousness, extraversion, and psychologic distress scores were observed between participants with no-to-mild and medium-to-high somatization. The association between TMD and somatization scores was weak but significant. Neuroticism and dysfunctional coping style were moderately correlated to general distress, depression, anxiety, and stress ($r_s = 0.44$ to 0.62). **Conclusions:** Findings suggest that anxiety is the main risk factor for the presence of TMDs and medium-to-high somatization in nonclinical young adults, while conscientiousness is a protective factor for somatization. *Int J Prosthodont* 2024;37:605–614. doi: 10.11607/ijp.8590

Temporomandibular disorders (TMDs) comprise a cluster of musculoskeletal conditions affecting the stomatognathic system. Classically described TMD features include temporomandibular joint (TMJ)/masticatory muscle pain, TMJ sounds, and limited or deviated jaw movements.^{1,2} TMD prevalence ranges from 6% to 16% based on protocolized diagnostic criteria and $\leq 75\%$ of the general population have TMD signs/symptoms.^{3,4} They can be broadly divided into pain-related and intra-articular conditions.⁵ Women, particularly those of reproductive age, are at higher risk of TMDs.^{6,7} TMDs can negatively affect both the quality of sleep and life.^{8,9} The multifactorial etiology of TMDs and their adherence to the “biopsychosocial model of illness” has been confirmed by various studies.^{10,11} Among the psychologic variables implicated are somatization, psychologic distress, personality, and coping.^{11–15}

High prevalence and levels of somatization (expression of psychologic distress through somatic symptoms), depression, anxiety, and stress were observed in clinical

Correspondence to:
Dr Carolina Marpaung
carolina@trisakti.ac.id

Submitted January 11, 2023;
accepted March 28, 2023.
©2024 by Quintessence
Publishing Co Inc.

and nonclinical samples with TMDs.^{12–14,16,17} The importance of psychologic distress in the etiology of TMDs, as well as the strong correlation between TMDs and somatic symptoms, has led some to posit that TMDs are a type of “central sensitization syndrome” that encompass conditions such as fibromyalgia, chronic fatigue, and irritable bowel syndromes.^{18,19} East and Southeast Asians appear to be more susceptible to somatization due to the stigma, interpersonal sensitivity, and low social support accompanying mental illness.^{17,20,21}

In addition to psychologic distress, personality and coping have also been linked to TMDs and somatization.^{15,22,23} Personality is the set of traits and distinctive patterns of thoughts, feelings, and behaviors that makes a person unique, whereas coping is the set of predictable strategies that a person uses to minimize stress and reduce negative emotions. Personality traits and coping styles have independent and interactive effects on the way psychologic distress is controlled and relieved.²⁴ People with TMDs were reported to have “distressed” and “neurotic” (propensity toward negative affect and self-doubt) personality traits.^{15,25,26} Furthermore, they had also been found to employ more dysfunctional coping behaviors.^{15,27,28} Nevertheless, studies in this area are infrequent, and only one addressed all three interrelated psychologic variables concurrently in a small cohort of TMD patients.¹⁵

With these premises, the present study’s objectives were to analyze the association of TMD and somatic symptoms with the psychologic variables of personality, coping, and distress in a nondclinical community-based sample of young adults. Physical and psychologic correlates were also explored along with the psychologic predictors of TMDs and somatization. The research hypotheses were: (a) young adults with TMD and somatic symptoms have certain personality traits and dispositional coping styles as well as higher levels of psychologic distress, (b) the various physical and psychologic variables are correlated, and (c) the presence of TMDs and somatization are associated with specific psychologic risk factors.

MATERIALS AND METHODS

Study Design and Participants

Approval for this study was granted by the ethics committee of the Faculty of Dentistry, Universitas Trisakti (project number: 013/S3/KEPK/FKG/9/2021). Potential participants were recruited from young adults attending a local university either via public internet postings or in-person invitations. A nonprobabilistic voluntary sampling method was applied. The inclusion criteria were individuals aged 18 to 24 years old and proficient in the English language. The exclusion criteria were individuals with prior orofacial trauma/orthognathic surgery,

uncontrolled autoimmune or metabolic diseases, major psychiatric disorders, and drug/substance abuse. At least 363 participants were required for the study based on an estimated TMD prevalence of 60%, 95% CI level, 5% margin of error, and a student enrollment of 20,638 students.¹⁶ All potential participants were provided with the study information and informed consent was obtained from eligible individuals. An online survey comprising demographic information; the Short-form Fonseca Anamnestic Index (SFAI); Patient Health Questionnaire-15 (PHQ-15); Big Five Personality Inventory-10 (BFI-10); brief-COPE (Coping Orientations to Problems Experienced) Inventory (BCI); and Depression, Anxiety, Stress-Scales-21 (DASS-21) were subsequently administered.^{29–33} Participants were given the survey link and completed the questionnaires independently.

Physical Symptom Measures

The presence of TMDs was determined with the SFAI, which contains two pain (TMJ and masticatory muscle pain) and three function-related (TMJ sounds, opening, and side-movement difficulties) items. The SFAI demonstrated high accuracy (AUC 0.97–0.99), sensitivity (91% to 98%), and specificity (93% to 97%) for pain and/or intra-articular conditions when referenced to the DC/TMD standard.²⁹ Recently, its reliability and validity for identifying people with TMDs (as determined by the DC/TMD) were independently confirmed.³⁴ The items are scored using the following response scale: “no” = 0 points, “sometimes” = 5 points, and “yes” = 10 points. Total SFAI scores ≥ 15 points indicate the presence of TMDs with higher scores suggesting greater TMD symptom severity.

The presence of somatization was ascertained with the PHQ-15 which comprises the 15 most common “DSM-IV somatization disorder” somatic symptoms.^{30,35} The psychometric properties of the PHQ-15 are well recognized, and it has been incorporated into Axis II of the DC/TMD.^{5,36} The items are scored using the following three-point Likert scale: “not bothered at all” = 0 points, “bothered a little” = 1 point, and “bothered a lot” = 2 points. Total PHQ-15 scores of 5, 10, and 15 points indicate low, medium, and high somatic symptom severity respectively.³⁰ Participants were then dichotomized into “no TMDs” (NT) and with TMDs” (WT) in addition to “no-to-low” (NL) and “medium-to-high” (MH) somatization groups. Both SFAI and PHQ-15 were appraised over 30 days.

Psychologic Measures

Personality was evaluated with the BFI-10, which consists of two items for each of the “big five” personality dimensions, namely, openness, conscientiousness, extraversion, agreeableness, and neuroticism (OCEAN).³¹ The BFI-10 has been validated for investigating both

16

Table 1 Demographic Characteristics of the Study Sample

Table 1. Demographic Characteristics of the Study Sample							
Variable	n (%)	Age			Gender		
		Mean (SD)	Median (IQR)	P*	Male n (%)	Female n (%)	P**
Total	455 (100)	22.5 (1.2)	22.0 (2)	–	66 (14.5)	389 (85.5)	–
TMDs: SFAI							
NT	372 (81.8)	22.5 (1.2)	23.0 (2)	.409	55 (14.8)	317 (85.2)	.863
WT	83 (18.2)	22.4 (1.2)	22.0 (3)		11 (13.3)	72 (86.7)	
Somatization: PHQ-15							
NL	429 (94.3)	22.5 (1.2)	23.0 (2)	< .001	61 (14.2)	368 (85.8)	.563
MH	26 (5.7)	21.5 (1.5)	21.0 (2)		5 (19.2)	21 (80.8)	

*Indicates Mann-Whitney U tests $P < .05$.**Indicates chi-square test $P < .05$.

between-person traits and within-person processes.³⁷ The items are scored using a five-point Likert scale ranging from “disagree strongly” = 1 point to “agree strongly” = 5 points, with an item in each dimension assessed in reverse. Dimension scores are calculated with higher scores indicating greater partiality toward the specific trait.

Dispositional coping styles were explored with BCI, which has two items for each of the fourteen coping strategies employed in response to daily life stresses.³² The BCI is widely used for measuring coping and attempts to condense it has yielded contentious factor structures.³⁸ The items are scored using a four-point Likert scale ranging from “I haven’t been doing this at all” = 1 point to “I’ve been doing this a lot” = 4 points. The different strategies are categorized into the following three coping styles: problem-focused coping (active coping, instrumental support, and planning), emotion-focused (acceptance, emotional support, humor, positive reframing, and religion), and dysfunctional coping (behavioral disengagement, denial, self-distraction, self-blame, substance use, and venting).^{32,39} Coping style scores are derived by totaling the respective strategy scores. Higher coping style scores indicate more extensive use of functional (problem and emotion-focused) and dysfunctional coping strategies.

Psychologic distress was examined with the DASS-21, which includes seven items for each of the emotional subscales, namely, depression, anxiety, and stress.³³ The good measurement properties of the DASS-21 are also well-established. Additionally, the DASS-21 has been shown to have a bifactor structure consisting of a general factor for distress (negative affect) and the three subscales.⁴⁰ The items are scored using a four-point Likert scale ranging from “did not apply to me at all” = 0 points to “applied to me very much, or most of the time” = 3 points. Total and subscale DASS-21 scores are calculated with greater scores indicating higher levels of general

distress (total DASS-21), depressive, anxiety, and stress symptoms. The cut-off points for the various subscale severity classification (normal to extremely severe) are presented in the DASS manual.³³

Statistical Assessment

Statistical evaluations were carried out using the Statistical Package for the Social Sciences software version 27.0 (IBM) with a significance level of .05. Categorical data were presented as frequencies with percentages and analyzed using chi-square tests. Numerical data were presented as means/medians with SDs/interquartile ranges (IQRs) and analyzed for normality using Kolmogorov-Smirnov test. Because non-normal distributions were observed, numerical data were assessed using Mann-Whitney U tests and Spearman rank-order correlation. Correlation coefficients (rs) of 0.1, 0.4, and 0.7 indicate weak, moderate, and strong associations between the various variables.⁴¹ Univariate and multivariate logistic regression analyses were conducted to identify the risk factors for the presence of TMDs and medium-to-high somatization. A step-wise variable selection procedure was employed with a threshold of $P < .10$ for removing insignificant ones. Outcomes were presented as odds ratios (ORs) together with 95% CIs.

RESULTS

Out of a total of 487 young adults who signed up for the study, 32 were declined because they met the exclusion criteria. The mean age of the remaining 455 participants was 22.7 ± 1.2 years and 85.5% were women. TMDs and medium-to-high somatization were present in 18.2% and 5.7% of the study sample respectively. While age and gender distribution between the NT and WT groups did not vary considerably, the NL group was considerably older than the MH group (Table 1).

Table 2 Physical and Psychologic Variable Scores for NT and WT Groups

Physical/psychologic variable		NT	WT	P*	Differences
Somatization	PHQ-15 scores				
	Mean	3.9 (3.1)	5.7 (4.1)		
	Median	4.0 (4)	5.0 (4)	< .001	WT > NT
Personality	Openness				
	Mean	6.3 (1.4)	6.5 (1.5)		
	Median	6.0 (2)	6.0 (2)	.478	
	Conscientiousness				
	Mean	6.7 (1.4)	6.4 (1.2)		
	Median	7.0 (2)	6.0 (1)	.064	
	Extraversion				
	Mean	6.9 (1.7)	6.8 (1.6)		
	Median	7.0 (2)	7.0 (2)	.794	
	Agreeableness				
	Mean	7.1 (1.4)	7.0 (1.4)		
	Median	7.0 (2)	7.0 (2)	.463	
Coping styles	Neuroticism				
	Mean	6.7 (1.7)	7.01 (1.7)		
	Median	7.0 (2)	7.0 (2)	.082	
	Problem-focused				
	Mean	17.9 (3.2)	17.4 (3.7)		
	Median	18.0 (4)	18.0 (5)	.360	
	Emotion-focused				
	Mean	28.1 (4.7)	27.8 (5.1)		
	Median	28.0 (6)	28.0 (6)	.708	
	Dysfunctional				
	Mean	23.6 (4.3)	24.5 (4.6)		
	Median	23.0 (6)	25.0 (7)	.080	
Psychologic distress	General				
	Mean	13.8 (9.9)	17.5 (10.1)		
	Median	12.0 (14)	16.0 (13)	.001	WT > NT
	Depression				
	Mean	3.2 (3.5)	3.8 (3.5)		
	Median	2.0 (4)	3.0 (4)	.045	WT > NT
	Anxiety				
	Mean	4.4 (3.4)	6.1 (3.6)		
	Median	4.0 (4)	6.0 (5)	< .001	WT > NT
	Stress				
	Mean	6.3 (4.3)	7.5 (4.2)		
	Median	6.0 (6)	7.0 (7)	.012	WT > NT

Values are expressed as means (SD) and medians (IQR). Bold indicates $P < .05$.

*Mann-Whitney U test.

11

Tables 2 and 3 show the mean and median physical and psychologic variable scores for the different groups. The WT group had substantially greater PHQ-15 scores than the NT group. While variations in personality

dimension and coping style scores were insignificant, the WT group exhibited considerably higher general distress, depression, anxiety, and stress scores than the NT group.

**Table 3** Physical and Psychologic Variable Scores for the NL and MH Somatization Groups

Physical/psychologic variable		NL	MH	P*	Differences
TMD	SFAI scores				
	Mean	6.1 (8)	11.2 (8.9)	.001	MH > NL
	Median	5.0 (10)	10.0 (10)		
Personality	Openness				
	Mean	6.3 (1.5)	6.7 (1.2)		
	Median	6.0 (2)	7.0 (1)	.101	
	Conscientiousness				
	Mean	6.6 (1.4)	5.9 (0.9)		
	Median	7.0 (2)	6.0 (1)	.001	NL > MH
	Extraversion				
	Mean	6.9 (1.7)	6.2 (1.3)		
	Median	7.0 (2)	6.0 (2)	.022	NL > MH
	Agreeableness				
	Mean	7.1 (1.4)	6.8 (1.4)		
	Median	7.0 (2)	7.0 (3)	.211	
Coping styles	Neuroticism				
	Mean	6.7 (1.7)	7.2 (1.3)		
	Median	7.0 (2)	7.0 (2)	.158	
	Problem-focused				
	Mean	17.8 (3.3)	16.9 (3.3)		
	Median	18.0 (4)	17.5 (5)	.156	
	Emotion-focused				
	Mean	28.2 (4.8)	26.7 (5.2)		
	Median	28.0 (6)	26.0 (8)	.089	
	Dysfunctional				
	Mean	23.7 (4.4)	24.7 (3.9)		
	Median	23.0 (7)	25.0 (7)	.220	
Psychologic distress	General				
	Mean	14.2 (1.0)	19.7 (10.3)		
	Median	13.0 (13)	19.0 (12)	.004	MH > NL
	Depression				
	Mean	3.2 (3.5)	4.9 (3.4)		
	Median	2.0 (4)	5.0 (5)	.004	MH > NL
	Anxiety				
	Mean	4.6 (3.4)	6.7 (4.1)		
	Median	4.0 (4)	6.0 (5)	.005	MH > NL
	Stress				
	Mean	6.4 (4.3)	8.2 (3.8)		
	Median	6.0 (6)	7.5 (4)	.022	MH > NL

Values are expressed as means (SD) and medians (IQR). Bold indicates $P < .05$.

*Mann-Whitney U test.

Table 4 Correlations Between Physical and Psychologic Variables

Variable	SFAI	PHQ-15	Openness	Conscientiousness	Extraversion	Agreeableness	Neuroticism
SFAI	-	-	-	-	-	-	-
PHQ-15	0.21*	0.27*	-	-	-	-	-
Openness	0.05	0.02					
Conscientiousness	-0.13*	-0.15*	-0.23**	-	-	-	-
Extraversion	-0.05	-0.08	-0.15**	0.18**	-	-	-
Agreeableness	-0.06	-0.11*	0.03	0.01	0.09	-	-
Neuroticism	0.11*	0.14**	0.22**	-0.21**	-0.28**	-0.01	-
Problem-focused	-0.04	-0.07	-0.06	0.17**	0.13**	0.04	-0.10**
Emotion-focused	-0.02	-0.05	-0.09	0.09	0.19**	0.11*	-0.15**
Dysfunction	0.08	0.18**	0.16**	-0.18**	-0.06	-0.10*	0.24**
General	0.15*	0.26**	0.27**	-0.26**	-0.29**	-0.04	0.59**
Depression	0.10*	0.22**	0.25**	-0.26**	-0.34**	-0.08	0.46**
Anxiety	0.18**	0.27**	0.23**	-0.22**	-0.19**	0.01	0.51**
Stress	0.13*	0.26**	0.24**	-0.22**	-0.24**	-0.05	0.62**
Variable	Problem-focused	Emotion-focused	Dysfunctional	General	Depression	Anxiety	
SFAI	-	-	-	-	-	-	
PHQ-15	-	-	-	-	-	-	
Openness	-	-	-	-	-	-	
Conscientiousness	-	-	-	-	-	-	
Extraversion	-	-	-	-	-	-	
Agreeableness	-	-	-	-	-	-	
Neuroticism	-	-	-	-	-	-	
Problem-focused	-	-	-	-	-	-	
Emotion-focused	0.73**	-	-	-	-	-	
Dysfunction	0.36**	0.32**	-	-	-	-	
General	0.02	-0.03	0.50**	-	-	-	
Depression	-0.10*	-0.10*	0.45**	0.84**	-	-	
Anxiety	0.08	0.04	0.44**	0.88**	0.60**	-	
Stress	0.05	-0.01	0.47**	0.95**	0.71**	0.79**	

Results of Spearman correlation. Bold indicates correlation coefficient > 0.4.

* $P < .05$.** $P < .01$.

The MH group had substantially greater SFAI scores than the NL group. Significant differences in conscientiousness and extraversion scores, but not coping style scores, were discerned. The MH group had considerably higher general distress, depression, anxiety, and stress scores than the NL group.

Tables 4 and 5 display the outcomes of correlational and regression analyses. The correlation between SFAI and PHQ-15 scores, although significant, was weak ($r_s = 0.21$) as with their associations with general distress, depression, anxiety, and stress ($r_s = 0.13$ – 0.27). Neuroticism and dysfunctional coping style were moderately associated with general distress, depression, anxiety, and stress ($r_s = 0.44$ – 0.62).

Strong correlations were noted between the two functional coping styles (problem and emotion-focused coping), as well as between general distress and all three DASS-21 subscales ($r_s = 0.73$ – 0.95). While univariate exploration showed that TMDs were related to general distress, anxiety, and stress, multivariate analysis indicated that only anxiety (OR = 1.15; 95% CI = 1.07–1.22) was a risk factor for the presence of TMDs. Conscientiousness, extraversion, general distress, depression, anxiety, and stress were all related to somatic symptoms with the univariate model. However, multivariate analysis indicated that conscientiousness (OR = 0.70; 95% CI = 0.51–0.95) and anxiety (OR = 1.13; 95% CI = 1.02–1.26) were protective and risk

Table 5 Risk Factors for the Presence of TMDs and Medium-to-High Somatization

Risk factor		Univariate		Multivariate	
		OR (95% CI)	P*	OR (95% CI)	P**
Presence of TMDs					
Gender	Male	Reference			
	Female	1.14 (0.57–2.28)	.720		
Personality	Openness	1.10 (0.93–1.30)	.273		
	Conscientiousness	0.86 (0.71–1.02)	.081		
	Extraversion	0.98 (0.84–1.13)	.739		
	Agreeableness	0.95 (0.81–1.14)	.601		
	Neuroticism	1.13 (0.98–1.31)	.091		
Coping	Problem-focused	0.89 (0.72–1.10)	.299		
	Emotion-focused	0.93 (0.73–1.19)	.932		
	Dysfunctional	1.34 (0.97–1.85)	.077		
Psychologic distress	General	1.04 (1.01–1.06)	.003		
	Depression	1.05 (0.99–1.12)	.131		
	Anxiety	1.15 (1.07–1.22)	< .001	1.15 (1.07–1.22)	< .001
	Stress	1.07 (1.01–1.12)	.014		
Presence of MH somatization					
Gender	Male	Reference			
	Female	0.70 (0.25–1.92)	.696		
Personality	Openness	1.23 (0.94–1.61)	.140		
	Conscientiousness	0.64 (0.47–0.87)	.004	0.70 (0.51–0.95)	.021
	Extraversion	0.76 (0.59–0.97)	.030		
	Agreeableness	0.83 (0.62–1.11)	.209		
	Neuroticism	1.20 (0.94–1.53)	.150		
Coping	Problem-focused	0.80 (0.56–1.13)	.197		
	Emotion-focused	0.73 (0.48–1.09)	.122		
	Dysfunctional	1.35 (0.80–2.29)	.263		
Distress	General	1.05 (1.01–1.09)	.007		
	Depression	1.12 (1.02–1.23)	.018		
	Anxiety	1.16 (1.05–1.28)	.003	1.13 (1.02–1.26)	.021
	Stress	1.09 (1.00–1.19)	.043		

Results of univariate and multivariate logistic regression analyses. Bold indicates $P < .05$.

factors for the presence of medium-to-high somatization correspondingly.

DISCUSSION

The present study is the first to establish the association of TMD and somatic symptoms with the psychologic variables of personality, coping, and distress in nonclinical community-based young adults. All three research hypotheses were partly supported because participants with TMD/somatic symptoms exhibited particular personality traits or greater psychologic distress, certain psychologic variables were moderate to strongly correlated, and anxiety was a risk factor for both TMDs

and somatization. Young adults were chosen because they represented the bulk of TMD patients and the peak age for the occurrence of TMD symptoms.^{2,42} In addition, university students were identified because they have high levels of life, social, and academic stress as well as high rates of psychologic distress.⁴³ Because somatic symptoms are frequently experienced by the general population, participants were dichotomized into no-to-low and medium-to-high somatization groups.^{44,45} Personality and coping can play both independent and interactive roles in influencing physical and psychologic symptoms and thus were analyzed synchronously together with general distress, depression, anxiety, and stress.⁴⁶

TMDs and Psychologic Variables

The prevalence of TMDs observed with the SFAI was consistent with that reported for the general population when established with protocolized diagnostic criteria.³ While personality dimension and coping style scores were statistically insignificant, participants with TMDs had substantially higher total and subscale DASS-21 scores than their peers without TMDs. Therefore, personality and coping may not have much bearing on the manifestation of TMD symptoms when compared to psychologic distress. Stress is the feeling of emotional or physical tension in response to pressure or adversity, whereas anxiety and depression are the persistent feelings of unease or apprehension and low mood and despair, respectively. Although significant differences in all psychologic distress variables were discerned between the WT and NT groups, the univariate model indicated that only general distress, anxiety, and stress were associated with TMDs. This result can be attributed to the largely normal level of depression (0 to 4 points) detected in the young adults examined.

The few studies conducted in clinical samples yielded disparate results with TMD patients having more “neurotic” personalities and dysfunctional or maladaptive coping styles.^{15,26–28} Ferrando et al,¹⁵ who explored all three psychologic variables concurrently, found that neuroticism (the tendency to experience negative emotions) and depression featured predominantly in TMD patients with muscle disorders, whereas conscientiousness was present in those with joint disorders. Furthermore, patients with muscle disorders also used functional coping strategies less regularly than the control group. The discrepancy in findings could be ascribed in part to the greater prevalence and severity of depression and anxiety among TMD patients, which is thought to be closely related to personality traits and mediated by dispositional coping.^{12–14,24} In addition to normal levels of depression, the nonclinical young adults with TMDs presented only moderate anxiety (6 to 7 points) and mild stress (8 to 9 points).

Somatization and Psychologic Variables

About 6% of the participants reported medium-to-high somatic symptoms, which was comparable to the estimated 8% prevalence of somatic symptom disorder in general practice.⁴⁷ Though substantial variations in conscientiousness (the tendency to be self-disciplined, well-organized, and goal-oriented), extraversion (the tendency to be friendly and outgoing), total, and subscale DASS-21 scores were noted between the NL and MH groups, no significant differences in coping style scores were perceived. The univariate analysis yielded similar results. Hence, dispositional coping styles may play a lesser role in the manifestation of somatic symptoms. Participants with medium-to-high somatization

were found to be less conscientious/extroverted and had considerably higher levels of general distress, depression, anxiety, as well as stress. Because individuals with more somatic symptoms have greater psychologic distress, they may not be in the best mental state to work hard or socialize. However, Rokvić et al,²³ in their initial investigation, stated that neuroticism and anxiety were most closely related to somatization in the general population. Notwithstanding, studies on the relationship of TMDs/somatization to personality and dispositional coping styles remain scant and further cross-cultural research in this area is warranted.

Correlations and Protective/Risk Factors

Though the WT and MH groups had significantly higher PHQ-15 and SFAI scores than the NT and NL groups, the correlation between SFAI and PHQ-15 scores was weak. This relationship is anticipated to be stronger in TMD patient populations given their high prevalence of comorbid chronic pain conditions and medium-to-high somatization.^{14,48} The normal levels of depression and lower severity of anxiety and stress explained the weak associations between SFAI/PHQ-15 and psychologic distress scores when compared to TMD patients.⁴⁹ Neuroticism and dysfunctional coping were found to be moderately correlated to general distress, depression, anxiety, and stress. Polygenic analysis has established the genetic connections between personality, particularly neurotic traits, and psychopathology.⁵⁰ Furthermore, it was found that personality and coping jointly accounted for up to 50% of the variance in psychopathology.⁵¹

Panayiotou et al²⁴ determined that neuroticism was related to dysfunctional coping, whereas conscientiousness and extraversion were associated with functional coping in a community sample. In the present study, the correlations between neuroticism and dysfunctional coping as well as conscientiousness and extraversion and functional coping were mostly significant albeit weak. The correlation between the two functional coping styles was strong, suggesting that individuals who employed problem-focused coping strategies also used emotion-focused ones. However, this relationship could be influenced by moderators such as type of illness, study design, and context, in addition to individual beliefs about coping and interventions.⁵² The strong correlations between general distress and the depression, anxiety, and stress subscales provided additional support for the bifactor structure of the DASS-21.⁴⁰

Multidimensional analysis of personality, coping, and distress with the multivariate model revealed that anxiety was the primary risk factor for the presence of TMDs and medium-to-high somatization, increasing their prospects by 15% and 13%, respectively. These findings corroborated other studies concerning nonclinical adolescent/young adult samples that showed anxiety (a



perceived stress response) and stress were predictors of TMDs and somatization.^{17,53,54} Moreover, polymorphism in catechol-O-methyltransferase was associated with both TMDs and anxiety in young people.⁵⁵ Conscientiousness was found to be a protective factor for medium-to-high somatization reducing it by 30%. This result supported the inverse relationship between conscientiousness and somatic symptoms reported in a prior study on university students.⁵⁶ Conscientiousness and two of its facets, specifically order and industriousness, can influence health outcomes by modifying the effects of daily stressors.⁵⁷ Given the connection between TMDs/somatization and psychologic distress, positive psychologic interventions including mindfulness-based stress reduction and cognitive-behavioral therapies that reduce anxiety and stress and promote resilience could conceivably reduce the incidence of TMD/somatic symptoms in young people. They have also been shown to improve pain, depression, and anxiety symptoms as well as the quality of life of patients.^{58,59}

Study Limitations

This observational study has its limitations. First, causal relationships among the various physical and psychologic variables cannot be established with the cross-sectional design employed. Longitudinal investigations are necessary to better explore causality and sequential interactions between the different factors. Second, the study sample involved only university students and consisted of more women. Other young, middle-aged, and older adult groups as well as more men must be incorporated in future work to allow for the generalization of findings and to minimize possible age-gender predilections. Third, only Asian young adults were studied, and results cannot be extrapolated to other racial and ethnic groups. Therefore, the study needs to be extended to other countries and cultures before absolute conclusions can be drawn. The research should also be repeated in TMD patient populations considering their probable variances in personality, coping, and distress when contrasted to nonclinical community samples. Lastly, as physical and psychologic variables were self-reported, they may be exposed to various information partialities including recall, social desirability, confirmation, and other biases.⁶⁰

CONCLUSIONS

This study is the first to determine the relationship of TMD/somatic symptoms with personality, coping, and psychologic distress in nonclinical community-based young adults. TMDs and medium-to-high somatization were present in 18.2% and 5.7% of the study sample. Young adults with TMDs had substantially higher levels of general distress, depression, anxiety, and stress than their counterparts with no TMDs. Those with medium-to-high somatization were low in conscientiousness as well as

extraversion and exhibited higher levels of psychologic distress compared to their peers with normal-to-low somatic symptoms. Neuroticism and dysfunctional coping style were found to be moderately correlated to general distress, depression, anxiety, and stress. Multivariate regression analysis determined that anxiety was the primary risk factor for the presence of TMDs and medium-to-high somatization and indicated that conscientiousness could be a protective factor for somatization in young adults. Because positive psychologic interventions were shown to be effective for managing both pain and psychologic symptoms, they hold promise as adjunctive therapies for various aspects of TMDs and somatization.

ACKNOWLEDGMENTS

The authors would like to thank I. Hanin, Y. Pragustine, and A. Fitryanur for their assistance with the data collection. This research was funded by grant no. 0142/PUF/FKG/2021-202 from Universitas Trisakti. The authors declare that they have no financial or personal interests related to the present work.

REFERENCES

1. Kapos FP, Exposto FG, Oyarzo JF, Durham J. Temporomandibular disorders: A review of current concepts in aetiology, diagnosis and management. *Oral Surg* 2020;13:321–334.
2. List T, Jensen RH. Temporomandibular disorders: Old ideas and new concepts. *Cephalalgia* 2017;37:692–704.
3. Manfredini D, Guarda-Nardini L, Winocur E, Piccotti F, Ahlberg J, Lobbezoo F. Research diagnostic criteria for temporomandibular disorders: A systematic review of axis I epidemiologic findings. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2011;112:453–462.
4. Ryan J, Akhter R, Hassan N, Hilton G, Wickham J, Ibaragi S. Epidemiology of Temporomandibular Disorder in the general population: A systematic review. *Adv Dent Oral Health* 2019;10:555787.
5. Schiffman E, Ohrbach R, Truelove E, et al. Diagnostic Criteria for Temporomandibular Disorders (DC/TMD) for clinical and research applications: Recommendations of the international rdc/tmd consortium network and orofacial pain special interest group. *J Oral Facial Pain Headache* 2014;28:6–27.
6. Bueno CH, Pereira DD, Pattussi MP, Grossi PK, Grossi ML. Gender differences in temporomandibular disorders in adult populational studies: A systematic review and meta-analysis. *J Oral Rehabil* 2018;45:720–729.
7. Warren MP, Fried JL. Temporomandibular disorders and hormones in women. *Cells Tissues Organs* 2001;169:187–192.
8. Roithmann CC, Silva CAGD, Pattussi MP, Grossi ML. Subjective sleep quality and temporomandibular disorders: Systematic literature review and meta-analysis. *J Oral Rehabil* 2021;48:1380–1394.
9. Pigozzi LB, Pereira DD, Pattussi MP, et al. Quality of life in young and middle age adult temporomandibular disorders patients and asymptomatic subjects: A systematic review and meta-analysis. *Health Qual Life Outcomes* 2021;19:83.
10. Slade GD, Fillingim RB, Sanders AE, et al. Summary of findings from the OPPERA prospective cohort study of incidence of first-onset temporomandibular disorder: Implications and future directions. *J Pain* 2013;14(12 suppl):T116–T124.
11. Fillingim RB, Ohrbach R, Greenspan JD, et al. Psychological factors associated with development of TMD: The OPPERA prospective cohort study. *J Pain* 2013;14(12 suppl):T75–T90.
12. Dos Santos EA, Peinado BRR, Frazão DR, et al. Association between temporomandibular disorders and anxiety: A systematic review. *Front Psychiatry* 2022;13:990430.

13. Reis PHF, Laxe LAC, Lacerda-Santos R, Münchow EA. Distribution of anxiety and depression among different subtypes of temporomandibular disorder: A systematic review and meta-analysis. *J Oral Rehabil* 2022;49:754–767.
14. De La Torre Canales G, Câmara-Souza MB, Muñoz Lora VRM, et al. Prevalence of psychosocial impairment in temporomandibular disorder patients: A systematic review. *J Oral Rehabil* 2018;45:881–889.
15. Ferrando M, Andreu Y, Galdón MJ, Durá E, Poveda R, Bagán JV. Psychological variables and temporomandibular disorders: Distress, coping, and personality. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2004;98:153–160.
16. Yap AU, Lee DZR, Marpaung C. Negative affectivity and emotions in youths with temporomandibular disorders across cultures. *Cranio* 2022;1–9.
17. Yap AU, Sultana R, Natu VP. Somatic and temporomandibular disorder symptom—Idioms of psychological distress in Southeast Asian youths. *Cranio* 2024;42:364–371.
18. Yap AU, Natu VP. Inter-relationships between pain-related temporomandibular disorders, somatic and psychological symptoms in Asian youths. *J Oral Rehabil* 2020;47:1077–1083.
19. Fantoni F, Salvetti G, Manfredini D, Bosco M. Current concepts on the functional somatic syndromes and temporomandibular disorders. *Stomatologia* 2007;9:3–9.
20. Sun Y, Chen G, Wang L, et al. Perception of stigma and its associated factors among patients with major depressive disorder: A multicenter survey from an Asian population. *Front Psychiatry* 2019;10:321.
21. Dreher A, Hahn E, Diefenbacher A, et al. Cultural differences in symptom representation for depression and somatization measured by the PHQ between Vietnamese and German psychiatric outpatients. *J Psychosom Res* 2017;102:71–77.
22. Franck E, Haegdorens F, Goossens E, et al. The role of coping behavior in healthcare workers' distress and somatization during the covid-19 pandemic. *Front Psychol* 2021;12:684618.
23. Rokvić N. Initial investigation of somatization in the general population of Serbia: Prevalence, manifestations and predictors. *Medicinski pregled* 2018;71:60–367.
24. Panayiotou G, Kokkinos CM, Kapsou M. Indirect and direct associations between personality and psychological distress mediated by dispositional coping. *J Psychol* 2014;148:549–567.
25. Gębska M, Dalewski B, Pałka Ł, Kołodziej Ł, Sobolewska E. The importance of Type D personality in the development of Temporomandibular Disorders (TMDs) and depression in students during the COVID-19 pandemic. *Brain Sci* 2021;12:28.
26. Michelotti A, Martina R, Russo M, Romeo R. Personality characteristics of temporomandibular disorder patients using M.M.P.I. *Cranio* 1998;16:119–125.
27. Reissmann DR, John MT, Schierz O, Seedorf H, Doering S. Stress-related adaptive versus maladaptive coping and temporomandibular disorder pain. *J Orofac Pain* 2012;26:181–190.
28. Galdón MJ, Durá E, Andreu Y, Ferrando M, Poveda R, Bagán JV. Multidimensional approach to the differences between muscular and articular temporomandibular patients: coping, distress, and pain characteristics. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2006;102:40–46.
29. Yap AU, Zhang MJ, Lei J, Fu KY. Diagnostic accuracy of the short-form Fonseca Anamnestic Index in relation to the Diagnostic Criteria for Temporomandibular Disorders. *J Prosthet Dent* 2022;128:977–983.
30. Kroenke K, Spitzer RL, Williams JB. The PHQ-15: Validity of a new measure for evaluating the severity of somatic symptoms. *Psychosom Med* 2002;64:258–266.
31. Rammstedt B, John, OP. Measuring personality in one minute or less: A 10 item short version of the Big Five Inventory in English and German. *J Res Pers* 2007;41:203–212.
32. Cooper C, Katona C, Livingston G. Validity and reliability of the brief COPE in carers of people with dementia: the LASER-AD Study. *J Nerv Ment Dis* 2008;196:838–843.
33. Lovibond SH, Lovibond PF. Manual for the Depression Anxiety & Stress Scales, ed 2. Psychology Foundation, 1995.
34. Zagalaz-Anula N, Sánchez-Torrelo CM, Acebal-Blanco F, et al. The short form of the Fonseca Anamnestic Index for the screening of Temporomandibular Disorders: Validity and reliability in a Spanish-speaking population. *J Clin Med* 2021;10:5858.
35. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders DSM-IV-TR, ed 4. American Psychiatric Press, 2000.
36. Sitnikova K, Dijkstra-Kersten SMA, Mookink LB, et al. Systematic review of measurement properties of questionnaires measuring somatization in primary care patients. *J Psychosom Res* 2017;103:42–62.
37. Ringwald WR, Manuck SB, Marsland AL, Wright AGC. Psychometric evaluation of a big five personality state scale for intensive longitudinal studies. *Assessment* 2022;29:1301–1319.
38. Solberg MA, Gridley MK, Peters RM. The factor structure of the brief cope: A systematic review. *West J Nurs Res* 2022;44:612–627.
39. Henry N, Kayser D, Egermann H. Music in mood regulation and coping orientations in response to COVID-19 lockdown measures within the United Kingdom. *Front Psychol* 2021;12:647879.
40. Lee J, Lee EH, Moon SH. Systematic review of the measurement properties of the Depression Anxiety Stress Scales-21 by applying updated COSMIN methodology. *Qual Life Res* 2019;28:2325–2339.
41. Dancy CP, Reidy J. Statistics Without Maths for Psychology, ed 7. Pearson, 2017.
42. Yap AU, Cao Y, Zhang MJ, Lei J, Fu KY. Age-related differences in diagnostic categories, psychological states and oral health-related quality of life of adult temporomandibular disorder patients. *J Oral Rehabil* 2021;48:361–368.
43. Regehr C, Glancy D, Pitts A. Interventions to reduce stress in university students: A review and meta-analysis. *J Affect Disord* 2013;148:1–11.
44. Hinz A, Ernst J, Glaesmer H, et al. Frequency of somatic symptoms in the general population: Normative values for the Patient Health Questionnaire-15 (PHQ-15). *J Psychosom Res* 2017;96:27–31.
45. Petrie KJ, Faasse K, Crichton F, Grey A. How common are symptoms? Evidence from a New Zealand national telephone survey. *BMJ Open* 2014;4:e005374.
46. Carver CS, Connor-Smith J. Personality and coping. *Annu Rev Psychol* 2010;61:679–704.
47. Lehmann M, Pohontsch NJ, Zimmermann T, Scherer M, Löwe B. Estimated frequency of somatic symptom disorder in general practice: Cross-sectional survey with general practitioners. *BMC Psychiatry* 2022;22:632.
48. Kleykamp BA, Ferguson MC, McNicol E, et al. The prevalence of comorbid chronic pain conditions among patients with temporomandibular disorders: A systematic review. *J Am Dent Assoc* 2022;153:241–250.e10.
49. Yap AU, Zhang MJ, Cao Y, Lei J, Fu KY. Comparison of psychological states and oral health-related quality of life of patients with differing severity of temporomandibular disorders. *J Oral Rehabil* 2022;49:177–185.
50. Sanchez-Roige S, Gray JC, MacKillop J, Chen CH, Palmer AA. The genetics of human personality. *Genes Brain Behav* 2018;17:e12439.
51. Mirmics Z, Heincz O, Bagdy G, et al. The relationship between the big five personality dimensions and acute psychopathology: Mediating and moderating effects of coping strategies. *Psychiatr Danub* 2013;25:379–388.
52. Hagger MS, Koch S, Chatzisarantis NLD, Orbell S. The common sense model of self-regulation: Meta-analysis and test of a process model. *Psychol Bull* 2017;143:1117–1154.
53. de Paiva Bertoli FM, Bruzamin CD, de Almeida Kranz GO, Losso EM, Brancher JA, de Souza JF. Anxiety and malocclusion are associated with temporomandibular disorders in adolescents diagnosed by RDC/TMD. A cross-sectional study. *J Oral Rehabil* 2018;45:747–755.
54. Lei J, Fu J, Yap AU, Fu KY. Temporomandibular disorders symptoms in Asian adolescents and their association with sleep quality and psychological distress. *Cranio* 2016;34:242–249.
55. Brancher JA, Spada PP, Meger MN, et al. The association of genetic polymorphisms in serotonin transporter and catechol-O-methyltransferase on temporomandibular disorders and anxiety in adolescents. *J Oral Rehabil* 2019;46:597–604.
56. Razagh-Pour M, Hoseinzadeh AA. The mediating role of Cognitive Emotion Regulation Strategies in the relationship between personality traits with somatization symptoms of female university students. *Quart Appl Psychol* 2019;13:295–316.
57. Gartland N, O'Connor DB, Lawton R, Ferguson E. Investigating the effects of conscientiousness on daily stress, affect and physical symptom processes: A daily diary study. *Br J Health Psychol* 2014;19:311–328.
58. Hilton L, Hempel S, Ewing BA, et al. Mindfulness meditation for chronic pain: Systematic review and meta-analysis. *Ann Behav Med* 2017;51:199–213.
59. Lakan SE, Schofield KL. Mindfulness-based therapies in the treatment of somatization disorders: A systematic review and meta-analysis. *PLoS One* 2013;8:e71834.
60. Althubaiti A. Information bias in health research: Definition, pitfalls, and adjustment methods. *J Multidiscip Healthc* 2016;9:211–217.

Copyright of International Journal of Prosthodontics is the property of Quintessence Publishing Company Inc. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.

Psychologic Factors in Temporomandibular Disorders and Somatization: A Multidimensional Analysis of Personality, Coping, and Distress Among Young Adults

ORIGINALITY REPORT

19%	12%	16%	3%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	mafiadoc.com Internet Source	3%
2	pdfslide.net Internet Source	2%
3	www.unboundmedicine.com Internet Source	1%
4	Submitted to University of Sydney Student Paper	1%
5	Adrian Ujin Yap, Ji Woon Park, Jie Lei, Chengge Liu, Seong Hae Kim, Byeong-min Lee, Kai Yuan Fu. "The influence of the COVID-19 pandemic, sex, and age on temporomandibular disorders subtypes in East Asian patients: a retrospective observational study", BMC Oral Health, 2023 Publication	1%
6	Adrian Ujin Yap, Jie Lei, Ji Woon Park, Chengge Liu, Seong Hae Kim, Byeong-Min Lee, Kai-	1%

Yuan Fu. "Age distribution of East Asian TMD patients and age-related differences in DC/TMD axis I findings", CRANIO®, 2024

Publication

7

Adrian Ujin Yap, Chengge Liu, Jie Lei, Ji Woon Park, Seong Hae Kim, Byeong-min Lee, Kai Yuan Fu. "DC/TMD axis I subtyping: generational and gender variations among East Asian TMD patients", BMC Oral Health, 2023

Publication

1 %

8

Adrian Ujin Yap, Brandon Ming Hong Peng, Su Yin Lim, Darren Zong Ru Lee. "Association of temporomandibular disorders severity with otologic and concomitant pain symptoms in Asian youths", CRANIO®, 2022

Publication

1 %

9

Adrian Ujin Yap, Muzaffar Apipi, Siti Mazlipah Ismail, Zamros Yuzadi Mohd Yusof, Kathreena Kadir. "Psychometric properties of the Malay Fonseca Anamnestic Index for temporomandibular disorders", CRANIO®, 2022

Publication

1 %

10

meridian.allenpress.com

Internet Source

1 %

11

Adrian Ujin Yap, Carolina Marpaung, Enrita Dian Rahmadini. "Psychological well-being

1 %

and distress: Their associations with Temporomandibular disorder symptoms and inter-relationships", Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology, 2021

Publication

12

Submitted to Henry M. Jackson High School

Student Paper

1 %

13

Adrian Ujin Yap, Carolina Marpaung, Enrita Dian Rahmadini. "Psychological well-being and distress: Their associations with temporomandibular disorder symptoms and interrelationships", Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology, 2021

Publication

1 %

14

Adrian Ujin Yap, Sharon Hui Xuan Tan, Carolina Marpaung. "Temporomandibular disorder symptoms in young adults: Three-dimensional impact on oral health-related quality of life", Journal of Oral Rehabilitation, 2022

Publication

<1 %

15

www.frontiersin.org

Internet Source

<1 %

16

mdpi-res.com

Internet Source

<1 %

17	www.dovepress.com Internet Source	<1 %
18	Adrian Ujin Yap, Ye Cao, Minjuan Zhang, Jie Lei, Kai-Yuan Fu. "Age-related differences in diagnostic categories, psychological states, and oral health-related quality of life of adult Temporomandibular Disorder patients", <i>Journal of Oral Rehabilitation</i> , 2020 Publication	<1 %
19	Adrian Ujin Yap, Henry Chee Wai Ho, Ye Choung Lai. "Analysing the psychosocial construct of Temporomandibular disorders: Implications for orthodontics", <i>Seminars in Orthodontics</i> , 2023 Publication	<1 %
20	Adrian Ujin Yap, Ye Cao, Min-Juan Zhang, Jie Lei, Kai-Yuan Fu. "Age-related differences in diagnostic categories, psychological states and oral health-related quality of life of adult temporomandibular disorder patients", <i>Journal of Oral Rehabilitation</i> , 2020 Publication	<1 %
21	Submitted to Nanyang Polytechnic Student Paper	<1 %
22	Submitted to University of Portsmouth Student Paper	<1 %

23 Adrian Ujin Yap, Vaishali Prakash Natu. "Inter-relationships between pain-related temporomandibular disorders, somatic and psychological symptoms in Asian youths", Journal of Oral Rehabilitation, 2020
Publication <1 %

24 www.medicinaoral.com
Internet Source <1 %

25 Adrian Ujin Yap, Carolina Marpaung. "Severity and form of temporomandibular disorder symptoms: Functional, physical, and psychosocial impacts", CRANIO®, 2021
Publication <1 %

26 Adrian Ujin Yap, Ye Cao, Min-Juan Zhang, Jie Lei, Kai-Yuan Fu. "Number and type of temporomandibular disorder symptoms: their associations with psychological distress and oral health-related quality of life", Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology, 2021
Publication <1 %

27 Submitted to University of Southampton
Student Paper <1 %
