

JOURNAL OF DENTISTRY INDONESIA

UNIVERSITAS INDONESIA









• Garuda

₩ebsite

3 Editor URL

History Accreditation

2018 2019 2020 2021 2022 2023 2024

Google Scholar

Treatment of Oral Vascular Malformations with Ethamolin: A Report of Two Cases

<u>UI Scholars Hub</u> <u>Nournal of Dentistry Indonesia</u>

<u>□ 2023</u> <u>□ DOI: -</u> <u>○ Accred : Sinta 2</u>

Ramsay Hunt Syndrome with Oral Findings: A Rare Case

<u>UI Scholars Hub</u> <u>Journal of Dentistry Indonesia</u>

<u>□ 2023</u> <u>□ DOI: -</u> <u>○ Accred : Sinta 2</u>

<u>Treatment of Gingival Fenestration using Mucoperiosteal Flap with Connective Tissue Graft: A</u> <u>Report of Two Cases</u>

<u>UI Scholars Hub</u> <u>Journal of Dentistry Indonesia</u>

<u>□ 2023</u> <u>□ DOI: -</u> <u>○ Accred : Sinta 2</u>

Do Nasal Anatomical Variations Affect the Maxillary Sinus? A CBCT Volumetric Analysis

<u>UI Scholars Hub</u> <u>Journal of Dentistry Indonesia</u>

<u>□ 2023</u> <u>□ DOI: -</u> <u>○ Accred : Sinta 2</u>

Assessment of DNA Damage of Oral Mucosa due to Ceramic Bracket Using Comet Assay and Mutagenicity of Orthodontic Bonding System Using Ames Test

<u>UI Scholars Hub</u> <u>Nournal of Dentistry Indonesia</u>

<u>□ 2023</u> <u>□ DOI: -</u> <u>○ Accred : Sinta 2</u>


<u>A Prospective-Randomized Study: The Impact of Four Different Caries Removal Method on Pain and Clinical Evaluations</u>

<u>UI Scholars Hub</u> <u>Journal of Dentistry Indonesia</u>

<u>□ 2023</u> <u>□ DOI: -</u> <u>○ Accred : Sinta 2</u>

Get More with SINTA Insight

Go to Insight



Journal By Google Scholar					
	All	Since 2019			
Citation	2176	1397			
h-index	20	16			
i10-index	61	36			

Studentsâ | Experience, Self-Confidence, and Perception Toward Endodontic Learning:

National Survey Among Malaysian Dental Schools

<u>UI Scholars Hub</u>

■ <u>Journal of Dentistry Indonesia</u>

□ 2023

<u> ■ DOI: -</u>

O Accred: Sinta 2

<u>Management of Tooth Hypersensitivity Under a Cantilever Fixed Partial Denture Using a Novel</u>
<u>Low-Level Laser Therapy Protocol: A Case Report</u>

<u>UI Scholars Hub</u>

Journal of Dentistry Indonesia

□ 2023

<u> ■ DOI: -</u>

O Accred: Sinta 2

<u>Class III Malocclusion Camouflage Treatment Using a Conventional Orthodontic Appliance in a Non-Growing Patient</u>

<u> UI Scholars Hub</u>

■ <u>Journal of Dentistry Indonesia</u>

2023

<u> ■ DOI: -</u>

O Accred: Sinta 2

<u>A Two-Year Retrospective Evaluation of Cone Beam Computed Tomography Indications in Pediatric, Adolescent and Adult Patients</u>

<u>UI Scholars Hub</u>

Journal of Dentistry Indonesia

2023

O Accred: Sinta 2



Get More with SINTA Insight

Go to Insight

Citation Per Year By Google Scholar

	Journal By Google Scholar	
	All	Since 2019
Citation	2176	1397
h-index	20	16
i10-index	61	36

P-ISSN 1693-9697 E-ISSN 2355-4800 Volume 30 Number 3 December 2023

Journal of Dentistry Indonesia

The Official Publication of The Faculty of Dentistry, Universitas Indonesia in Collaboration with Indonesian Dental Association

Journal of **Dentistry** Indonesia

Home > Journals > Faculty of Dentistry > JDI

Editorial Team

EDITOR IN CHIEF

Yuniardini Septorini Wimardhani, (SCOPUS ID=56784363100) Department of Oral Medicine, Faculty of Dentistry, Universitias Indonesia, Indonesia, Indonesia

SECTION EDITORS

Lisa Rinanda Amir, (SCOPUS ID = 14048047100) Department of Oral Biology, Faculty of Dentistry Universitas Indonesia,

Indonesia

Bramma Kiswanjaya, (SCOPUS ID = 26664624100) Department of Dentomaxillofacial Radiology, Faculty of Dentistry,

Universitas Indonesia

Erik Idrus, (SCOPUS ID = 51763635000) Department of Oral Biology, Faculty of Dentistry Universitas Indonesia, Indonesia

Sunarso, (SCOPUS ID = 55191249900) Department of Dental Materials Science, Faculty of Dentistry, Universitas Indonesia

EDITORIAL TEAM

Anne-Marie Kuijpers-Jagtman, (SCOPUS ID = 35237602100) Department of Orthodontics, University Medical Center

Log In Journal Home About This Journal

Focus and Scope Editorial Team

Editorial Policies Publication Ethics

Author Guidelines

Reviewers

Indexing and Abstracting

View Readership View Citations

Submit Article

Most Popular Papers

Receive Email Notices or RSS

Select an issue:

Receive Email Notices or RSS

Select an issue:

All Issues

Browse

Search

Enter search terms:

Search

in this journal

Advanced Search

ISSN: 1693-9697

E-ISSN: 2355-4800

VISITORS

00833367

View Statistics



This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 License

EDITORIAL TEAM

Groningen, The Netherlands

Anne-Marie Kuijpers-Jagtman, (SCOPUS ID = 35237602100) Department of Orthodontics, University Medical Center

Adrian Yap, (SCOPUS ID = 7103215953) National University of Singapore, Singapore

Benny Mulyono Soegiharto, (SCOPUS ID = 24478917200) Department of Orthodontics, Faculty of Dentistry, Universitas Indonesia. Indonesia

Anak lamaroon, (SCOPUS ID = 6701786341) Faculty of Dentisry, Chiang Mai University, Thailand, Thailand

<u>Budi Kusnoto</u>, (SCOPUS ID = 6603330742) Department of Orthodontics, College of Dentistry, University of Illinois at Chicago, United States

BSM Samadarani Siriwardena, (SCOPUS ID = 16246416800) Department of Oral Pathology, Faculty of Dental Sciences,

University of Peradeniya, Sri Lanka

Chun Hung Chu, (SCOPUS ID= 7404345729) The University of Hong Kong, Hong Kong

e-en assembles well-development processes and an experimental processes and the end of t

Clive Wright, (SCOPUS ID = 55453620500) University of Sydney, Australia

Cortino Sukotjo, (SCOPUS ID = 6508194317) Department of Restorative Dentistry, College of Dentistry, University of Illinois at Chicago, United States

<u>Diah Ayu Maharani</u>, (SCOPUS ID = 32867911000) Department of Preventive and Public Health Dentistry Faculty of Dentistry Universitas Indonesia, Indonesia

Diana Cassi, (SCOPUS ID = 56062239400) Department of Orthodontics University of Modena and Reggio Emilia, Italy

Hamid Nurrohman, Hamid Nurrohman, (SCOPUS ID = 52564067000) Department of Restorative and Prosthodontics; The University of Texas School of Dentistry, Houston, Texas, USA.



This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 License.

Diah Ayu Maharani, (SCOPUS ID = 32867911000) Department of Preventive and Public Health Dentistry Faculty of Dentistry Universitas Indonesia, Indonesia

Diana Cassi, (SCOPUS ID = 56062239400) Department of Orthodontics University of Modena and Reggio Emilia, Italy

Hamid Nurrohman, Hamid Nurrohman, (SCOPUS ID = 52564067000) Department of Restorative and Prosthodontics, The University of Texas School of Dentistry, Houston, Texas, USA.

Kobkan Thongprasom, (SCOPUS ID = 6601959811) Department of Oral Medicine, Faculty of Dentistry, Chulalongkorn

Izzet Yayuz, (SCOPUS ID = 35967243400) Faculty of Dentistry, University of Dicle, Turkey, Turkey

Koichi Kawabuchi, (SCOPUS ID = 7003523337) Tokyo Medical and Dental University, Japan

Marco Meleti, (SCOPUS ID = 55926088200) Dental School at the University of Parma, Italy

Masita Mandasari, (SCOPUS ID= 57008957900) Department of Oral Medicine, Faculty of Dentistry, Universitas Indonesia

Nieka Adhara Wahono, (SCOPUS ID= 57204112724) Department of Pediatric Dentistry, Faculty of Dentistry, Universitas Indonesia

Nina Ariani, (SCOPUS ID = 55301181500) Department of Prosthodontics, Faculty of Dentistry, Universitas Indonesia

Pham Anh Vu Thuy, (SCOPUS ID = 51562199400) University of Medicine and Pharmacy, Viet Nam



About | FAQ | My Account | Accessibility Statement

University, Thailand, Thailand

Journal of Dentistry Indonesia

Log In

Journal Home

About This Journal

Focus and Scope Editorial Team

Editorial Policies
Publication Ethics

Author Guidelines

Reviewers
Indexing and Abstracting

View Readership

Submit Article

Most Popular Papers

Receive Email Notices or RSS

Select an issue:

∨ Browse

Home > Journals > Faculty of Dentistry > JDI

Current Issue: Volume 30, Number 3 (2023)

Articles

PDF Surface Roughness of Restorative Materials After Simulated Toothbrushing with Toothpastes Containing Theobromine and Arginine: An In Vitro Study

Fikri Öcal, Burak Dayi, and Zeynep Küçükakçalı

https://doi.org/10.14693/jdi.v30i3.1438

▶ PDF Comparison of EDTA and Boric Acid Irrigation in Terms of Root Canal Dentin Microhardness and Sealer Penetration Depth Selen Înce Yusufoğlu, Esma Saricam, Neslihan Bulak, Mustafa Güngörmüs, Fatih Tulumbacı, and Suat Özcan https://doi.org/10.14693/idi.v30i3.1472

PDF Comparison and Assessment of Design, Handling Characteristics, and Efficacy of Hand Instruments in Calculus Removal Mohan Kumar Pasupuleti, Krishna Ganesh, Gautami S Penmetsa, Boddeda Anusha, Haripriva Narukurthi, and Kommina Mouna Sahith



Follow

This issue (JDI Volume 30 Issue 3 Year 2023) has been finalized and launched

Receive Email Notices or RSS Select an issue: All Issues Browse Search Enter search terms: Search in this journal Advanced Search ISSN: 1693-9697 E-ISSN: 2355-4800 VISITORS 00833368 View Statistics This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 License

Comparison and Assessment of Design, Handling Characteristics, and Efficacy of Hand Instruments in Calculus Removal Mohan Kumar Pasupuleti, Krishna Ganesh, Gautami S Penmetsa, Boddeda Anusha, Haripriya Narukurthi, and Kommina Mouna Sahith https://doi.org/10.14693/idi.v30i3.1514 Effect of Triple Antibiotic Paste, Double Antibiotic Paste, and Calcium Hydroxide on Antibiotic Resistance of Tet Repressor Proteins, Tetracycline Resistance Gene W. and Tetracycline Resistance Gene Q: A Randomized Controlled Clinical Trial Ertuğrul Karataş, Özgen Kırmızıbekmez, Mustafa Özkan Baltacı, and Ahmet Adiaüzel https://doi.org/10.14693/jdi.v30i3.1516 Evaluation of Trabecular Bone Around the Impacted Maxillary Canine on CBCT Images by Fractal Analysis Zeynep Betül Arslan and Berrin Celik https://doi.org/10.14693/jdi.v30i3.1524 Evaluation of Morphologic and Morphometric Characteristics of the Mental Foramen and Anterior Loop in a Turkish Subpopulation Using Cone Beam Computed Tomography Fatma Büşra Doğan and Faruk Akgünlü https://doi.org/10.14693/idi.v30i3.1531 Integration of Massive Open Online Course (MOOC) in Conventional Teaching and Learning for Fixed Prosthodontics - A Case Study Nor Faharina Abdul Hamid, Rohana Ahmad, Tong Wah Lim, Nur Hafizah Kamar Affendi, Tengku Fazrina Tengku Mohd Ariff, Melati Mahmud, and Siti Mariam Ab Ghani https://doi.org/10.14693/jdi.v30i3.1538 Factors Affecting the Preference of Orthodontic Appliances Among Undergraduate Students in West Jakarta Joko Kusnoto, Lia H. Andayani, and Budi Kusnoto

https://doi.org/10.14693/idi.v30i3.1590

This issue (JDI Volume 30 Issue 3 Year 2023) has been finalized and launched on 26th December 2023. All articles in this issue (11 original research articles and 4 case reports articles) include 63 Authors from 8 countries of origin (Australia, Brazil, Hong Kong SAR, India, Indonesia, Malaysia, Türkiye, and United States of America).

License.

PDF Factors Affecting the Preference of Orthodontic Appliances Among Undergraduate Students in West Jakarta Joko Kusnoto, Lia H. Andayani, and Budi Kusnoto https://doi.org/10.14693/jdi.v30i3.1590 PDF Prevalence and Clinical Characteristics of Temporomandibular Disorders in Adults: An Epidemiological Study in the Mediterranean Region of Turkiye Esra Yavuz, Selmi Yardimci, and Humeyra Tercanli https://doi.org/10.14693/jdi.v30i3.1579 PDF The Effect of Several Electric Cigarette Puffs on Nanohybrid Composite Resin Surface Roughness Alvi Rizgi Amaliyah, Haris Budi Widodo, and Irfan Dwiandhono https://doi.org/10.14693/jdi.v30j3.1508

PDF Elastic Recovery Properties of Alginate Impression Materials Based

Agus Sumono, Niken Probosari, and Sulistiyani Sulistiyani

Didin Erma Indahyani, Izzata Barid, Nafisah Nafisah, Depi Praharani,

on Red Algae Kappaphycus alvarezii

https://doi.org/10.14693/jdi.v30i3.1492

https://doi.org/10.14693/jdi.v30i3.1557

Case Reports

PDF Clinically Effective Regenerative Therapy for Primary Endodontic Lesions with Secondary Periodontal Lesions: A Case Report Martina Amalia, Budi Budi, Iceu Estu Kurmaena, Nevi Yanti, Hesty Nurcahyanti, and Aini Hariyani Nasution https://doi.org/10.14693/jdi.v30i3.1485 PDF Management of Ankyloglossia in a Non-Syndromic Patient Using a Super Pulsed Laser: A Life-Changing Treatment- A Case Report Urfi Jawed, Afaf Zia, Ramniwas Kumawat, Afshan Bey, and Maahin Mahmood

Case Reports

Clinically Effective Regenerative Therapy for Primary Endodontic Lesions with Secondary Periodontal Lesions: A Case Report Martina Amalia, Budi Budi, Iceu Estu Kurmaena, Nevi Yanti, Hesty Nurcahyanti, and Aini Hariyani Nasution

Mahmood

Imperfecta

Report

Amanda Phoon Nguyen

PDF The Effect of Several Electric Cigarette Puffs on Nanohybrid

Alvi Rizgi Amaliyah, Haris Budi Widodo, and Irfan Dwiandhono

▶ PDF Elastic Recovery Properties of Alginate Impression Materials Based

Agus Sumono, Niken Probosari, and Sulistiyani Sulistiyani

Management of Ankyloglossia in a Non-Syndromic Patient Using a Super Pulsed Laser: A Life-Changing Treatment- A Case Report Urfi Jawed, Afaf Zia, Ramniwas Kumawat, Afshan Bey, and Maahin

▶ PDF Case Report and X-ray Microtomography and Scanning Electron Microscopy Analysis of Teeth with Hypocalcified Amelogenesis

M B Rumbelsperger, and Antonio Adilson Soares de Lima

PDF Enigmatic Cervicofacial Actinomycosis Post Radiotherapy - A Case

Rodrigo W. Marinoni, Thiago G. Silva, Leonardo S. Benato, Anelize

Didin Erma Indahyani, Izzata Barid, Nafisah Nafisah, Depi Praharani,

Composite Resin Surface Roughness

https://doi.org/10.14693/jdi.v30i3.1508

on Red Algae Kappaphycus alvarezii

https://doi.org/10.14693/jdi.v30i3.1557

https://doi.org/10.14693/idi.v30i3.1485

https://doi.org/10.14693/jdi.v30i3.1492

https://doi.org/10.14693/jdi.v30i3.1438

https://doi.org/10.14693/jdi.v30i3.1589

doi: 10.14693/jdi.v30i3.1590

ORIGINAL ARTICLE

Factors Affecting the Preference of Orthodontic Appliances Among Undergraduate Students in West Jakarta

Joko Kusnoto^{1*}, Lia Hapsari Andayani², Budi Kusnoto³

- ¹Department of Orthodontics, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia
- ²Department of Dental Public Health and Preventive Dentistry, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia
- ³Department of Orthodontics, Faculty of Dentistry, University of Illinois, Chicago, USA

ABSTRACT

Malocclusion's negative impact is not only limited to dentofacial appearance but also related to quality of life. Nowadays, orthodontic treatment can be done using conventional fixed appliances, or removable appliances called clear aligners. The preference for orthodontic appliances can be affected by several factors, including patient characteristics, self-perception, psychosocial, and oral-disorder factors. **Objective:** This study aims to evaluate the factors affecting the preference for orthodontic appliances among undergraduate students in West Jakarta. **Methods:** In this cross-sectional study, 244 undergraduate students were retrieved by purposive sampling from 7 universities located in West Jakarta. A self-administered questionnaire was used to collect data, and logistic regression analysis was carried out. **Results:** Fixed orthodontic appliances were commonly used (79.5%), and the majority of treatment providers were orthodontists (64.8%). Students' belief that poor dental alignment is related to decreased social attractiveness was significantly associated with the preference for orthodontic appliances (p = 0.026; 95% CI = 1.170-11.317). Orthodontic treatment provider was significantly associated with a preference for orthodontic appliances (p = 0.000; 95% CI = 0.678-3.156). **Conclusion:** Fixed orthodontic appliances still become the main preference for malocclusion treatment among undergraduate students in West Jakarta. Besides that, the qualification of orthodontic treatment providers affects the preference for orthodontic appliances among undergraduate students in West Jakarta.

Key words: clear aligners, fixed orthodontic appliances, malocclusion, orthodontic treatment, preference

How to cite this article: Kusnoto J, Andayani LH, Kusnoto B. Factors affecting the preference of orthodontic appliances among undergraduate students in West Jakarta. J Dent Indones. 2023;30(3): 212-218

INTRODUCTION

Malocclusion is considered as the misalignment of teeth and/or jaws that results from disorders in the dynamic process of craniofacial growth and development. This condition may cause psychosocial problems due to impaired dentofacial aesthetics such as difficulties of social interaction or disturbances in oral functions, such as mastication, swallowing and speech. Untreated malocclusions are connected to an increased negative impact on the health profile and psychosocial aspects, while good dental appearance can lead to a better social function. Malocclusion can be treated using orthodontic appliances, that is best carried out by an orthodontist in order to provide appropriate treatment.

The upsurge in orthodontic treatment demand continuously rising based on awareness to improve

dentofacial appearance and quality of life. Currently, orthodontic appliances are known to be divided into conventional fixed appliances (braces) or types of removable appliances commonly known as clear aligners.³ Clear aligners was firstly introduced in 1997 (Invisalign, Align Technology, Santa Clara, CA) and have been evolving in material, design and software simulation.4 Clear aligners provide more esthetic and comfortable treatment experience, simplify oral hygiene treatment, causing less pain as well as emergency visits as compared to fixed appliances. Despite the benefits and increased number of cases treated, it is improbable for clear aligners to treat all types of malocclusions. Clear aligners may be appropriate for mild to moderate crowding cases, but vigilance still needed in more complex cases.5 Clear aligners is associated with shorter treatment, better periodontal health status, and lower risk of

^{*}Correspondence e-mail to: joko.k@trisakti.ac.id

root resorption, yet there were still indistinct results regarding pain.⁶

Orthodontic treatment uptake might be influenced by many predictors such as gender, socioeconomic class, type of financial responsibility, self-perception, and dissatisfaction with dental appearance.7 Family income, cause of the problem, and self-esteem were also important attributes influencing patient decision to have orthodontic treatment.8 Although many patients take personal appearance into consideration, preference of orthodontic appliance is also determined by financial issues. Previous studies found that there are statistically significant differences in regard to pain, psychological discomfort, psychological disability, social ability and oral health quality of life between different groups of orthodontic treatment.9,10 Less psychological discomfort, functional limitation, and physical disability were found in clear aligners users, though income level may impact pain perception and quality of life in populations with different socioeconomic status.9

Preference for orthodontic appliances can also be associated with certain outcomes, including sleeping, absences for work or school, difficulties in daily psychosocial improvement, social performance, and concentration during work or studies. Preference factors such as attractiveness, acceptability, or monetary value may also vary across different populations. Although previous studies have examined the efficacy and efficiency of treatment using fixed appliances compared to clear aligners, yet there are still lack of findings regarding the characteristic or self-perception of user. This study aimed to evaluate the characteristic, self-perception, psychosocial and oral disorder factors affecting preference of orthodontic appliance among undergraduate students in West Jakarta.

METHODS

This cross-sectional study was conducted in October until December 2022. The seven main universities in West Jakarta based on the number of students body were included in the study population. In accordance with the minimum sample size calculation, the study sample consists of 244 undergraduate students receiving clear aligners or fixed orthodontic appliance, taken by purposive sampling. Following their informed consent, participants were asked to complete a validated and modified version of a previously published questionnaire¹³ by completing the Google Forms (Google LLC, Mountain View, CA). Self-administered questionnaire that was sent through social media (WhatsApp or Instagram direct messenger) required participants to send photo of their

orthodontic appliances. If there are any questions that the participant is unable to understand, the investigator is on hand to assist them throughout the process.

The questionnaire was organized into 4 sections representing characteristic of students, self-perception, psychosocial, and oral disorder factors. Characteristic of students consist of age, gender, study major, parents' educational level, parents' monthly income, provider of orthodontic appliances, first person who noticed the teeth misalignment, and suggested orthodontic treatment. This study was approved by the Ethics Committee of the Faculty of Dentistry Universitas Trisakti with the ethical clearance No.589/S1/KEPK/FKG/8/2022.

Data analysis

The statistical analysis was performed using the Statistical Package of the Social Sciences 24.0 (SPSS Inc., Chicago, IL). Descriptive statistics was obtained for all measured variables. Logistic regression was carried out to determine the preference of orthodontic appliance based on student' characteristic, self-perception, psychosocial, and oral disorder factors. A level of significance (p < 0.05; 95% CI) was used for the statistical test.

RESULTS

The distribution of students' characteristics was presented in Table 1. The majority of the students were female (71.7%), aged older than 20 years (65.2%), and undergoing study in non-medical faculty (63.5%). Based on parental educational level, 69.7% of the fathers and 68.4% of the mothers had completed at least a diploma degree or higher. Based on parental monthly income, 86.1% of the fathers and 47.5% of the mothers had monthly income higher than IDR 3,500,000. Most of the students used fixed orthodontic appliances (79.5%) and selected orthodontists as the providers of orthodontic treatment (64.8%). As much as 43% of students realized that they had teeth misalignment prior to treatment, while orthodontic treatment was mostly suggested by family, friends, or dentists (77.9%).

Logistic regression between self-perception factors and preference for orthodontic appliances was presented in Table 2. Student' awareness regarding their poor dental arrangement (p = 0.222; 95% CI = 0.582-10.315), their dissatisfaction of dental arrangement (p = 0.227; 95% CI = 0.743-3.495), the expectation that their appearance will improve after orthodontic treatment (p = 0.646; 95% CI = 0.182-15.581), as well as their self-confidence (p = 0.805; 95% CI = 0.436-2.917), were not significantly associated with their preference of orthodontic appliance.

Table 1. Characteristic of undergraduate students using orthodontic appliance in West Jakarta

Variables	n	%
Gender		
Female	175	71.7
Male	69	28.3
Age		
≤ 20 years	85	34.8
>20 years	159	65.2
Faculty		
Medical	89	36.5
Non-Medical	155	63.5
Father's education level		
Low - Intermediate	74	30.3
High	170	69.7
Mother's education level		
Low - Intermediate	77	31.6
High	167	68.4
Father's monthly income		
≤IDR 3,500,000	34	13.9
>IDR 3,500,000	210	86.1
Mother's monthly income		
≤IDR 3,500,000	128	52.5
>IDR 3,500,000	116	47.5
Provider of orthodontic appliances		
General Dentist	86	35.2
Orthodontist	158	64.8
First person who noticed teeth misalignment		
Myself	105	43.0
Others (family/friends/dentists)	139	57.0
First person who suggested orthodontic treatment		
Myself	54	22.1
Others (family/friends/dentists)	190	77.9
Type of orthodontic appliance used		
Fixed Orthodontic	194	79.5
Clear Aligners	50	20.5

Table 2. Association between self-perception factors and preference of orthodontic appliance among undergraduate students in West Jakarta

Questions	Fixed Orthodontic	Clear Aligners	p	Exp (B)	95% CI
Do you realize having poor den	tal arrangement?				
No	23 (11.9%)	3 (6%)			
Yes	171 (88.1%)	47 (94%)	0.222	2.449	0.582-10.315
Are you satisfied with your den	tal arrangement?				
No	146 (75.3%)	36 (72%)			
Yes	48 (24.7%)	14 (28%)	0.227	1.1611	0.743-3.495
Do you think your dental and fa	acial appearance will imp	rove after orthod	lontic treati	ment?	
No	9 (4.6%)	1 (2%)			
Yes	185 (95.4%)	49 (98%)	0.646	1.684	0.182-15.581
Do you think your self-confiden	ce will improve after und	lergoing orthodoi	ntic treatmo	ent?	
No	33 (17%)	7 (14%)			
Yes	161 (83%)	43 (86%)	0.805	1.127	0.436-2.917
Total	194 (100%)	50 (100%)			

^{*}p-value from logistic regression (<0.05)

Table 3. Association between psychosocial factors and preference of orthodontic appliance among undergraduate students in West Jakarta

Questions	Fixed Orthodontic			Exp (B)	95% CI
Are you sure that you will get good	d career opportunities	after orthodonti	ic treatment	?	
No	96 (49.5%)	20 (40%)			
Yes	98 (50.5%)	30 (60%)	0.318	1.414	0.717-2.791
Do you expect good social interact	ion after orthodontic	treatment?			
No	59 (30.4%)	15 (30%)			
Yes	135 (69.6%)	35 (70%)	0.256	0.643	0.300-1.378
Do you think poor dental alignme	nt is related to decreas	sed social attract	iveness?		
No	43 (22.2%)	4 (8%)			
Yes	151 (77.8%)	46 (92%)	0.026*	3.638	1.170-11.317
Total	194 (100%)	50 (100%)			

Table 4. Association between oral disorder factors and preference of orthodontic appliance among undergraduate students in West Jakarta

Questions	Fixed Orthodontic	Clear Aligners	p	Exp (B)	95% CI
Do you have any problem while s	peaking?				
No	144 (74.2%)	35 (70%)			
Yes	50 (25.8%)	15 (30%)	0.211	1.626	0.759-3.484
Do you sense any pain/clicking so	ound around your ears	s (at the TMJ join	nt)?		
No	110 (56.7%)	34 (68%)			
Yes	84 (43.3%)	16 (32%)	0.068	0.504	0.241-1.052
Do you think your dental health	will improve after ort	hodontic treatme	ent?		
No	7 (3.6%)	3 (6%)			
Yes	187 (96.4%)	47 (94%)	0.475	0.592	0.140-2.499
Do you think your biting and che	wing will improve aft	er orthodontic tr	eatment?		
No	39 (20.1%)	7 (14%)			
Yes	155 (79.9%)	43 (86%)	0.230	1.727	0.708-4.216
Total	194 (100%)	50 (100%)			

^{*}p-value from logistic regression (<0.05)

Logistic regression between psychosocial factors and preference for orthodontic appliances was presented in Table 3. There was a significant association between awareness that poor dental alignment is related to decreased social attractiveness and preference for orthodontic appliances (p = 0.026). Students think that poor dental alignment is related to decreased social attractiveness and had higher number in the fixed orthodontic appliance group (3.638; 95% CI = 1.170-11.317).

Logistic regression between oral disorder factors and preference for orthodontic appliance was presented in Table 4. There was no significant association between any problem while speaking (p = 0.211; 95% CI = 0.759-3.484) or sense of pain/clicking sound around

the temporomandibular joint (p = 0.068; 95% CI = 0.241-1.052), with the preference of orthodontic appliance. Student' beliefs regarding their dental health improvement (p = 0.475; 95% CI = 0.140-2.499), as well as their biting and chewing improvement (p = 0.230; 95% CI = 0.708-4.216), were also not significantly associated with preference of orthodontic appliance.

Multivariate analysis between student' characteristics and preference of orthodontic appliance was presented in Table 5. Logistic regression analysis revealed that most of characteristic variables were not significant in predicting preference of orthodontic appliances. The only significant predictor for preference of orthodontic appliances was provider of orthodontic appliances (p = 0.000; 95% CI = 0.678-3.156).

Table 5. Association between personal characteristics and preference of orthodontic appliance among undergraduate students in West Jakarta

Variables	Fixed Orthodontic	Clear Aligners	Total (100%)	p	Exp (B)	95% CI
Gender						
Female	145 (82.9)	30 (17.1)	175		1	
Male	49 (71.0)	20 (29.0)	69	0.092	1.974	0.895-4.353
Age						
≤ 20 years	68 (80.0)	17 (20.0)	85		1	
>20 years	126 (79.2)	33 (20.8)	159	0.729	0.874	0.409-1.867
Faculty						
Medical	76 (85.4)	13 (14.6)	89		1	
Non-Medical	118 (76.1)	37 (23.9)	155	0.332	1.463	0.678-3.156
Father's education level						
Low - Intermediate	56 (75.7)	18 (24.3)	74		1	
High	138 (81.2)	32 (18.8)	170	0.101	0.357	0.104-1.224
Mother's education level						
Low - Intermediate	60 (77.9)	17 (22.1)	77		1	
High	134 (80.2)	33 (19.8)	167	0.827	0.871	0.254-2.988
Father's monthly income						
≤IDR 3,500,000	31 (91.2)	3 (8.8)	34		1	
>IDR 3,500,000	163 (77.6)	47 (22.4)	210	0.308	2.094	0.506-8.670
Mother's monthly income						
≤IDR 3,500,000	106 (82.8)	22 (17.2)	128		1	
>IDR 3,500,000	88 (75.9)	28 (24.1)	116	0.419	1.362	0.644-2.881
Provider of orthodontic appliances						
General Dentist	82 (95.3)	4 (4.7)	86		1	
Orthodontist	112 (70.9)	46 (29.1)	158	0.000*	11.574	0.678-3.156
First person who noticed teeth misalignment						
Myself	79 (75.2)	26 (24.8)	105		1	
Others (family/friends/dentists)	115 (82.7)	24 (17.3)	139	0.252	0.625	0.280-1.397
First person who suggested orthodontic treatment						
Myself	40 (74.1)	14 (25.9)	54		1	
Others (family/friends/dentists)	154 (81.1)	36 (18.9)	190	0.173	0.510	0.193-1.343

^{*}p-value from logistic regression (<0.05)

DISCUSSION

According to this study, the majority of undergraduate students undergoing orthodontic treatment were female, aged older than 20 years, and undergoing study in non-medical faculty. Based on parental educational level, majority had completed at least diploma degree or higher. Based on parental monthly income, majority of the fathers had high monthly income but on the contrary majority of the mothers had low monthly income. Most of students using fixed orthodontic appliance and selected orthodontist as

the provider of orthodontic treatment. Several studies showed that women have a significantly higher rate of interest in orthodontic treatment,^{7,14} and were incline to choose clear aligners as orthodontic treatment.^{15,16} Decision for orthodontic treatment is generally based on esthetic improvement, other than physical and oral health well-being. The interest and consideration in women due to orthodontic treatment may be caused by higher aesthetic standards and self-perception.^{2,15} Since family income was suspected as the main influencing factor in preference of orthodontic appliance, special attention was given to that variable by classifying the family income according to Indonesian Central Agency

on Statistics classification on Indonesian population monthly income¹⁷ and also dividing parents' income to the single income or double income.

Student awareness regarding their self-perception factors, the psychosocial aspects, and oral disorder factors were not associated with preference of orthodontic appliance. That finding is consistent with earlier research that found there is currently not enough evidence to identify a decisive element in relation to patient's preference for one orthodontic appliance over another.¹¹

Multivariate analysis showed that there was no significant association between age, study major, parents' educational level, and parents' monthly income, with preference of orthodontic appliance. This finding is in contrast with previous study that stated family income, cause of the problem, age and self-esteem were found to be the most important attributes influencing patient decision in orthodontic treatment.⁸

Based on the orthodontic treatment, most of students were using fixed orthodontic appliance. It is known that clear aligner is not effective in controlling several cases such as anterior extrusion movement or rotations. Clear aligners also require the use of other auxiliaries such as attachments, inter-arch elastics, or altered aligner geometries to improve the predictability of orthodontic movement.¹⁸ All these considerations might affect students' decision in determining treatment options. Though a prediction of interest in European Union found that Invisalign was chosen to represent all orthodontic aligners as the most searched term in Google Trends, there might be different results in developing countries.¹⁹ Majority of orthodontists currently use clear aligners in their practice stated that it was not because they believe clear aligners are more effective or more comfortable than braces or because they are more profitable, but rather to have prestige in the community and not to lag behind in technology. Therefore, it seems that fixed appliance treatment will maintain its place in orthodontic practice as an option for the near future.20

This study found a significant association between provider of orthodontic treatment with preference of orthodontic appliance. Orthodontist had become the most choices among undergraduate students in West Jakarta, and higher number of them underwent fixed orthodontic appliance. Patients with the highest level of interest in pursuing orthodontic care tend to prefer orthodontists, primarily because of treatment quality.²¹ Orthodontists were seen as being better at identifying and managing complications as well as delivering reliable results. Patients with higher educational level and incomes preferred treatment by orthodontists although they knew that they would be charged up. Preference for care by orthodontists was based on functional reasons and anticipating complications.²²

This study showed that the undergraduate students in West Jakarta self-awareness regarding malocclusion and the needs to seek orthodontic treatment is still rather low. That finding suggest that orthodontists in West Jakarta need to be more proactive in public campaign regarding awareness about malocclusion and orthodontic treatment. However, when they seek orthodontic treatment, they already have a good understanding to have the treatment done by orthodontic specialists.

The main limitation of this study was the small sample size and only taken from several universities in West Jakarta, which might be insufficient to represent the general population of undergraduate students. Further research with larger and more representative samples to evaluate the preference of orthodontic appliance is required. Other variables such as ethnicity, daily oral hygiene practice, treatment expenses, and dental health insurance, need to be investigated.

CONCLUSION

The findings in this study indicate that fixed orthodontic appliance still become the main preference for malocclusion treatment among undergraduate students in West Jakarta. Besides that, qualification of orthodontic treatment provider affecting the preference of orthodontic appliances among undergraduate students in West Jakarta.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest regarding the publication of this paper.

REFERENCES

- 1. Peres MA, Antunes JLF, Watt RG. Oral epidemiology: A textbook on oral health conditions, research topics and methods. Switzerland: Springer Nature Switzerland AG; 2021.
- Saccomanno S, Saran S, Laganà D, Mastrapasqua RF, Grippaudo C. Motivation, perception, and behavior of the adult orthodontic patient: A survey analysis. Biomed Res Int. 2022; 2022:2754051.
- 3. Yovanka V, Kusnoto J, Andayani LH. Characteristic of orthodontic appliance users based on demographics, self-self-perception factors, psychosocial, and oral disorders (Study among undergraduate students in West Jakarta). J Indones Dent Association. 2023; 6(1):9-14.
- Borda AF, Garfinkle JS, Covell DA, Wang M, Doyle L, Sedgley CM. Outcome assessment of orthodontic clear aligner vs fixed appliance treatment in a teenage population with mild

- malocclusions. Angle Orthod. 2020; 90(4):485-90.
- 5. Tamer İ, Öztaş E, Marşan G. Orthodontic treatment with clear aligners and the scientific reality behind their marketing: A literature review. Turk J Orthod. 2019; 32(4):241-6.
- Yassir YA, Nabbat SA, McIntyre GT, Bearn DR. Clinical effectiveness of clear aligner treatment compared to fixed appliance treatment: An overview of systematic reviews. Clin Oral Investig. 2022; 26(3):2353-70.
- 7. Badran SA, Al-Khateeb S. Factors influencing the uptake of orthodontic treatment. J Public Health Dent. 2013; 73(4):339-44.
- Fleming PS, Colonio-Salazar F, Waylen A, Sherriff M, Burden D, O Neill C, Ness A, Sandy J, Ireland T. Factors affecting willingness to pay for NHSbased orthodontic treatment. Br Dent J. 2022. Online ahead of print.
- 9. Gao M, Yan X, Zhao R, Shan Y, Chen Y, Jian F, Long H, Lai W. Comparison of pain perception, anxiety, and impacts on oral health-related quality of life between patients receiving clear aligners and fixed appliances during the initial stage of orthodontic treatment. Eur J Orthod. 2021; 43(3):353-9.
- Antonio-Zancajo L, Montero J, Albaladejo A, Oteo-Calatayud MD, Alvarado-Lorenzo A. Pain and oral-health-related quality of life in orthodontic patients during initial therapy with conventional, low-friction, and lingual brackets and aligners (Invisalign): A prospective clinical study. J Clin Med. 2020; 9(7):2088.
- 11. Ben Gassem AA. Does clear aligner treatment result in different patient perceptions of treatment process and outcomes compared to conventional/traditional fixed appliance treatment: A literature review. Eur J Dent. 2022; 16(2):274-85.
- Alansari RA, Faydhi DA, Ashour BS, Alsaggaf DH, Shuman MT, Ghoneim SH, Linjawi AI, Marghalani HY, Dause RR. Adult perceptions of different orthodontic appliances. Patient Prefer Adherence. 2019; 13:2119-28.
- Laakshmi GM. Reasons for seeking orthodontic treatment among south Indian population - A questionnaire study. J Pharm Sci Res. 2016; 8(7):692-5.

- 14. Shetty SK, Kumara V, Panicker VG. Self-perception of malocclusion and perception towards orthodontic treatment according to age in Indian population: a questionnaire study. Int J Innov Sci Res Technol. 2022; 7(11):2046-50.
- 15. Basanagouda CP, Chaitra C. Self-perception of dental appearance and apprehension towards orthodontic treatment and knowledge about clear aligners among undergraduates and interns-a questionnaire study. Sch J Dent Sci. 2023; 10(6):113-20.
- d'Apuzzo F, Perillo L, Carrico CK, Castroflorio T, Grassia V, Lindauer SJ, Shroff B. Clear aligner treatment: Different perspectives between orthodontists and general dentists. Prog Orthod. 2019; 20(1):10.
- 17. Rakasiwi LS, Kautsar A. Pengaruh faktor demografi dan sosial ekonomi terhadap status kesehatan individu di Indonesia. Kajian Ekonomi dan Keuangan. 2021; 5(2):146-57.
- Rossini G, Parrini S, Castroflorio T, Deregibus A, Debernardi CL. Efficacy of clear aligners in controlling orthodontic tooth movement: A systematic review. Angle Orthod. 2015; 85(5):881-9
- Sycińska-Dziarnowska M, Szyszka-Sommerfeld L, Woźniak K, Lindauer SJ, Spagnuolo G. Predicting interest in orthodontic aligners: A google trends data analysis. Int J Environ Res Public Health. 2022; 19(5):3105.
- Ozkan TH, Dursun D. An investigation of the opinions and preferences of orthodontists in Turkey regarding the use of clear aligners in orthodontic treatment. Anatol Clin J Med Sci. 2023; 28(1):72-80.
- 21. Olson JC, Shroff B, Carrico C, Boyle J, Lindauer SJ. Comparison of patient factors influencing the selection of an orthodontist, general dentist, or direct-to-consumer aligners. Am J Orthod Dentofacial Orthop. 2020; 157(4):526-32.e2.
- 22. Chambers DW, Zitterkopf JG. How people make decisions about whether or not to seek orthodontic care: Upstream in the treatment chain. Am J Orthod Dentofacial Orthop. 2019; 155(6):826-31.

(Received September 14, 2023; Accepted November 5, 2023)

Factors Affecting the Preference of Orthodontic Appliances Among Undergraduate Students in West Jakarta

by Joko Kusnoto


Submission date: 03-Jan-2024 10:27AM (UTC+0700)

Submission ID: 2266300612

File name: Paper_Joko_Kusnoto_-_13_SINTA_2_-_TAMBAHAN_31_DESEMBER_2023.pdf (1.34M)

Word count: 3993

Character count: 22448



Journal of Dentistry Indonesia

The Official Publication of The Faculty of Dentistry, Universitas Indonesia in Collaboration with Indonesian Dental Association

Journal of Dentistry Indonesia

Log in Journal Home About This Journal Focus and Scope Editorial Team Editorial Policies Publication Ethics Author Guidelines Reviewers Indexing and Abstracting View Readership

Subme Article

Most Popular Papers

Receive Email Notices or RSS

Select an Issue.

Home > Journals > Faculty of Denástry > JDI

Editorial Team

EDITOR IN CHIEF

Yuniardini Septorini Wimardhani, (SCOPUS ID=56784363100) Department of Oral Medicine, Faculty of Dentistry, Universitas Indonesia, Indonesia, Indonesia

SECTION EDITORS

Lisa Rinanda Amir, (SCOPUS ID = 14048047100) Department of Oral Blology, Faculty of Dentistry Universitas Indonesia, Indonesia

Bramma Kiswaniaya. (SCOPUS ID = 26684624100) Department of Dentomaxillofacial Radiology, Faculty of Dentistry, Universitas Indonesia Enk Idrus. (SCOPUS ID = 51783635000) Department of Oral Biology, Faculty of Dentistry Universitas Indonesia. Indonesia

Sunarso (SCOPUS ID = 55191249900) Department of Dental Materials Science, Faculty of Dentistry, Universitas Indonesia

EDITORIAL TEAM

Anne-Mare Kuiners Jagraan. (SCOPUS ID = 35237602100) Department of Orthodonbos, University Medical Center

Receive Email Notices or RSS

Select an issue.

All Issues V Browse

Search

Enter search terms

in this journal *

Advanced Search

ISSN: 1693-9697

E-18SN: 2355-4800

VISITORS

08833367

View Statistics



This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 License.

EDITORIAL TEAM

Anne-Mane Kuilpers-Jagiman (SCOPUS ID = 35237602100) Department of Orthodontics, University Medical Center Groningen, The Netherlands

Adrian Yap, (SCOPUS ID = 7103215953) National University of Singapore, Singapore

Anak lamaroon, (SCOPUS ID = 6701786341) Faculty of Dentisry, Chiang Max University, Thailand, Thailand

Benny Mulyono Sogginarto, (SCOPUS ID = 24478917200) Department of Orthodontics, Faculty of Dentistry, Universities Indonesia, Indonesia

Bud Kusnoto, (SCOPUS ID = 6603330742) Department of Orthodontics, College of Dentistry, University of Illinois at Chicago, United States BSM Samadarani Sinwardena, (SCOPUS ID = 16246416800) Department of Oral Pathology, Faculty of Dental Sciences, University of Peradeniya, Sri Lanka

Chun Hung Chu, (SCOPUS ID= 7404345729) The University of Hong Kong, Hong Kong

Cine Whight (SCOPUS ID = 55453620500) University of Sydney, Australia

Cortino Sukollo, (SCOPUS ID = 6508194317) Department of Restorative Dentistry, College of Dentistry, University of Illinois at Chicago, United States Dah Ayu Maharani, (SCOPUS ID × 32867911000) Department of Preventive and Public Health Dentistry Faculty of Dentistry Universitas Indonesia, Indonesia

Diana Cassi. (SCOPUS ID = 56062239400) Department of Orthodontics University of Modena and Reggio Emilia, Italy

Hamid Nurrohman, Hamid Nurrohman, (SCOPUS ID = \$2564067000) Department of Restorative and Prosthodontics, The University of Texas School of Dentistry, Houston, Texas, USA,



This work is licensed under a Creative Commons Attribution-ShareAlike 4.0

Ucense.

Diah Ayu Maharani, (SCOPUS ID = 32867911000) Department of Preventive and Public Health Dentistry Faculty of Dentistry Universitas Indonesia, Indonesia

Diana Cassi (SCOPUS ID = 56062239400) Department of Orthodoniics University of Moderna and Reggio Emilia, Italy

Hamid Nurrohman, Hamid Nurrohman, (SCOPUS ID = 52564067000) Department of Restorative and Prosthodontics, The University of Texas School of Dentistry, Houston, Texas, USA

ZZELYRYLZ (SCOPUS ID * 35967243400) Faculty of Dentistry, University of Dicle, Turkey, Turkey

Kobkan Thongorasom. (SCOPUS ID = 660/1959811) Department of Oral Medicine, Faculty of Dentistry, Chulalongkorn University, Thailand, Thailand

Kolchi Kawabuchi, (SCOPUS ID = 7003523337) Tokyo Medical and Dental University, Japan

Marco Meleti. (SCOPUS ID = 55926088200) Dental School at the University of Parma, Italy

Masta Mandasari, (SCOPUS ID: 57008957900) Department of Oral Medicine, Faculty of Dentistry, Universitas Indonesia

Neisa Adhara Wahang, (SCOPUS ID= 57204112724) Department of Pediatric Dentistry, Faculty of Dentistry, Universitas

NINE Ariani, (SCOPUS ID = 55301181500) Department of Prosthodontics, Faculty of Dentistry, Universitas Indonesia

Pham Anh Vu Thuy, (SCOPUS ID = 51552199400) University of Medicine and Pharmacy, Viet Nam



Digital Commons

Home | About | FAQ | My Account | Accessibility Statement

Copyright Privacy

Journal of Indonesia Dentistry

Journal Home

About This Journal

Editorial Team

Focus and Scope

Editorial Policies

Publication Ethics

Author Guidelines

Reviewers

Indexing and Abstracting

View Readership

View Citations

Submit Article

Receive Email Notices or ASS

Most Popular Papers

Select an Issue

All Issues

Srowse >

Home > Journals > Faculty of Dentistry > JDI

Current Issue: Volume 30, Number 3

Follow

Articles

(2023)

Toothbrushing with Toothpastes Containing Theobromine and Surface Roughness of Restorative Materials Affer Simulated Arginine: An In Vitro Study A POF

Fikn Ocal, Burak Dayi, and Zeynep Küçükakçalı https://doi.org/10.14893/jdi.v30i3.1438

A POF

Selen Ince Yusufoğlu, Esme Sancam, Neslihan Bulak, Mustafa Comparison of EDTA and Bork: Acid Irrigation in Terms of Root Canal Dentin Microhardness and Sealer Penetration Depth Güngörmüs, Fatih Tulumbacı, and Suat Occan https://doi.org/10.14893/jdi.v30/3.1472

A POF

Boddeda Anusha, Harpriya Narukurth, and Kommina Mouna Sahith Mohan Kumar Pasupuleti, Krishna Ganesh, Gautami S Penmetsa, Comparison and Assessment of Design, Handling Characteristics. and Efficacy of Hand Instruments in Calculus Removal



Indonesia Dentistry

This issue (JDI Volume 30 issue 3 Year 2023) has been finalized and launched

Receive Email Notices or RSS

Select an Issue:

All Issues Y Browse

Search

Enter search terms.



Advanced Search

SSN: 1693-9697

E-188N: 2355-4800

VISITORS

88833388

View Statistics



This work is licensed under a Creative Commons Attribution-ShareAlike 4.0

License

A PDF Comparison and Assessment of Design, Handling Characteristics, and Efficacy of Hand Instruments in Calculus Removal Mohan Kumar Pasupulett, Krishna Ganesh, Gautamy S Penmetra, Bodoleda Anusha, Harpinya Narukurthi, and Kommina Mouna Sahith https://doi.org/10.14693/jdi.y30i3.1514

A PDF Effect of Triple Antibiotic Paste, Double Antibiotic Paste, and Calcium Hydroxide on Antibiotic Resistance of Tel Repressor Proteins, Tetracycline Resistance Gene W, and Tetracycline Resistance Gene O. A. Randomized Controlled Clinical Trial Enugrui Karakas, Ozgen Kirmizibekmez, Mustafa Ozkan Balfaci, and Anmet Adiguzei.

App. Evaluation of Trabecular Bone Around the Impacted Maxillary Canine on CBCT Images by Fractal Analysis
Zeymen Bettil Arsian and Berrin Çelik
https://doi.org/10.14693/jdi.v3003.1524

A PDF. Evaluation of Morphologic and Morphometric Characteristics of the Mental Foramen and Anterior Loop in a Turkish Subpopulation Using Cone Beam Computed Tomography. Fatma Büşra Doğan and Faruk Akgünü https://doi.org/10.14693/jdi.v30i3.1531

A PDE. Integration of Massive Open Online Course (MOOC) in Conventional Teaching and Learning for Fixed Prosthodoritics - A Case Study. Nor Faharina Abdul Hamid. Rohana Ahmad. Tong Wah Lim. Nur. Haftzah Kamar Affendi. Tengku Fazinna Tengku Mohd Anff, Melati Mehmud, and Str. Marlem Ab Ghani https://doi.org/10.14693/jdi.v3013.1538

▲ PDF Factors Affecting the Preference of Orthodordic Appliances Among Undergraduate Students in West Jakarta Joko Kusnoto Lie H. Andeyavir, and Budi Kusnoto nttps://doi.org/10.14693/jdi.v30i3.1590

This issue (JDI Volume 30 Issue 3 Year 2023) has been finalized and launched on 26th December 2023. All articles in this issue (11 original research articles and 4 case reports articles) include 53 Authors from 8 countries of origin (Australia, Brazil, Hong Kong SAR, India, Indonesia, Malaysia, Türkiye, and United States of America).

PDE Factors Affecting the Preference of Orthodortic Appliances Among Undergraduate Students in West Jakarta Joko Kusnoto, Lie H. Andayan, and Budi Kusnoto https://doi.org/10.14693/jdi.v30l3.1590

A PDF. The Effect of Several Electric Cigarette Puffs on Nanohybrid Composite Resin Surface Roughness
Avv Rzgr. Amakyart, Hans Budi Widodo, and Irfan Dwiandhono https://doi.org/10.14693/jdi.v30/3.1508

PDF. Elastic Recovery Properties of Alginate Impression Materials Based on Red Algae Kappaphycus alvarezii Didin Erma Indehyani Izzata Band, Nafisah Nafisah, Depi Praharani, Agus Sumono, Niken Probosan, and Sulistiyani Sulistiyani https://doi.org/10.14693/jdiv30i3.1557

Case Reports

A POF Clinically Effective Regenerative Therapy for Primary Endodontic Lesions with Secondary Periodontal Lesions. A Case Report Martina Amalia, Budi Budi, Ibeu Estu Kurmaena, New Yand, Heaty Nurcehyanti, and Aur Hanyani Nasuton https://doi.org/10.14693.jdi.v30i3.1485. A PDF. Management of Ankylogiossia in a Non-Syndromic Patient Using a Super Pulsed Laser ALrife-Changing Treatment: A Case Report Urfl Jawed, Afaf Zia, Ramawas Kumawat, Afshan Bey, and Mashin Mahmood https://doi.org/10.14693/jdi.y30i3.1492 APDF The Effect of Several Electric Cigarette Puffs on Nanohybrid Composite Realin Surface Roughness Anv Rizov Amalyah, Hans Buck Widodo, and Irlan Diviandhono https://doi.org/10.14693/jdiv/30i3.1508

PDF Elastic Recovery Properties of Alginate Impression Materials Based on Red Algae Kappaphycus alvarezii Didrin Erme Indahyani, 122ata Band, Nafisah Nafisah, Depi Praharani, Agus Sumono, Niken Probosen, and Sulistiyani Sulistiyani https://doi.org/10.146930fi.v30i3.1557

Case Reports

A PDE Clinically Effective Regenerative Therapy for Primary Endodomtic Lesions with Secondary Periodomial Lesions. A Case Report Martina Amalia. Budi Budi, Iceu Esta Kurmaena, New Yanti, Hesty Nurcahyanti, and Aim Harryani Nasution https://doi.org/10.14893/jdi.v30i3.1485 A PDF Management of Ankylogiossia in a Non-Syndromic Patient Using a Super Pulsed Laser A Life-Changing Treatment- A Case Report Lift Jawed, Afaf 2ig, Ramniwas Kumawat, Afshan Sey, and Mashim Mahmood https://doi.org/10.14893/di.v30i3.1492

A PDE Case Report and X-ray Microtomography and Scanning Electron
Microscopy Analysis of Teeth with Hypocalcified Amelogenesis
Imperfecta
Rodrigo W Merinoni, Thiago G. Silva, Leonardo S. Benato, Anelize
M. B. Rumbelsperger, and Antonio Adilson Spares de Lima
https://doi.org/10.14693/jdi.v30/3.1438

A PDF Enigmatic Cervicotadial Actinomycosis Post Radiotherapy - A Case Report
Amanda Phoon Ngojen
https://doi.org/10.14693/jdi.y30i3.1589

ORIGINAL ARTICLE

Factors Affecting the Preference of Orthodontic Appliances Among Undergraduate Students in West Jakarta

Joko Kusnoto", Lia Hapsari Andayani2, Budi Kusnoto3

Department of Orthodontics. Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia

Department of Dental Public Health and Preventive Dentistry, Faculty of Dentistry, Universitas Trisakti, Jakawa, Indonesia

Department of Orthodontics, Faculty of Destistry, University of Illinois, Chicago, USA

*Correspondence e-mail to: Joko k@trisakti.ac.id

ABSTRACT

Malocclusion's negative impart is not only limited to dentoficial appearance but also related to quality of life, Nowadays, orthodontic treatment, can be done using conventional fixed appliances, or removable appliances called clear aligners. The preference for orthodontic appliances can be affected by several factors, including patient characteristics, self-perception, psychosocial, and oral-disorder factors. Objective: This study aims to evaluate the factors affecting the preference for orthodontic appliances among undergraduate students in West Jakara. Methods: In this cross-sectional study, 244 undergraduate students were retrieved by purposive sampling from 7 universities located in West Jakarta. A self-administered questionnaire was used to collect data, and logistic regression analysis was carried out. Results: Fixed orthodontic appliances were commonly used (79,7%), and the majority of treatment providers were orthodontics (64.8%). Students' belief that poor dental alignment is related to decreased social attractiveness was significantly associated with the preference for orthodontic appliances (p = 0.026; 95% Cl = 1.170-11.317). Orthodontic treatment provider was significantly associated with a preference for orthodontic appliances (p = 0.000; 95% Cl = 0.678-3.156). Conclusion: Fixed orthodontic appliances still become the main preference for malocclusion treatment among undergraduate students in West Jakarta. Besides that, the qualification of orthodontic treatment providers affects the preference for orthodontic appliances among undergraduate students in West Jakarta.

Key words: dear aligners, fixed orthodontic appliances, malocclusson, orthodontic treatment, preference

How to cite this article: Kusnoto J. Andayasi LH, Kusnoto B. Factors affecting the preference of orthodontic appliances among undergraduate students in West Jakarta. J Dent Indones. 2023;30(3): 212-218

INTRODUCTION

Malocelusion is considered as the munigiment of teeth and or jaws that results from disorders in the dynamic process of craninficial growth and development. This condition may cause psychosocial problems due to impaired demofacial aesthetics such as difficulties of social interaction or disturbances in oral functions, such as mastication, swalkowing and speech! Untreated malocelusions are connected to an increased negative impaction the health profile and psychosocial aspects, while good dental appearance can lead to a better social function. Malocelusion can be treated using orthodomic appirances, that is best carried out by an orthodomic in order to provide appropriate treatment.

The upsurge in orthodontic treatment demand communusly rising based on awareness to improve

dentofacial appearance and quality of life. Currently, orthodoutic appliances are known to be divided into conventional fixed appliances (braces) or types of removable appliances commonly known as clear aligners.3 Clear aligners was firstly introduced in 1997 (Invisalign, Align Technology, Santa Clara, CA) and have been evolving in material, design and software simulation.4 Clear aligners provide more esthetic and comfortable treatment experience, simplify oral hygiene treatment, causing less pain as well as emergency visits as compared to fixed appliances. Despite the benefits and increased number of cases treated, it is improbable for clear aligners to treat all types of malocclusions. Clear aligners may be appropriate for mild to moderate crowding cases, but vigilance still needed in more complex cases.3 Clear aligners is associated with shorter treatment, better periodontal health status, and lower risk of root resurption, yet there were still indistinct results regarding pain.⁵

Orthodontic treatment uptake might be influenced by many predictors such as gender, sociocconomic class, type of financial responsibility, self-perception, and dissatisfaction with dental appearance. Family income, cause of the problem, and self-esteem were also important attributes influencing patient decision to have orthodontic treatment. Although many patients take personal appearance into consideration, preference of orthodontic appliance is also determined by financial issues." Previous studies found that there are statistically significant differences in regard to pain, psychological discomfort, psychological disability, social ability and oral health quality of life between different groups of orthodontic treatment." Less psychological discomfort, functional limitation, and physical disability were found in clear aligners users. though income level may impact pain perception and quality of life in populations with different socieeconomic status."

Preference for orthodontic appliances can also be associated with certain outcomes, including sleeping, absences for work or school, difficulties in daily psychosocial improvement, social performance, and concentration during work or studies." Preference factors such as attractiveness, acceptability, or monetary value may also vary across different populations. Although previous studies have examined the efficacy and efficiency of treatment using fixed appliances compared to clear alignen, yet there are still lack of indings regarding the characteristic or self-perception of user. This study aimed to evaluate the characteristic self-perception, psychosocial and avaldicander factors affecting preference of orthodontic appliance among undergraduate students in West Jakarta.

METHODS

This cross-sectional study was conducted in October until December 2022. The seven main universities in West Jakarra based on the number of students hody were included in the etudy population. In accordance with the minimum sample size calculation, the study sample consists of 244 undergraduate students receiving clear aligners or fixed orthodontic appliance, taken by purposive sampling. Following their informed consent, participants were acked to complete a validated and modified version of a previously published questionnaire by completing the Google Forms (Google LLC, Mountain View, CA) Self-administered questionnaire that was sent through social media (WhotsApp or Instagram direct messenger) required participants to send photo of their

orthodontic appliances. If there are any questions that the participant is unable to understand, the investigator is on hand to assist them throughout the process.

The questionnaire was organized into 4 sections representing characteristic of students, self-perception, psychosocial, and oral disorder factors. Characteristic of students consist of age, gender, study major, parents' educational level, parents' monthly income, provider of orthodontic appliances, first person who noticed the teeth misa againest, and suggested orthodontic treatment. This study was approved by the Ethics Committee of the Faculty of Dentistry Universitas Trisakti with the ethical clearance No.589/SI/KEPK/FKG58/2022.

Data analysis

The statistical analysis was performed using the Statistical Package of the Social Sciences 24.0 (SPSS Inc., Chicago, IL). Descriptive statistics was obtained for all measured variables. Logistic regression was carried out to determine the preference of orthodomic appliance based on student characteristic, self-perception, psychosocial, and oral disorder factors. A level of significance (p < 0.05; 95% CI) was used for the statistical tast.

RESULTS

The distribution of students' characteristics was presented in Table 1. The majority of the students were female (71.7%), aged older than 20 years (65.2%), and undergoing study in non-medical faculty (63.5%). Based in parental educational level, 69.7% of the fathers and 68.4% of the mothers had completed at least a diploma degree or higher. Based on parental monthly income, 86.1% of the fathers and 47.5% of the mothers had monthly income higher than IDR 3,500,000. Most of the students used fixed orthodontic appliances (79.5%) and selected orthodontics as the providers of orthodonic treatment (64.8%). As much as 42% of students realized that they had teeth misalignment prior to treatment, while orthodontic treatment was mostly suggested by family, friends, or dentists (77.9%).

Logistic regression between self-perception factors and preference for orthodontic appliances was presented in Table 2. Student' awareness regarding their poor dental arrangement ($p=0.222, 95\% \, \text{CI}=0.582\text{-}10.315$), their dissolitification of dental arrangement ($p=0.227; 95\% \, \text{CI}=0.743-3.495$), the expectation that their appearance will improve after orthodoxitic treatment ($p=0.646; 95\% \, \text{CI}=0.182\text{-}15.381$), as well as their self-confidence ($p=0.805; 95\% \, \text{CI}=0.436\text{-}2.917$), were not significantly associated with their preference of orthodoxitic appliance.

Table 1. Characteristic of undergradume students using orthodomic appliance in West Jakarta

Variables	10	56
Gender		
Female	175	71.7
Male	69	24.3
Age		
≤20 years	85	34.8
>20 years	159	65.2
Faculty		
Modical	939	36.5
Non-Medical	158	63.5
Father's education level		
Low - Interinediate	74:	30.3
High	170	69.7
Mother's education level		
Low - Intermediate	73	31.6
High 6	167	68.4
Father's monthly income		
≤IDR 2,500,000	34	13.9
>IDR 3.500,000	210	86.1
Mother's monthly income		
≤IDR 3.500,000	128	52.5
>IDR 3,500,000	356	47.5
Provider of orthodontic appliances		
General Dentist	86	35.2
Orthodomisi	158	64.8
First person who naticed teeth misalignment		
Myself	105	43.0
Others (family/friends/dentists)	139	57.0
First person who suggested orthodontic treatment		
Myself	- 54	22.1
Others (family friends dentises)	190	77.9
Type of orthodontic appliance used		
Fixed Orthodontic	194	29.5
Clear Aligners	50	20.5

Table 2. Association between self-perception factors and preference of orthodoxic applicance among undergraduate students in West Jakarta

Questions	Fixed Orthodontic	Aligners	p	Exp(B)	95% CI
Do you realize having po	or dental arrangement?				
No	23 (11.0%)	3 (6%)			
Yes	171 (88.1%)	47 (94%)	0.222	2:449	0.582-10.315
Are you satisfied with yo	our deatal arrangement?				
Na	146 (75.3%)	36 (72%)			
2 Yes	48 (24.7%)	14 (28%)	0.227	1.1601	0.743-3.495
Do you think your denta	I and facial appearance will imp	rove after orthod	londic treat	ment?	
Na	9 (4 (6%)	1 (2%)			
2 Yes	185 (95.4%)	49 (9854)	0.646	1.684	0.182-15.581
Do you think your self-e-	onfidence will improve after a no	lergoing or tho do	otic treatme	ent?	
No	33 (17%)	7 (14%)			
Yes	168 (83%)	43 (80%)	0.805	1.127	0.436-2.917
Total	194 (106%)	30 (100%)			

^{*}p-value from logistic regression (=0.08)

Table 3. Association between psychonocial factors and preference of orthodoxic appliance among undergraduate students in World Association.

Opertions	Fixed	Clear	p	Exp (B)	95% CT	
	Orthodontic	Aligners				
Are you mure that you will ge		after orthodosti	c treatment	2		
No	96 (49,5%)	20 (40%)				
Ver	98 (50.5%)	30 (60%)	0.318	1.414	0.717-2.791	
Do you expect good social int	eraction after orthodontic	treatment?				
No	59 (30.4%)	15 (3096)				
Yes	1.35 (69.6%)	35 (70%)	0.256	0.643	0.3004.379	
Do you think pour dental alig	pament is related to decrea	sed secial attract	liveness?			
No.	43 (22.2%)	4 (8%)				
Yes	151 (77.8%)	46 (9276)	0.020+	9.038	1.170-11.31	
Total	194 (900%)	50 (100%)				

Table 4. Association between and disorder factors and preference of orthodostic appliance among undergraduate students ≡ West likerta

Questions	Fixed Orthodontic	Clear Aligners	p	Exp (B)	95% CI
Do you have any probl-	em while speaking?				
No	144 (74.2%)	35 (70%)			
2 Yes	50 (25.8%)	15 (30%)	0.211	1:626	0.750-3.484
Do you sense any pain	elicking sound around your ear	s (at the TMJ join	107		
No	110 (56.7%)	34 (58%)			
2 Yes	84 (43.3%)	15 (32%)	8.000	0.504	0:241-1.052
Do you think your den	tal health will improve after ort	hodontic treatme	nt?		
No	7 (3,6%)	3 (6%)			
2 Ves	197 (96.4%)	47 (94%)	0.475	0.592	0.140-2.499
Do you thick your bitic	ng and chewing will improve all	er or the dontie tr	eatment?		
No	39 (20.1%)	7 (14%)			
Yes	155 (79.9%)	43 (86%)	0.230	1.72.7	0.708-4-250
Total	194 (100%)	50 (100%)			

^{*}p-v due from logistic regression (+005)

Logistic regression between psychosocial factors and preference for orthodontic appliances was presented in Table 3. There was a significant association between awareness that poor dental alignment is related to decreased social attractiveness and preference for orthodontic appliances (p = 0.026). Studests think that poor dental alignment is related to decreased social attractiveness and had higher number in the fixed orthodontic appliance group (3.638; 98% CI = 1.176-11.317).

Logistic regression between oral disorder factors and preference for orthodontic appliance was presented in Table 4. There was no significant association between any problem while speaking (p = 0.211; 95% C1 = 0.759-3.484) in sease of pain/clicking sound around the temperomandibular joint (p = 0.068; 95% CI = 0.241-1.052), with the preference of orthodontic appliance. Student beliefs regarding their dental health improvement (p = 0.475, 95% CI = 0.140, 2.899), as well as their biting and chewing improvement (p = 0.230; 95% CI = 0.708-4.216), were also not significantly associated with preference of orthodontic appliance.

Multivariate analysis between ctudent' characteristics, and preference of orthodontic appliance was presented in Table 5. Logistic regression analysis revealed that most of characteristic variables were not significant in predicting preference of orthodontic appliances. The only significant predictor for preference of orthodontic appliances (p = 0.008, 95% C1 = 0.678-3.156).

Table 5. Association between personal characteristics and preference of orthodonic appliance among undergradume students in West Jakarta.

Variables	Fixed Orthodomic	Clear Aligners	Totat (100%)	P	Exp (B)	95% C1
Gender						
Fenule	145 (82.9)	39 (17.1)	175		4	
Male	49 (71,0)	20 (29.0)	69	0.092	1.974	0.895-4.353
Age						
≤ 20 years	68 (80.0)	17 (20.0)	15		21	
>20 years	126 (79.2)	33 (20.8)	159	0.729	0.874	0.409-1.867
Faculty						
Medical	76 (85.4)	17 (14.6)	10		1	
Non-Medical	118 (76.1)	37 (23.9)	155	0.332	1.463	0.678-3.156
Father's education level		050000000				
Low - Intermediate	56 (75.7)	18 (24.3)	74		1	
High	138 (81,2)	32 (18.8)	170	0.101	0.357	0.104-1.224
Mother's education level						
Low - Intermediate	60 (77.9)	11 (22.1)	37		84	
High 6	134 (80.2)	32 (19.8)	163	0.827	0.871	0.254-1.988
Father's monthly income						
SIDR 3,510,000	51 (91.2)	2 (8.8)	34		1	
>IDR 3,580,000	163 (77,6)	47 (22.4)	210	0.308	2.094	0.506-8.670
Mother's monthly income						*115000000
≤IDR 3,510,000	106 (82.8)	22 (17.2)	12.8		4	
>IDR 3,590,000	88 (75.9)	28 (24.1)	116	0.419	1362	0.644-2.891
Provider of orthodontic appliances						
General Dentist	82 (95.3)	4 (4.7)	16		1	
Orthodomist	11.2 (70.9)	46 (29.1)	158	0.000*	11.574	0.678-3.156
First person who noticed teeth misslignment						
Myself	79 (75.2)	26 (24.8)	105		1	
Others (family friendsalentist)	11.5 (82.7)	24 (17.3)	130	0.257	0.625	0.280-1.307
First person who suggested orthodoxic treatment						
Myself	40 (74.1)	14 (25.9)	54		11	
Others (family Triends de itsis)	154 (80.1)	36 (18.9)	190	0.173	0.510	0.193-1.343

^{*}p-value from logistic regression (<0.05)

DISCUSSION

According to this study, the majority of undergraduate students undergoing orthodontic treatment were female, aged older than 20 years, and undergoing study in non-medical faculty. Based on parental educational level, majority had completed at least diploma degree or higher. Based on parental monthly income, majority of the fathers had high monthly income but on the contrary majority of the mothers had low monthly income. Most of students using fixed orthodontic appliance and selected orthodontist as

the provider of orthodontic treatment. Several studies showed that women have a significantly higher rate of interest in orthodontic treatment, 15 and were incline to choose clear aligners as orthodontic treatment, 15 and becision for orthodontic treatment is generally based on eithetic improvement, other than physical and ortal health well-being. The interest and consideration in women due to orthodontic treatment may be caused by higher aesthetic standards and self-perception. 15 Since family income was suspected as the main influencing factor in preference of orthodontic appliance, special attention was given to that variable by classifying the family income according to Indonesian Central Agency

on Statistics classification on Indonesian population monthly income!" and also dividing parents" income to the single income or double income.

Student awareness regarding their self-perception factors, the psychosocial aspects, and oral disorder factors were not associated with preference of orthodontic appliance. That finding is consistent with earlier research that found there is currently not enough evidence to identify a decisive element in relation to patient's preference for one orthodontic appliance over another."

Multivariate analysis showed that there was no significant association between age, study major, parents' educational level, and parents' mouthly meome, with preference of orthodoritic appliance. This finding is in confrast with previous study that stated family meome, cause of the problem, age and self-esteem were found to be the most important attributes influencing patient decision in orthodoritic treatment."

Based on the orthodontic treatment, most of students were using fixed orthodontic appliance. It is known that clear aligner is not effective in controlling several cases such as unterior extrusion movement or rotations. Clear aligners also require the use of other auxiliaries such as attachments, inter-arch elastics, or altered aligner geometries to improve the predictability of orthodontic movement." All these considerations might affect students' decision in determining treatment options. Though a prediction of interest in European Union found that Invisalign was chosen to represent all orthodontic aligners as the most searched term in Google Trends, there might be different results in developing countries. 4 Majority of orthodontists currently use clear aligners in their practice stated that it was not because they believe clear aligners are more effective or more comfortable than braces or because they are more profitable, but rather to have prestige in the community and not to lag behind in technology. Therefore, it seems that fixed appliance treatment will maintain its place in orthodontic practice as an option for the near future."

This study found a significant association between provider of orthodontic treatment with preference of orthodontic appliance. Orthodontist had become the most choices among undergraduate students in West Jakurta, and higher number of them underwent fixed orthodontic appliance. Patients with the highest level of interest in pursuing orthodontic care tend to prefer orthodontists, primarily because of treatment quality. Orthodontists were seen as being better at identifying and managing complications as well as delivering reliable results. Patients with higher educational level and incumes preferred treatment by orthodontists although they knew that they would be charged up. Preference for case by orthodontists was based on functional reasons and unicipating complications.²²

This study showed that the undergraduate students in West Jahurta self-awareness regarding maleculusion and the needs to seek orthodontic treatment is still rather low. That finding suggest that orthodontists in West Jakarta need to be more proactive in public compatign regarding awareness about malocelusion and orthodontic treatment. However, when they seek orthodontic treatment, they already have a good understanding to have the treatment done by orthodontic specialists.

The main limitation of this study was the small sample size and only taken from several universities in West Jakarta, which might be insufficient to represent the general population of undergraduate students. Further research with Jarger and more representative samples to evaluate the preference of orthodontic appliance is required. Other variables such as ethnicity, daily oral hygiene practice, treatment expenses, and dental health insurance, need to be investigated.

CONCLUSION

The findings in this study indicate that fixed orthodontic appliance still become the main preference for malocelusion treatment among undergraduate students in West Jakarta. Besides that, qualification of orthodontic treatment provider affecting the preference of orthodontic appliances among undergraduate students in West Jakarta.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest regarding the publication of this paper.

REFERENCES

- Peres MA, Antunes JLF, Watt RG. Oral epidemiology: A textbook on oral health conditions, research topics and methods. Switzerland: Springer Nature Switzerland AG; 2021.
- Saccomanno S, Saran S, Lagana D, Mastrapisqua RF, Grippaudo C. Motivation, perception, and behavior of the adult orthodontic patient: A survey analysis. Biomed Res Int. 2022; 2022;2754051.
- Yovanka V, Kusnote J, Andayani LH. Characteristic of onhodomic appliance users based on demographics, self-self-perception factors, psychosocial, and oral disorders (Study among undergraduate students in West Jakarta). J Indones Dem Association. 2023; 6(1):9-14.
- Borda AF, Garfinkle JS, Covell DA, Wang M, Doyle L, Sedgley CM. Outcome assessment of orthodonic clear aligner vs fixed appliance treatment in a teenage population with mild

- malocdusions. Angle Orthod. 2020; 90(4):485-90.
- Tumer I, Oznaş E, Murşan G. Orthodentic treatment with clear aligners and the scientific reality behind their marketing: A liberature review. Turk J Orthod. 2019; 32(4):241-6.
- Yassir YA, Nabbat SA, McIntyre GT, Bearn DB. Clinical effectiveness of clear aligner treatment compared to fixed appliance treatment: An overview of systematic reviews. Clin Oral Investig, 2022; 26(3):2953-70.
- Badrau SA, Al-Khateeb S. Factors influencing the uptake of orthodontic treatment. J Public Health Dent. 2013; 73(4):339-44.
- Fleming PS, Colonio-Salazar F, Waylen A, Sherriff M, Burden D, O Neill C, Ness A, Sandy J, Ireland T. Factors affecting willingness to pay for N115based orthodomic treatment. Br Dent J. 2022. Online ahead of print.
- Gao M, Yan X, Zhao R, Shan Y, Chen Y, Jian F, Long H, Lui W. Comparison of pain perception, anxiety, and impacts on oral health-related quality of life between patients receiving clear aligness and fixed appliances during the initial stage of orthodontic treatment. Eur J Orthod. 2021; 43(3):333-9
- Aosenio Zanenjo L, Montero J, Albaladejo A, Oteo -Calatayud MD, Alvarado -Lorenzo A. Pain and oral-health-related quality of life in orthodontic patients during initial therapy with convertional, low-friction, and lingual brackets and aligners (Invisalign): A prospective clinical study. J Clin Med. 2020; 9(7):2088.
- Ben Gassen, A.A. Does clear aligner treatment result in different patient perceptions of treatment process and outcomes compared to conventional/ traditional fixed appliance treatment: A literature review. Eur J Dent. 2022; 16(2):274-85.
- Alansari RA, Faydhi DA, Ashour BS, Alsaggaf DH, Shuman MT, Ghoneim SH, Linjawi AL, Marghalani HY, Dause RR. Adult perceptions of different orthodontic appliances. Patient Profer Adherence 2019; 13:2119-28.
- Laakshmi GM, Reasons for seeking orthodontic treatment among south Indian population - A questionnaire study. J Pharm Sci Res. 2016; 8(7):692-5.

- Sherty SK., Kamara V, Panicker VG. Selfperception of malocolusion and perception towardsorthodontic treatment according to age in Indian population: a questionnaire study. Int J Innov Sci Res Technol. 2022; 7(1):e2046-50.
- Basanagouda CP, Chaitra C. Self-perception of dental appearance and apprehension towards orthodontic treatment and knowledge about clear aligners among undergraduates and interns-a questionnaire study. Sch J Dent Sci. 2023; 10(6):113-20.
- d'Apuzzo F, Perillo L, Carrico CK, Castrofforio T, Grassia V, Lindauer SJ, Shroff B. Clear aligner treatment: Different perspectives between orthodontists and general deatists: Prog Orthod. 2019; 20(1):10.
- Rakusiwi LS, Kautsar A. Pengaruh faktor demografi dan sosial ekonomi terhadap status kesehatan individu di Indonesia. Kajian Ekonomi dan Keuangan, 2021. 5/2/kl46-57.
- Rossini G, Parrini S, Castroflorio T, Deregibus A, Debernardi CL. Efficacy of clear aligners in controlling orthodontic tooth movement: A systematic review. Angle Orthod. 2015; 85(5):881-9
- Syemika-Deiarnowska M, Sayseka Semmerfeld L, Wożniak K, Lindauer SJ, Spagnuolo G. Predicting interest in orthodontic aligners: A google trends data analysis, Int J Environ Res. Public Health. 2022; 19(5):3105.
- Ozkun TH, Dursun D. An investigation of the opinions and preferences of orthodontists in Turkey regarding the use of clear aligners in orthodontic treatment. Anatol Clin J Med Sci. 2023; 28(1):72-80.
- Olsen JC, Shroff B, Carrico C, Hoyle J, Lindauer SJ. Comparison of patient factors influencing the selection of an orthodontist, general dentist, or direct-to-consumer aligners. Am J Orthod Demoficial Orthop. 2020; 157(4):526-32.e2.
- Chambers DW, Zitterkopf IG. How people make decisions about whether or not to seek orthodontic care: Upstream in the treatment chain. Am J Orthod Dentofacial Orthop. 2019; 155(6):826-31.

(Received September 14, 2023; Accepted November 5, 2023)

Factors Affecting the Preference of Orthodontic Appliances Among Undergraduate Students in West Jakarta

ORIGINALITY REPORT SIMILARITY INDEX **INTERNET SOURCES PUBLICATIONS** STUDENT PAPERS **PRIMARY SOURCES** discovery.researcher.life 2% Internet Source innovpub.org Internet Source Submitted to University of Ulster Student Paper link.springer.com **Internet Source** pubmed.ncbi.nlm.nih.gov 5 Internet Source Wahyu P. Nugraheni, Asri Hikmatuz Zahroh, 0% 6 Risky Kusuma Hartono, Ryan Rachmad Nugraha, Chang Bae Chun. "National Health Insurance Deficit in Indonesia: Identification of Causes and Solutions for Resolution", Global Journal of Health Science, 2020 **Publication** www.hindawi.com

Internet Source


8	Submitted to Universitas Indonesia Student Paper	1%
9	www.ncbi.nlm.nih.gov Internet Source	1%
10	David W. Chambers, Jacob G. Zitterkopf. "How people make decisions about whether or not to seek orthodontic care: Upstream in the treatment chain", American Journal of Orthodontics and Dentofacial Orthopedics, 2019 Publication	1 %
11	openaccess.leidenuniv.nl Internet Source	1 %
12	article.sciencepublishinggroup.com Internet Source	1 %
13	bmcoralhealth.biomedcentral.com Internet Source	1%
14	www.opendentistryjournal.com Internet Source	1 %
15	www.readkong.com Internet Source	1 %
16	fatcat.wiki Internet Source	<1%
17	trace.tennessee.edu Internet Source	<1%


Exclude quotes On Exclude matches < 15 words


Exclude bibliography On


Factors Affecting the Preference of Orthodontic Appliances Among Undergraduate Students in West Jakarta


GRADEMARK REPORT	
FINAL GRADE	GENERAL COMMENTS
/0	
PAGE 1	
PAGE 2	
PAGE 3	
PAGE 4	
PAGE 5	
PAGE 6	
PAGE 7	
PAGE 8	
PAGE 9	
PAGE 10	
PAGE 11	
PAGE 12	
PAGE 13	
PAGE 14	
PAGE 15	











Review by: Reviewer #1

Review date: Oct 1 2023 09:53 AM PDT

The review:

Thank you very much for inviting me to review this manuscript. I have greatly enjoyed reading this well written manuscript. The area which the authors were searching were still lacking in evidence, hence I welcome more research into this field especially considering the growing number of promoted clear aligners in the orthodontic market. At first, I was slightly concerned with regards to the sample size as well as the disproportion number between the genders. However, I was glad that the authors have acknowledged this in the Discussion section. I would gladly take this study as a pilot study for the specific population and therefore follow up study with larger sample size, especially with better gender distribution would be greatly welcomed. However, there was a very minor typo issue in the 3rd paragraphs of the Intro section (last sentence) it was written "...12saimpact...". Despite this, all are fine for me.

RESPONSE:
Typo corrected

Review by: Reviewer #2

Review date: Oct 14 2023 11:07 PM PDT

The review:

Please explain in the Discussion:

The research population includes 7 PTs in West Jakarta. Were they selected based on private PTs or State PTs? The number of students being treated with orthodontics is only 244, or are there also those who refuse?

RESPONSE:

Research population includes 5 universities in West Jakarta with largest student body. The number of 244 samples included in this study based on the minimum sample size calculation. **EXPLANATION ADDED IN METHOD**

2. Are the questions in the questionnaires regarding self-perception, psychological factors, oral disorder factors, and personal characteristics adopted from other researchers' questionnaires that have been published? or was the questionnaires created by the researcher?

RESPONSE:

The questionnaire used in this study is a modified version of questionnaire utilized in previously published study by Laakshmi **EXPLANATION ADDED IN METHOD** Muthu Laakshmi G. Reasons for Seeking Orthodontic Treatment among South Indian Population-A Questionnaire Study. J Pharm Sci & Res. 2016;8(7) \rightarrow ADD TO REFERENCES

3. Why is standard income used Rp. 3.500.000 based on what? Why separate the income of father and mother?

RESPONSE:

Central Agency on Statistics classification on Indonesian population monthly income EXPLANATION ADDED IN DISCUSSION

Rakasiwi S, Kautsar LA. Pengaruh Faktor Demografi dan Sosial Ekonomi terhadap Status Kesehatan Individu di Indonesia. J KEK. 2021;5(2):146-157) \rightarrow ADD TO REFERENCES

4. Can all the questionnaire questions be understood by the respondent, for example the question: Do you feel any pain/clicking sound around your ears (at the TMJ point)? So there is either "some kind of explanation to this question"?

When the respondent fills the questionnaire, the investigator guides the respondent if there is a question that cannot be understood by the respondent

RESPONSE:

EXPLANATION ADDED IN METHOD

5. Please explain in the discussion explaining the contents of the table, written (Table 1, 2 or 3).

RESPONSE:

EXPLANATION ADDED IN DISCUSSION

