

VOL. 45 NO. 3, DECEMBER 2022

ACCREDITED RISTEKDIKTI NO. 30/E/KPT/2018

EDITORIAL TEAM

Ensured Editor

1. Ariana Soemanto, Testing Center for Oil and Gas LEMIGAS, Indonesia

Chief Editor

1. Prof. Dr. Bambang Widarsono, National Research and Innovation Agency, Indonesia

Chief Deputy Editor

- 1. Basuki Basuki, Testing Center for Oil and Gas LEMIGAS, Indonesia
- 2. Dr. Setyo Widodo, Testing Center for Oil and Gas LEMIGAS, Indonesia

Managing Editor

- 1. Purwo Subagiyo, Testing Center for Oil and Gas LEMIGAS, Indonesia
- 2. Nurhadi Setiawan, Testing Center for Oil and Gas LEMIGAS, Indonesia

Associate Editor

- 1. Dr. Junita Trivianty, Testing Center for Oil and Gas LEMIGAS, Indonesia
- 2. <u>Dr. Humbang Purba</u>, Testing Center for Oil and Gas LEMIGAS, Indonesia
- 3. Dr. Tri Muji Susantoro, National Research and Innovation Agency, Indonesia
- 4. Dr. Cut Nanda, Testing Center for Oil and Gas LEMIGAS, Indonesia
- 5. Dr. Rachmi Kartini, Testing Center for Oil and Gas LEMIGAS, Indonesia
- 6. Dr. Christina Ani Setvaningsing, Testing Center for Oil and Gas LEMIGAS, Indonesia
- 7. Dr Riesta Anggarani, Testing Center for Oil and Gas LEMIGAS, Indonesia
- 8. <u>Dr. Usman Usman</u>, National Research and Innovation Agency, Indonesia
- 9. <u>Himawanto Himawanto</u>, Testing Center for Oil and Gas LEMIGAS, Indonesia

International Editorial Boards

- 1. Prof. Dr. Bambang Widarsono, National Research and Innovation Agency, Indonesia
- 2. Prof. Dr. Istadi M.T., University of Diponegoro, Indonesia
- 3. Prof. Dr. Maizar Rahman, Testing Center for Oil and Gas LEMIGAS, Indonesia
- 4. Prof. Dr. M. Kholil, Bogor Agricultural University/USAHID, Indonesia
- 5. <u>Prof. Septoratno Siregar, Ph.D</u>, Bandung Institute of Technology, Indonesia
- 6. Prof. M. Udiharto, Testing Center for Oil and Gas LEMIGAS, Indonesia
- 7. Prof. Dr. R. P. Koesoemadinata, Bandung Institute of Technology, Indonesia
- 8. Prof Pudji Permadi, Ph.D, Bandung Institute of Technology, Indonesia
- 9. Prof. Renanto Ph.D., Technology Institute of Surabaya, Indonesia
- 10. Dr. Anda Lucia, Testing Center for Oil and Gas LEMIGAS, Indonesia
- 11. Dr. Ulrike Schacht, Geo Marine Chemist, University of Adelaide, Australia
- 12. Dr. Robert John Morley, Palinologist and Startigrapher, Palynova, United Kingdom
- 13. Dr. Nur Suhascaryo, University of National Development "Veteran" Yogyakarta, Indonesia
- 14. Dr. Hyundon Shin, Inha University, Korea, Republic of

- 15. Dr. Suhendra Suhendra, Sunan Kalijaga State Islamic University, Indonesia
- 16. Dr. Usman Usman, National Research and Innovation Agency, Indonesia
- 17. <u>Dr. Restu Juniah</u>, University of Lampung, Indonesia
- 18. Dr. Dedy Kristanto, University of National Development "Veteran" Yogyakarta, Indonesia
- 19. Dr. Eko Budi Lelono, Palynologist, Agency for Geology, Indonesia
- 20. Dr. Mudjito Mudjito, Testing Center for Oil and Gas LEMIGAS, Indonesia
- 21. Dr. Adiwar Adiwar, Testing Center for Oil and Gas LEMIGAS, Indonesia
- 22. Dr Oberlin Sidjabat, Testing Center for Oil and Gas LEMIGAS, Indonesia
- 23. <u>Valentina Alvarez, Ph. D.</u>, Centro de Investigación en Ecosistemas de la Patagonia (CIEP), Chile
- 24. <u>Ryozo Noguchi, Ph. D.</u>, University of Kyoto, Japan
- 25. Bagas Pudji Laksono, Ph.D, Gadjah Mada University, Indonesia
- 26. Jonathan Setyoko Hadimuljono, Testing Center for Oil and Gas LEMIGAS, Indonesia
- 27. John G Kaldi, Ph.D, University of Adelaide, Australia
- 28. Dr. Nuning Vita Hidayati, University of Soedirman, Indonesia
- 29. Dr. Noor Cahyo D. Aryanto, Exploration Geology and Petrology Geochemistry, Indonesia

Language Editor

Ferry Imanuddin Sadikin, Testing Center for Oil and Gas LEMIGAS, Indonesia

Windy Permata Luvian, Testing Center for Oil and Gas LEMIGAS, Indonesia

Design Layout and OJS

Andry Hidayat, Testing Center for Oil and Gas LEMIGAS, Indonesia

Circulation

Wulandari Dianningtyas, Testing Center for Oil and Gas LEMIGAS, Indonesia, Indonesia

OJS Website Administrator

Yasdi Pramesti, Testing Center for Oil and Gas LEMIGAS, Indonesia

Photographer

Dicky Christian Fransiscus, Testing Center for Oil and Gas LEMIGAS, Indonesia

TABEL OF CONTENTS

| Articles | |
|---|------------|
| Downstreaming Buton Asphalt Into Heavy Oil Production: A Techno-Economic Analysis | <u>PDF</u> |
| Approach | |
| Danang Sismartono, Bambang Widarsono, Arie Rahmadi, Usman Usman, Wanda Ali Akbar, | 95-113 |
| Djoko Sunarjanto, Aziz M Lubad, Herizal Herizal, Atyanto D Atmoko, Nurkamelia Nurkamelia, | |
| Rudi Suhartono, Sunting Kepies | |
| Trap Prevention in Machine Learning in Prediction of Petrophysical Parameters: A Case Study | <u>PDF</u> |
| in The Field X | 115-127 |
| Adam Putra Pratama Zainuri, Pahala Dominicus Sinurat, Dedy Irawan, Hari Sasongko | |
| A Bibliometric Analysis of Carbonate Paragenesis | <u>PDF</u> |
| Moehammad Ali Jambak, Riezky Kokky Mahendra | 129-144 |
| Relative Amplitude Preservation Analysis on Interpolation Methods of The Unaliased F-K | <u>PDF</u> |
| Trace Interpolation and Regularized Nonstationary Autoregression | 145-159 |
| Wahyu Triyoso, Sunawar Kunaifi | |
| Best Practices to Achieve Optimal Geothermal Drilling Performance in A Cost-Effective | <u>PDF</u> |
| Manner: Case Study of the Fastest Geothermal Well Drilling in Java and Sumatra | 161-171 |
| Bambang Yudho Suranta, Irfan Rasyid, Akhmad Sofyan, Arif Rahutama | |

OIL AND GAS

Scientific Contributions Oil & Gas, Vol. 46. No. 3, December: 129-144

SCIENTIFIC CONTRIBUTIONS OIL AND GAS Testing Center for Oil and Gas LEMIGAS

Journal Homepage:http://www.journal.lemigas.esdm.go.id ISSN: 2089-3361, e-ISSN: 2541-0520



A Bibliometric Analysis of Carbonate Paragenesis

Moehammad Ali Jambak and Riezky Kokky Mahendra

Geological Engineering Department, Faculty of Earth Technology and Energy, Trisakti University Kyai Tapa street No. 1, Jakarta 11440, Indonesia Corresponding author: ali@trisakti.ac.id Manuscript received: July 24st, 2023; Revised: Oktober 23st, 2023 Approved: December 14st, 2023; Available online: December 23st, 2023

ABSTRACT - Carbonate rock deposits play an important role in the oil and gas industries because it is a good reservoir for hydrocarbon due to the porosity related to diagenesis and paragenesis processes which produce many pore types. Paragenesis is a mineral sequence that is formed in rocks or ore deposits. The analysis that is done to determine the paragenesis is Petrography analysis, Scanning Electron Microscopy, and Cathodoluminescence. This paper discussed the bibliometric analysis in paragenesis subjects using VOSviewer, Open Refine, and Tableau. The analysis showed a result of 987 articles published on the topic of paragenesis from 1999 until 2022 with 537 authors from 46 different countries and around 308 publications with the predefined keyword. VOS viewer could generate a few bibliometric maps such as Co-occurrence, Co-citation of authors and countries. It shows the correlation between each keyword in the bibliometric map showing topics related to paragenesis and also the relations between the country origins. It shows that paragenesis is an important subject to study in not only the ore deposits field but also needs to be used in oil and gas fields because the prospects for paragenesis in the science world are promising.

Keywords: carbonate, paragenesis, hydrocarbon, bibliometric map.

© SCOG - 2023

How to cite this article:

Moehammad Ali Jambak and Riezky Kokky Mahendra, 2023, A Bibliometric Analysis of Carbonate Paragenesis, Scientific Contributions Oil and Gas, 46 (3) pp. 129-144. DOI.org/10.29017/SCOG.46.3.1590.

INTRODUCTION

Earth's crust is mainly composed of sedimentary rocks that either turn as a form of carbonate rocks or are reduced as organic matter, the process of sedimentation burial could result in thermal reactions that produce hydrocarbons in liquid and gaseous form that usually come from organic-rich sediments with high content of carbons. That makes carbonaceous deposits one of the most important objects in oil and gas industries as well as the world's main energy sources (Birger et al., 2019). Carbonates are naturally formed as sediments or reefs mostly in marine environments with tropical and temperate seas, also as important mineral deposits. Not only that, but carbonate rocks also could be an excellent porous and permeable groundwater aquifer. That means the porosity in carbonate rocks is also an important object to analyze because we know that the fluid storage and transmission through the porosity pores could be influenced by the condition of the environment around during the burial processes of the rocks. Carbonate reservoirs may have a variation of pore system and it is necessary to understand the origin of it because from it can be collected the depositional characteristics data (Ahr, 2008). One of the factors that affects the carbonate reservoirs are diagenesis process that is usually associated with the geological structural systems such as faults and fractures with the product of strong dolomitization silicification and dissolution. Those products are also a part of the paragenetic evolution of the diagenetic processes (Lima et al., 2020).

Diagenetic processes could also affect the porosity of the reservoir resulting in multiple pore types such as vuggy or some microporosity and the different relationship between the porosity. Meanwhile, Paragenesis is a term that is sometimes used as a reference to characteristics of ore mineral assemblages, or it can also be used as a sequential formation of associated minerals (Craig et al., 1994). Meanwhile, according to Britannica Encyclopedia, Paragenesis itself is a sequence of minerals that are formed in rocks or an ore deposit. Minerals in the same ore deposit can have several variations in the temperatures and pressure changes that affect the minerals and the chemical elements from the hydrothermal solution. In particular, the location of the ore deposit has been formed by more than one period of hydrothermal activity, which makes the paragenesis sequences more complicated. Even if the decision of this paragenesis is not too vital in ore and hydrocarbon exploitation paragenesis can be an important benefit to explain the geological history of the minerals deposits and to determine when hydrocarbon migrates and is placed inside the porosity of the reservoir (Dani, 2012).

Porosity is one of the important aspects in the petrophysical study and reservoir rock because through porosity we can determine the storage capacity of the original hydrocarbon in place inside the reservoir rock (Widarsono, 2014). Porosity can be evaluated by using petrography, image analysis, and core analysis. Porosity is also heavily controlled by mechanical and chemical compaction or diagenesis processes that can result in porosity reduction (Musu, 2007). Especially limestone porosity affects the quality of the reservoir rather than cementation, whether the pore system is vuggy, fractured, or connected system limestone reservoir porosity is much more complex than sandstone porosity (Widarsono, 2011) limestone reservoir tends to be influenced not only by the tectonic but also by the dissolution process associated with sea level change and also exchanging of ions that result in recrystallization and effects from hydrothermal circulation, which is why the pore structures in limestone reservoirs are complex and extremely challenging but also a good reservoir for hydrocarbon

migration and placement (Usman, 2014).

Thin section, SEM (Scanning Electron Microscopy), and Cathodoluminescence are a very important step in analyzing the paragenetic evolution from the diagenetic processes and products. Hydrothermal flow also plays an important part in paragenesis, which could cause petroleum inclusion, and to testify the typical oil generation and migration because the hydrothermal flow, could create an alteration with a strong impact on the porosity, permeability, and heterogeneity of the host rocks. Usually carbonate reservoirs have quite an elaborate heterogeneous diagenetic evolution and interact with many fractures, which results in strong dolomitization, silicification, or dissolution. The figure below shows a paragenetic sequence interpreted from the calcite crusts in the northern Campos Basin, Brazil. Therefore, the paragenetic sequence has been divided into four stages, which are the Syngenetic, Eodiagenetic, Mesodiagenetic, and Hydrothermal phases (Figure 1) (Lima et al., 2020).

The application of paragenesis study in Indonesia is still relatively small one of the case examples is Ponorogo, East Java used the term paragenesis to point to the mineralization sequence. The analysis of the ore mineral deposit structure present in veins indicates that there are two phases of mineralization. The early phase of mineralization was related to hypogene processes meanwhile the second phase resulted in supergene enrichment processes (Table 1) (Hastuti and Wiwik, 2017).

From the case above show the use of paragenesis in the ore minerals fields it means that most publications about paragenesis in Indonesia are from the mineralogical discussion because the use of paragenesis in the oil and gas industries is still little unknown. However, apart from that, some cases discuss the use of paragenesis in the oil and gas field in Indonesia. This case example is from research with Jambak in 2016 that was done in the Miocene-aged carbonate reservoir rocks in the Northwest Java Basin. The research was done in a few formations of the Northwest Java Basin, which are the Middle Cibulakan Formation, Upper Cibulakan Formation, and Parigi Formation. Each of these formations had a different diagenesis and paragenetic sequence that is based on how the tectonic processes occur. The data used in this research were obtained from laboratory results from petrography analysis, SEM, XRD, Cathodolumination, and geochemistry analysis.

Figure 1

Paragenetic Sequences of calcite crusts in northern Campos Basin, Brazil. Thick lines show the intensity and abundance of the process, red lines show less porosity and blue lines show a porosity increase. Tha and Tho show the average homogenization temperature of primary aqueous and oil inclusions each. (Lima et al., 2020)

 Table 1

 Paragenesis in the study area, which is the Southern Mountain Range of Ponorogo, East Java (Hastuti and Wiwik, 2017)

| | | HY | POGENE (200 – 3 | 60°C) |
|--------------|-------|-------|------------------------|---------------------|
| ORE MINERAL | 400°C | 300°C | 200°C | 100°C |
| Magnetite | | | | |
| Pyrite | | | | |
| Sphalerite | | | | |
| Galena | | | | |
| Chalcopyrite | | | | _ |
| Bonite | | | | |
| Arsenopyrite | | | I. | |
| | | | | SUPERGENE <125°C |
| Covelite | | | | |
| Bornite | | | | |
| Digenite | | | | |
| Chalcosite | | | | |
| Limonite | | | | |

In the Middle Cibulakan Formation, the early start of diagenesis or the syndepositional stage is shown from the cement, which is a marine deposited cement consisting of High-Magnesium calcite with radian or fibrous cement and then followed by a non-Fe calcite with isopachous cement and in this early stage the reservoir porosity is still good-very good. The next stage which is the Shallow Phreatic stage has a few processes that happen which as Karstification, Chemical and physical compaction that happen until the end of the diagenesis stage, Recrystallization, Dolomitization and a slight hydrocarbon placing in the reservoir with the porosity poor-good. Then the last stage of diagenesis which is the burial stage, starts to form a stylolite, fractures, and much more hydrocarbon placed inside the reservoir rock with poor-good porosity and in this last stage mainly formed because of the sediment overburden.

The Upper Cibulakan Formation mostly presents cementation of the non-Fe Calcite and allochems which are formed in the early stage (Syndepositional) and with reservoir porosity of good-very good. The next stage of diagenesis which is in the Shallow Phreatic regime occurs a few processes which are karstification, chemical, and physical compaction until the last stage of diagenesis, recrystallization, dolomitization, stylolite, and some oil placing with reservoir porosity of fair-good until the end of the diagenesis process. then the last stage, which is the burial stage, formed fractures and the continuous oil placed inside the porosity.

Meanwhile, in the Parigi Formation, the diagenesis process happened in a marine environment because the marine cement formed as a non-Fe fibrous radial calcite cement as a product of the early diagenesis stage, the next stage occurred some processes such as karstification which resulted in an aragonite dissolution then occur chemical compaction that formed a non-Fe Calcite cement as a filling in porosity, then physical compaction that happened until the end of diagenesis process, recrystallization, dolomitization, and then starting a formation of stylolite and oil placing. For the last stage as the end of the diagenesis, that is last fractures and oil placing start to decrease. The porosity of the reservoir from the start is still good-very good, then for the middle of the process, it reduces to fair-good then at the last stage the porosity goes up again from poor - very good. From the explanation of each formation above it can be concluded that the quality of the porosity from good to excellent and

the permeability are affected by the depositional environment during the diagenesis process and also the tectonics that happened in the research can be one of the factors involved which resulted in oil placement inside the porosity.

METHODOLOGY

The data used in this analysis comes from articles contained in Scopus and ScienceDirect and also from a few open-access journal sites, which are systematically arranged to minimize possible errors when selecting articles according to the topic of the analysis (Deyer and Tranfield, 2009). This analysis is divided into lists of the occurrence network with the topic, country of origin, and the year the articles were published. Content Analysis is also measured quantitatively through many keywords (Ellegaard and Wallin, 2015).

The proposed method for this bibliometric analysis is structured and consists of four phases in stages which are determining the keywords or search criteria, data gathering and compilation, Data processing and refinement of the criteria, and analysis from the result to determine the development of paragenesis in the world especially in Indonesia to expand and develop the knowledge about paragenesis. The words used as search criteria and keywords are Paragenesis and Carbonate as a reference to the research topics. This part of the research allows the compilation of the basic data that will be used for the next analysis (Figure 2) (Hiatt, 2000).

RESULT AND DISCUSSION

After doing the search results about a geological term called "paragenesis", show the study on this topic has produced a total of 987 publications for the period of 1999 to 2022. The number of publications shows an increase each year in the period investigated but experienced a decline every five years in 2001, 2006, and 2010 and also a decline every three years in 2013 and 2016. The analysis reveals that 1999 was the birth period of paragenesis. This result shows that paragenesis is a scientific field with an unstable number of publications and interest every year, unlike other general fields with stable publication numbers increasing each year. Therefore, the space for paragenesis studies is still potentially large for future studies.

A Bibliometric Analysis of Carbonate Paragenesis (Moehammad Ali Jambak et al.)

Figure 2 Diagram of the methodology used in this article

Distribution of Publications in Paragenesis year 1999-2022

Publications with 23 articles then experienced a publication increase in the next year reaching the peak publication in 2017 with 80 publications but then there was a decline in publications and then experienced a slow growth of publications in the year 2018 until 2021 and this year, 2022 there is only eleven publication (Figure 3). Meanwhile the term "Carbonate Paragenesis" showed 308 publications. 2017 was the birth year period for carbonate paragenesis with 70 articles but then declined to 46 in 2018 and then slowly increased each year until it reached the peak period in 2021 with 71 articles. In 2022 right now, there is only one publication yet with the hope that it will increase again (Figure 4). Another thing that is the most important aspect to analyze is the number of citations of journals that discuss the same topic as the used keyword, which is paragenesis, and the country origin of the publications to determine which country is the most advanced and has the most publications in this study.

The figure below shows the percentage of document topics in paragenesis and citation analysis, the most cited publication was by deMelo et al with 60 citations, with the title Temporal evolution of the giant Salobo IOCG deposit, Carajás Province (Brazil): constraints from paragenesis of hydrothermal alteration and U-Pb geochronology. In this publication deMelo et al state that the paragenetic

Figure 4 Distribution of Publications in Carbonate Paragenesis year 2017-2022

Figure 5 Paragenesis topic classification and percentage pie chart

sequence represents a decreasing temperature in the hydrothermal system that appeared from the hydrolytic alteration during or after the mineralization stage and also in this paper presents the new paragenetic data for the Salobo deposit with new insight on the temporal evolution of this IOCG (Iron oxide-copper-gold) Deposits which are economically important, these Salobo deposits are believed to be associated with the alkaline emplacement and crystallization of the deposit itself (deMelo et al., 2017).

Paragenesis has been widely used in many different areas of study, the figure above shows a rough percentage of paragenesis studies in different topics from the literature used. It has been categorized into a few classifications such as minerals, hydrothermal, ore mineral, gold, and oil and gas or in carbonate. The highest percentage can be found in the mineral's topic, which is 39%, and there are 12 out of 31 publications, it can also be divided again according to the method used in each research. The methods that are usually used generally are petrographic analysis, isotope analysis or mapping, SEM (Scanning Electron Microscope), and XRD (X-ray diffraction). But the specifically used can be found in the second highest percentage which is the oil and gas or carbonate topics with 29% (9 out of 31 publications) The special method used is Cathodoluminescence (CL) shows certain minerals components from the luminescence that appeared and lastly are fluorescence analysis or UVF (Ultra Violet Fluorescence) to detect the hydrocarbons or how oil dispersed inside the sample taken which can be seen from the visible light that emitted from the sample. Next are hydrothermal topics with 13% and there are 4 publications out of 31 publications. With 10% there are ore mineral topics with three publications and then followed by gold topic which is 6% with two publications and the last one is a geochemical topic with one publication and topic with the lowest percentage which is 3%. If the other paragenesis topics are combined as one and compared with the carbonate paragenesis it shows that there are still many studies that used paragenesis in the minerals, hydrothermal, ore, and others topics with 71% percentage rather than the carbonate field with 29% percentage (Figure 5).

Figure 6 Co-occurrence bibliometric map of author keywords

In this step VOS viewer were used to determine and prepare the bibliometric map of author keywords and analyze the strong correlation between the keywords. The network resulted in 44 keywords that make up to five clusters (Figure 6) and the correlation shows through nodes and links. Keywords are represented by nodes and the bigger the node's size means that the relation with the main keyword is also very related and vice versa (Zupic and Čater, 2015). The most developed research area is Deposit in the yellow nodes where the research focuses on knowing mostly about the deposit of the energy resource. For example, Peng et al discuss in their publications the Formation mechanism of saddle dolomite hydrothermal fluid in the Middle-Upper Cambrian, he then explained that the Lower Cambrian Deposits of thick mudstone in Niutitang and Qiongzhusi Formations have multiple sets of volcanic detritus deposits which supplied the necessary material condition with fluid that are rich in ore and Mg⁺ (Peng et al., 2018). Meanwhile, another developed area that correlates with paragenesis is in the Paragenetic Sequence research area with red nodes that connected to a few other research areas such as Formation, Stage, Mineralogy, Deposits, etc. In Dongpu Depression, China showed that there is a heavy shale oil that was a product of biodegradation trapped in ferrous dolomite to know

Figure 7

Figure 8 Paragenetic sequence (Ping et al., 2020)

A Bibliometric Analysis of Carbonate Paragenesis (Moehammad Ali Jambak et al.)

Figure 9 Paragenetic sequence observed in research outcrops (Blouet et al., 2017)

when the diagenetic happened, the oil charge events are through paragenetic sequence, and provable from the result of photographic analysis in figure 6, the oil charge events were divided into 3 stages. The oil 1 charge event trapped in ferrous calcite with a similar wavelength than the next stage oil inclusions trapped in late ferrous dolomite meaning that more mature oil charge happened are grouped as oil 2 charge, the last stage is oil inclusions that primarily trapped in secondary fractures that cut through ferrous calcite and dolomite crystals this suggesting that oil 3 charge happened (Figure 7 and Figure 8) (Ping et al., 2020). Paragenetic sequence was also mentioned in a published paper written by Blouet et al in 2017, studying about biogenic gas mechanism produced by a seep carbonate paragenesis in Panoche Hills, California. The author, Blouet identified from the two outcrops analyzed located in the Moreno Formational and Cima Sandstone lentils found that there is a recurrent paragenetic sequence that could be used to evaluate the paleo fluid flow in the area. The paragenetic sequence was grouped into two stages of diagenetic, which are early and late. From the thin section analysis, Blouet et al interpreted that the seepage process from the paragenetic sequence was started by Authigenic carbonate precipitation inside a burrow that acted as a pathway for the hydrocarbon flow then coated by micrite from the mixture of methane with the surrounding sediments. From the preservation of the micrite, shows that the carbonate is deposited at a shallow depth before the compaction occurs. From Figure 8 there is a three cyclic fluid flow sequence or elementary sequence of micritic rim, botryoidal aragonite, fans of aragonite, and laminated micrite, thus indicating the early stage of diagenesis. Moving to the next stage which is the late diagenesis was showed by a formation of microsparite as an in-situ replacement of the earlier micrite, at this stage there is also a shift in the isotope number from the methane-derived carbonate into the marine carbonate signature. There are also a few changes from marine to meteoric fluids that can be seen from the isotropic trend. So, the three kinds of diagenetic fluids that could reviewer be diverse to methane, marine, and meteoric waters. The sequence event that happened during late diagenesis started with the first marine cement formation of sparite, micrite, hematite, microsparite, quartz, and ends as gypsum precipitation indicating a telogenesis stage (Figure 9 and 10) (Blouet et al., 2017).

Another publication from (Rosales et al., 2018) discusses Microfacies, diagenesis, and oil emplacement of the Upper Jurassic Arab-D carbonate reservoir in an oil field in central Saudi Arabia (Khurais Complex) also using paragenetic sequence to study the productivity of the reservoir. The result of the paragenetic sequence is obtained from a few analyses such as petrographic of microfacies, mineralogy, neomorphism or replacement, and the relationship between porosity and cementation. The study is accomplished by combining observations using petrographical, epifluorescence (UVF), and cathodoluminescence (CL) microscopes. The diagenetic and paragenetic study indicate a sequence that includes three events of dissolution, four stages of calcite cement, and two last stages of dolomite formation that involve the 5 dolomite zones. From the result of UVF (fluorescence), the presence of oil is trapped in various carbonate cements suggesting that there are about two phases of oil charging (Figure

11) (Rosales et al., 2018). VOSviewer analysis also can be used to determine the bibliometric map with the author of the article which showed the range of impact from each article, the maximum numbers parameter set to 25 and from 37,848 authors only 537 authors meet the threshold and were grouped into 8 clusters with different colors that indicating each cluster. The red Clusters have the most cited authors with 133 authors and the smallest group of nodes is cluster 8-colored brownish gray with 13 authors. (Figure 12).

This analysis of Co-citation from the country, which the articles were from, is grouped into nine clusters with different colors and there are 46 countries in total. The biggest node group is cluster one with red nodes which has nine countries, and the smallest group of nodes is cluster nine with a brownish-gray color and has around 2 countries, which are India and Denmark. Indonesia is in cluster two (blue) with six countries such as Ireland, Japan, Malaysia, Singapore, Sweden, Switzerland, and the United Kingdom. From this map, we can also see the line on each node that represents the connection between countries, the bigger the node shows that the country has made many contributions and publications in the paragenesis world and if we look closely at the Indonesia node, it shows that Indonesia had a close relation to some country which is United Kingdom, France, Sweden, and Iran but Indonesia represented by a small node which means that Indonesia still have a little publications in the paragenesis study (Figure 13).

Figure 10

Result of the petrographic analysis from the minerals. A) Elementary sequence laminated micrite indicates the early diagenesis stage. B) Late diagenetic minerals, burrow-coated micrite cross-cut by crack-filled microsparite and later by a septarian crack (Blouet et al., 2017)

Figure 11

Paragenetic sequence of diagenetic events against time for the Arab-D study oil field. There are two phase of oil charging (H1-H2) that occurred in late Cretaceous (H1) and Oligocene (H2) (Rosales et al., 2018).

Figure 12 Co-citation bibliometric map of the cited author

140 | DOI.org/10.29017/SCOG.46.3.1590

A Bibliometric Analysis of Carbonate Paragenesis (Moehammad Ali Jambak et al.)

CONCLUSIONS

This Article shows the overall summary about carbonate paragenesis obtained from secondary data, which is a few articles published years before. Graphic from distribution of publications in paragenesis showed a stable and increasing trend line that indicates a consistent publication and study about this topic. A few bibliometric, analyses that were done resulted in a few bibliometric maps, which are a Cooccurrence bibliometric map of author keywords, a Co-citation bibliometric map of the cited author, and a Co-citation bibliometric map from the network of countries. Paragenesis study especially paragenetic sequence is very connected to many study fields. There are about 987 articles published on the topic of paragenesis from 1999 until 2022 with 537 authors from 46 different countries. The statistical analysis that was done according to the secondary data resulted in a consistent trend line that is overall increasing even if there is a drop in some years, but the graphic line keeps on increasing.

This bibliographic analysis can also show the correlation between each keyword in the co-occurrence map of keywords showing topics related to paragenesis, especially paragenetic sequence. The map showed that paragenetic sequence connected to a few other keywords such as Formation, Stage, fracture, mineralogy, deposit, phase, process, and many other keywords. Another thing is that bibliometric analysis also can determine the mapping of the author's co-citation analysis of each publication and from the country it originates from and is divided into nine clusters with the biggest node, which is the United States, United Kingdom, China, and Russia. Meanwhile, Indonesia is in cluster two and mostly related to some countries such as the United Kingdom, France, Sweden, and Iran.

Based on the explanation above shows that paragenesis is an important subject to continue studying and researching. Paragenesis and diagenesis not only need to be studied in the ore deposits field, but they also need to be used in the oil and gas fields, especially to determine when the oil or hydrocarbon is placed inside the pores and also from paragenesis we can know for sure about the geological history from the mineral deposits, formed fractures, compactions, and others products formed because of the diagenesis burials. Paragenesis also growing in many countries, especially in Indonesia, which is, still little publication but has promising growth. Therefore, the paragenesis study needs to be improved and much discussed because the future of paragenesis is still wide and promising.

ACKNOWLEDGEMENT

The authors would like to thank the Faculty of Earth and Energy Technology, Trisakti University, and the entire person involved in the making of this article.

| Symbol | Definition | |
|--------------|---|--|
| | is a set of methods to quantitatively | |
| Bibliometric | analyze scientific and technological | |
| | literature. | |
| | is a salt of carbonic acid, | |
| Carbonate | characterized by the presence of the carbonate ion, CO_2^{-3} . | |
| | is changes in sediment or | |
| | sedimentary rocks during and after | |
| | rock formation, at temperatures | |
| | and pressures less than that | |
| Diagenesis | required for the formation of | |
| | metamorphic rocks or melting. It | |
| | does not include changes from | |
| | weathering. | |
| | is an organic compound consisting | |
| Hydrocarbon | entirely of hydrogen and carbon. | |
| | is an open-source desktop | |
| | application for data cleanup and | |
| | transformation to other formats, | |
| | commonly known as data | |
| OpenRefine | wrangling. Similar to spreadsheet | |
| | applications and can handle | |
| | spreadsheet file formats such as | |
| | CSV, but it behaves more like a | |
| | database. | |
| | is a petrologic concept meaning an | |
| Paragenesis | equilibrium sequence of mineral | |
| | phases. | |
| | is a software tool for constructing | |
| VOS viewer | and visualizing bibliometric | |
| v OS viewer | networks including journals, | |
| | researchers, or individuals. | |

GLOSSARY OF TERMS

REFERENCES

- Ahr, M.W., (2008), Geology of carbonate reservoirs: the identification, description, and characterization of hydrocarbon reservoirs in carbonate rocks, John Wiley & Sons, Inc., Hoboken, New Jersey.
- Birger R., Janet R. & Muhling., (2019), 'Evidence for widespread oil migration in the 1.88 Ga Gunflint Formation, Ontario, Canada', Geology, Vol. 47, No. 10, pp.899–903. Available from GeoScienceWorld [20 August 2022].
- Blouet, J., Imbert, P., and Foubert, A., (2017), 'Mechanisms of biogenic gas migration revealed by seep carbonate paragenesis, Panoche Hills', California. AAPG Bulletin, Vol. 101, No.8, pp. 1309–1340.
- Britannica, The Editors of Encyclopedia, Paragenesis. viewed 26 September (2022), from https:// www.britannica.com/science/paragenesis.
- Craig, James, R., and Vaughan, (1994), Ore Microscopy and Ore Petrography, John Wiley & Sons, Inc., USA.
- **Dani, W.I.,** (2012), Studi Alterasi dan Mineralisasi Emas Berdasarkan Analisis Petrografi Conto Inti Pemboran Aaerah Arinem, kabupaten garut, Jawa Barat. Undergraduate Thesis, Bandung Institute of Technology.
- deMelo, G., Monteiro, L., Xavier, R., Moreto, C., Santiago, E., Dufrane, S.A., Aires, B. and Santos, A., (2017), 'Temporal evolution of the giant Salobo IOCG deposit, Carajás Province (Brazil): constraints from paragenesis of hydrothermal alteration and U-Pb geochronology', Mineralium Deposita, Vol. 52. DOI: 10.1007/ s00126-016-0693-5.
- **Denyer, D. and Tranfield, D,** (2009), Handbook of Organizational Research Methods: Producing a Systematic Review, Sage Publications Ltd., Thousand Oaks, CA, USA.
- Ellegaard, O., and Wallin, J.A., (2015), 'The bibliometric analysis of scholarly production: How great is the impact?', Scientometrics Vol. 105, pp. 1809–1831.
- Hastuti, D., and Wiwik, E., (2017), 'The study of ore minerals parageneses in Ponorogo area, East Java', Sriwijaya International Conference on Engineering, Science and Technology. Avail-

able from: MATEC Web of Conferences. [16 September 2022]

- Hiatt, E., (2000), Fluids and Basin Evolution: Links between Depositional and Diagenetic Processes in Basin Analysis: Porosity and Permeability Evolution in Sedimentary Rocks. Mineralogical Association of Canada, Canada.
- Jambak, M.A., (2016), Microfacies And Diagenetic History Of Miocene Carbonate Rocks In The Onshore Region North West Java Basin, Doctoral dissertation, Universitas Padjadjaran.
- Lima, B.E.M., Tedeschi, L.R., Pestilho, A.L.S., Santos, R.V., Vazquez, J.C., Guzzo, J.V.P., Ros, L.F.D., (2020), 'Deep-burial hydrothermal alteration of the Pre-Salt carbonate reservoirs from northern Campos Basin, offshore Brazil: Evidence from petrography, fluid inclusions, Sr, C and O isotopes', Marine and Petroleum Geology. Vol. 113.
- Musu, J., (2007), 'To Explain the Nature of Core Porosity Using Results of Petrography Analysis', Lemigas Scientific Contributions, Vol. 30, No. 3, pp. 43-51.
- Peng, B., Li, G., and Li, Z., (2018), 'Discussion of multiple formation mechanisms of saddle dolomites—Comparison of geochemical data of Proterozoic–Paleozoic dolomites', Energy Exploration & Exploitation, Vol. 36, No. 1, pp. 66-96.
- Ping, H., Li, C., Chen, H., Simon, C., George, and Gong, S., (2020), Overpressure release: Fluid inclusion evidence for a new mechanism for the formation of heavy oil, Geology, Vol. 48, No. 8, pp. 803–807.
- Rosales, I., Pomar, L., and Al-Awwad, S.F., (2018), 'Microfacies, diagenesis and oil emplacement of the Upper Jurassic Arab-D carbonate reservoir in an oil field in central Saudi Arabia (Khurais Complex)', Marine and Petroleum Geology.
- Usman, (2014), 'Comparative Study of Porosity Determination Methods for OGIP in Fractured Basement Reservoir', Scientific Contributors Oil & Gas, Vol. 37, No. 2, pp. 79-86.
- Widarsono, B., (2014), 'Porosity Versus Depth Characteristics of Some Reservoir Sandstones in Western Indonesia', Scientific Contribution Oil & Gas, Vol. 37, No. 2, pp. 87-104.

- Widarsono, B., (2011), 'The Importance of Litho-Facies Distinction in Determining the Most Representative Cementation Factors for Well-Log Evaluation: An Old Issue Persistently Neglected', Scientific Contribution Oil & Gas, Vol. 34, No. 2, pp. 139-148.
- Zupic, I., and Čater, T., (2015), 'Bibliometric Methods in Management and Organization', Organizational Research Methods, Vol. 18, No. 3, pp. 429–472.

Article Test by Riezky Mahendra

Submission date: 22-Jul-2023 04:52PM (UTC+0700) Submission ID: 2134927631 File name: A_Bibliometric_Analysis_Of_Paragenesis_LEMIGAS.pdf (1.11M) Word count: 5380 Character count: 29831

A BIBLIOMETRIC ANALYSIS OF CARBONATE PARAGENESIS

ANALISIS BIBLIOMETRIK MENGENAI PARAGENESIS BATUAN KARBONAT

Moehammad Ali Jambak^{a,*}, Riezky Kokky Mahendra^a

^a Geological Engineering Department, Faculty of Earth Technology and Energy, Trisakti University, Jalan Kyai Tapa No. 1, Jakarta 11440, Indonesia. *ali@trisakti.ac.id

ABSTRAK

Endapan batuan karbonat memiliki peran yang penting dalam industri minyak dan gas dan umumnya ditemukan sebagai reservoar hidrokarbon. Porositas pada batuan reservoar berhubungan dengan proses diagenesis dan paragenesis yang menghasilkan berbagai macam tipe porositas. Paragenesis merupakan sikuen mineral yang terjadi pada suatu batuan atau endapan. Analisis yang dilakukan untuk menentukan paragenesis dapat menggunakan analisis petrografi, SEM (Scanning Electron Microscopy) dan Cathodoluminescence. Naskah ini membahas tentang analisis bibliometrik pada paragenesis dengan menggunakan VOSviewer, Open Refine, dan Tableau. Dari analisis yang telah dilakukan diperoleh 987 artikel terpublikasi dengan topik paragenesis pada tahun 1999 hingga 2022 dengan 537 penulis yang berasal dari 46 negara berbeda dan sekitar 308 publikasi mengenai paragenesis karbonat. Dengan menggunakan VOSviewer dapat diperoleh beberapa data peta bibliometrik yaitu dari penggunaan kata kunci, sitasi penulis, dan dari hubungan antar negara. Sehingga, paragenesis merupakan salah satu topik yang penting untuk dipelajari dan tidak hanya dalam bidang mineral bijih namun juga perlu untuk dipelajari pada bidang minyak dan gas serta prospek masa depan paragenesis di dunia sains cukup menjanjikan.

Kata Kunci: Karbonat, Paragenesis, Hidrokarbon, Peta Bibliometrik

ABSTRACT

Carbonate rock deposits played an important role in the oil and gas industries because it's a good reservoir for the hydrocarbon due to the porosity that related to diagenesis and paragenesis processes which produces many pore types. Paragenesis is a mineral sequence that formed in rocks or ore deposit. The analysis that are done to determine the paragenesis is Petrography analysis, SEM (Scanning Electron Microscopy) and also Cathodoluminescence. This paper discussed about the bibliometric analysis in paragenesis subject using VOSviewer, Open Refine, and Tableau. From the analysis, it showed a result of 987 articles published with the topic of paragenesis since 1999 until 2022 with 537 authors from 46 different country and around 308 publications with the predefined keyword. VOSviewer can generate a few bibliometric maps such as Co-occurrence, Co-citation of authors and countries. It shows the correlation between each key words in the bibliometric map showing topics related to paragenesis and it also the relations between the country origin. It shows that paragenesis is an important subject to study and not only in the ore deposits field, but it also needs to be used in the oil and gas fields because the future prospects for paragenesis in the science world is promising.

Keywords: Carbonate, Paragenesis, Hydrocarbon, Bibliometric map

A. INTRODUCTION

Earth crust is mainly composed of sedimentary rocks that either turn as a form of carbonate rocks

or reduced as organic matter, the process of sedimentation burial can result a thermally reactions that produces hydrocarbons in liquid and

gaseous form that usually comes from organicrich sediments with high content of carbons. That make carbonaceous deposits are one of the most important objects in oil and gas industries as well as the world's main energy sources [1]. Carbonates are naturally formed as sediments or reefs mostly in marine environment with tropical and temperate seas, also as important mineral deposits. Not only that, carbonate rocks also can be an excellent porous and permeable groundwater aquifer. That means, porosity in carbonate rocks also an important object to analyze because we can know about the fluid storage and transmission through the porosity, also pores can be influenced from the condition of the environment around during the burial processes from the rocks. Carbonate reservoirs may have a variation of pore system and it is necessary to understand the origin of it, because from it can be collected the depositional characteristics data [2]. One of the factors that affecting the carbonate reservoirs are diagenesis process that usually associated with the geological structural systems such as faults and fracture with the product of strong dolomitization silicification and dissolution. Those products also a part of the paragenetic evolution of the diagenetic processes [3].

Diagenetic processes can also be affecting the porosity of the reservoir that resulted in multiple of pore types such as vuggy or some microporosity and the difference relationship between the porosity. Meanwhile, Paragenesis is a term that sometime used as a reference to characteristic of ore mineral assemblages, or it can also be used as a sequential formation of associated minerals [4]. Meanwhile, Paragenesis itself is a sequence in minerals that are formed in rocks or in an ore deposit. Minerals in the same ore deposit can have several variations resulted due to the temperatures and pressure changes that affecting the minerals and in the chemical elements from the hydrothermal solution. In particular location if the ore deposit has been formed by more than one period of hydrothermal activity it makes the paragenesis sequences more complicated [5]. Even if the decision of this paragenesis is not too vital in ore and hydrocarbon exploitation but paragenesis can be an important benefit to explain about the geological history from the minerals deposits and to determine at when does hydrocarbon migrate and placed inside the porosity of the reservoir [6].

Porosity is one of the important aspects in petrophysical study and reservoir rock, because through porosity we can determining the storage capacity of original hydrocarbon in place inside the reservoir rock [7] Porosity can be evaluated by using petrography, image analysis and core analysis. Porosity also heavily controlled by mechanical and chemical compaction or diagenesis process that can be resulted in porosity reduction [8]

Thin section, SEM (Scanning Electron Microscopy) and Cathodoluminescence are a very important step to do in order to analyses the paragenetic evolution from the diagenetic processes and products. Hydrothermal flow also played an important part in paragenesis which can cause a petroleum inclusion and to testify the typical oil generation and migration because from the hydrothermal flow, it can create an alteration with a strong impact to the porosity, permeability, and heterogeneity of the host rocks. Usually in carbonate reservoirs have quite an elaborate heterogenous diagenetic evolution and interacted with many fractures which resulted in strong dolomitization, silicification, or dissolution. The figure below shows a paragenetic sequences interpretated from the calcite crusts in northern Campos Basin, Brazil. And so, the paragenetic sequence has divided into four stage which are Syngenetic, Eodiagenetic, Mesodiagenetic, and Hydrothermal phases (Figure 1) [9].

Figure 1. Paragenetic Sequences of calcite crusts in northem Campos Basin, Brazil. Thick lines show intensity and abundance of the process, red lines show less porosity, blue lines show porosity increase. Tha and Tho show the average homogenization temp of primary aqueous and oil inclusions each. (Lima et al, 2019)

The application of paragenesis study in Indonesia is still relatively small one of the case examples are in Ponorogo, East used the term paragenesis to point the mineralization sequence. From the analysis of the ore mineral deposit structure presents in veins indicates that there are two phases of mineralization. The early phase of mineralization related to hypogene processes meanwhile the second phase resulted in supergene enrichment processes (Table 1) [10].

Table 1. Paragenesis in the study area, which isSouthern Mountain Range of Ponorogo, East Java(Hastuti, 2017)

From the case above show the use of paragenesis in the ore minerals fields which shows that mostly publication about paragenesis in Indonesia are from the mineralogical discussion. But apart from that, there are also some cases that discuss about the use of paragenesis in the oil and gas field in Indonesia. This case example is from research with Jambak in 2016 that is done in the Miocene-aged carbonate reservoir rocks in the Northwest Java Basin. The research was done in a few formations of the Northwest Java Basin, which is Central Cibulakan Formation, Upper Cibulakan Formation, and Parigi Formation. Each of this formation have a different diagenesis and paragenetic sequence that based on how the tectonic processes occur. The data used in this research were obtained from laboratory results from petrography analysis, SEM, XRD, Cathodolumination and geochemistry analysis.

In the Central Cibulakan Formation the early start of diagenesis or the syndepositional stage is showed from the cement which is a marine deposited cement and consisted of High-Magnesium calcite with radian or fibrous cement and then followed by a non-Fe calcite with isopachous cement and in this early stage the reservoir porosity is still good-very good. Next the next stage which is the Shallow Phreatic stage have a few processes that happen which is Karstification, Chemical and physical compaction (physical compaction happened until the end of diagenesis stage), Recrystallization, Dolomitization and a slight hydrocarbon placing in the reservoir with the porosity poor-good. Then the last stage of diagenesis which is the burial stage are starting to formed a stylolite, fractures, and much more hydrocarbon placing inside the reservoir rock with poor-good porosity and in this last stage mainly formed because of the sediment overburden.

For the **Upper Cibulakan Formation** mostly present cementation of the non-Fe Calcite and allochems which is formed in the early stage (Syndepositional) and with reservoir porosity of good-very good. Next stage of diagenesis which is in the Shallow Phreatic regime occur a few processes which is karstification, chemical and physical compaction until the last stage of diagenesis, recrystallization, dolomitization, stylolite, and some oil placing with reservoir porosity of fair-good until the end of the diagenesis process. And then the last stage which is the burial stage formed fractures and the continuous oil placing inside the porosity.

Meanwhile in the **Parigi Formation** the diagenesis process happened in a marine environment because of the marine cement formed as a non-Fe fibrous radial calcite cement as a product of the early diagenesis stage, the next stage occur some process such as karstification which resulted as a aragonite dissolution then occur a chemical compaction that formed a non-Fe Calcite cement as a filling in porosity, then a physical compaction that happened until the end

of diagenesis process, recrystallization, dolomitization, and then starting a formation of stylolite and oil placing. For the last stage as the end of the diagenesis that is last fractures and oil placing starts to decrease. The porosity of the reservoir from the start are still good-very good, then for the middle of the process it reduces to fairgood then at the last stage the porosity went up again from poor-very good.

From the explanation of each formation above it can be concluded that the quality of the porosity from good to excellent and the permeability are affected by the depositional environment during the diagenesis process and also the tectonics that happened in the research are can be one of the factors involved which resulted in oil implacement inside the porosity.

B. MATERIAL AND METHOD

The data used in this analysis comes from articles contained in Scopus and Sciencedirect also from a few open access journal sites which are systematically arranged in order to minimize possible errors when selecting articles according to the topic of the analysis [11]. This analysis divided into lists of the occurrence network with the topic, country of origin, and the year the articles published. Content Analysis also measured quantitatively through many keywords [12].

The proposed method for this bibliometric analysis is structured, and consist from four phases in stages which is Determining the keywords or search criteria, data gathering and compilation, Data processing and refinement of the criteria, and analysis from the result to determine the development of paragenesis in the world especially in Indonesia in order to expand and develop the knowledge about paragenesis. The word used as a search criteria and keyword are Paragenesis and Carbonate as a reference to the research topics. This part of research is allowed compilation of the basic data that will be used for the next analysis (Figure 2) [13].

C. RESULT

After doing the search results about geological term called "paragenesis", shows the study about this topic has produced a total of 987 publications for the period of 1999 to 2022. The number of publications show an increase each year in the period investigated but experiencing a decline every five year in 2001, 2006, and 2010 also a decline every three year in 2013 and 2016. The analysis reveals that 1999 was the birth period of paragenesis

Figure 2. Diagram of the methodology used in this article

publications with 23 articles then experiencing a publication increase in the next year and reaching the peak publication in 2017 with 80 publications but then there was a decline of publication and then experiencing a slow growth of publications in the year 2018 until 2021 and in this year 2022 there are only eleven publication (Figure 3). Meanwhile with the term "Carbonate Paragenesis" showed 308 publications. 2017 were the birth year period for carbonate paragenesis with 70 articles but then decline to 46 in 2018 and then slowly increasing each year until it reaches the peak period in 2021 with 71 articles. In 2022 right now there are only one publication yet with the hope of it will increase again (Figure 4).

Another thing that is the most important aspects to analyze is the number of citations of journals that discuss the same topic as the used keyword which is paragenesis and the country origin of the publications to determine which country is the most advanced and has the most publications in this study.

Figure 4. Distribution of Publications in Carbonate Paragenesis year 2017-2022

The figure below shows percentage document topics in paragenesis and citation analysis, the most citated publication were by deMelo et al with 60 citations, with the tittle of Temporal evolution of the giant Salobo IOCG deposit, Carajás Province (Brazil): constraints from paragenesis of hydrothermal alteration and U-Pb geochronology. In this publication deMelo et al state that the paragenetic sequence represents a decreasing temperature in the hydrothermal system that appeared from the hydrolytic alteration during or after mineralization stage and also in this paper presents the new paragenetic data for the Salobo deposit with new insight on the temporal evolution of this IOCG (Iron oxide-copper-gold) Deposits which are economically important, this Salobo deposits are believed to be associated with the alkaline emplacement and crystallization of the deposit itself [14].

Figure 5. Paragenesis topic Classification and Percentage Pie Chart

Paragenesis have been widely used in many different areas of study, the figure above shows roughly percentage of paragenesis study in different topics from the literature used. It has been categorized into a few classifications such as minerals, hydrothermal, ore mineral, gold, and oil and gas or in carbonate. The highest percentage can be found in minerals topic which is 39% and there are 12 out of 31 publications, it can also be divided again according to method used in each research. The method that usually used generally are from petrographic analysis, isotope analysis or mapping, SEM (Scanning Electron Microscope) and from XRD (X-Ray Diffraction). But, the specifically used can be found in the second highest percentage which is the oil and gas or carbonate topics with 29% (9 out of 31

publications) The special method used are Cathodoluminescence (CL) that shows certain minerals component from the luminescence that appeared and lastly are fluorescence analysis or UVF (Ultra Violet Fluorescence) in order to detect the hydrocarbons or how oil dispersed inside the sample taken which can be seen from the visible light that emitted from the sample. Next are hydrothermal topic with 13% and there are 4 publications out of 31 publications. With 10% there are ore mineral topic with 3 publications and then followed by gold topic which is 6% with 2 publications and last one is geochemical topic with 1 publications and topic with the lowest percentage which is 3%. And if the other paragenesis topic are combined as one and compared with the carbonate paragenesis it shows that there are still many studies used paragenesis in the minerals, hydrothermal, ore and others topic with 71% percentage rather than the carbonate field with 29% percentage (Figure 5).

In this step VOSviewer were used to determine and prepare the bibliometric map of author keywords and analyzing the strength correlation between the keywords. The network resulted in 44 keywords that makes up to 5 clusters (in Figure 6) and the correlation shows through nodes and links. Keywords are represented by nodes and the bigger the nodes size means that the relation with the main keyword are also very related and vice versa [15].

Figure 6. Co-occurrence bibliometric map of author keywords

The most developed research area is Deposit in yellow nodes where the research focuses on knowing mostly about the deposit of the energy resource. For example, Peng et al discuss in his publications about the Formation mechanism of saddle dolomite hydrothermal fluid in the Middle-Upper Cambrian which he then explained that the Lower Cambrian Deposits of thick mudstone in Niutitang and Qiongzhusi Formations there are some areas that have a multiple set of volcanic detritus deposits which supplied the necessary material condition with fluid that are rich in ore and Mg⁺ [16].

Meanwhile, another developed area that corelates with paragenesis are in the Paragenetic Sequence research area with red nodes that

connected to a few another research area such as Formation, Stage, Mineralogy, Deposits etc. In Dongpu Depression, China showed that there is a heavy shale oil that was a product of biodegradation that trapped in ferroan dolomite to know when the diagenetic happened and the oil charge events are through paragenetic sequence and provable from the result of photographic analysis in Figure 6, the oil charge events were divided into 3 stages. The oil 1 charge event trapped in ferroan calcite with similar wavelength then the next stage oil inclusions trapped in late ferroan dolomite meaning that more mature oil charge happened are grouped as oil 2 charge, last stage are oil inclusions that primarily trapped in secondary fractures that cut through ferroan

calcite and dolomite crystals this suggesting that oil 3 charge happened (Figure 7 and 8) [17].

Figure 7. Photomicrographs analysis. A) Cathodoluminescence and B) UV light. Showed abundant fractures trapping nonfluorescent oil inclusions indicating that it was formed at late stage and the latest stage of oil charge (Ping, H. et al. 2020).

Figure 8. Paragenetic Sequence (Ping, H. et al. 2020)

Paragenetic sequence also mentioned in a published paper written by Blouet et al in 2017, studying about biogenic gas mechanism produced by a seep carbonate paragenesis in Panoche Hills, California. The author, Blouet identified from the two outcrops analysed that located in the Moreno Formationa and Cima Sandstone lentil were found that there is a recurrent paragenetic sequence happened which can be used to evaluates the paleo fluid flow in the area. The paragenetic sequence were grouped into two stages of diagenetic which is early and late. From the thin section analysis Blouet et al interpreted that the seepage process from the paragenetic sequence were started by a Authigenic carbonate precipitation inside a burrow that acted as a pathway for the hydrocarbon flow then coated by micrite from the mixture of methane with the surrounding sediments. From the preservation of the micrite it shows that the carbonate is deposited at shallow depth before the compaction occurred. From Figure 8 there is a three cyclic fluid flow sequence or elementary sequence of micritic rim, botryoidal

© 2023 R&D Centre for Oil and Gas Technology LEMIGAS All rights reserved

aragonite, fans of aragonite, and laminated micrite happened, thus indicates the early stage of diagenesis. Moving to the next stage which is the late diagenesis were showed by a formation of microsparite as a in situ replacement of the earlier micrite, at this stage there is also a shift in the isotope number from the methane derived carbonate into the marine carbonate signature. There are also a few changes from marine to meteoric fluids that can be seen from the isotropic trend. So, the three kind diagenetic fluids that can be diverse to methane, marine, and meteoric waters. The sequence event that happened during late diagenesis are started by the first marine cement formation of sparite, micrite, hematite, microsparite, quartz, and ended as gypsum precipitation indicating a telogenesis stage (Figure 9 and 10) [18].

Figure 9. Paragenetic sequence observed in research outcrops (Blouet et al, 2017)

Jambak et al. / A Bibliometric Analysis of Carbonate Paragenesis XX (20XX) XX-XX

Figure 10. Result of the petrographic analysis from the minerals. A) Elementary sequence laminated micrite indicates early diagenesis stage. B) Late diagenetic minerals, burrow coated micrite cross-cut by crack filled microsparite and later by a septarian crack (Blouet et al, 2017)

Another publication from Rosales et al, 2018 discussing about Microfacies, diagenesis, and oil emplacement of the Upper Jurassic Arab-D carbonate reservoir in an oil field in central Saudi Arabia (Khurais Complex) also using paragenetic sequence to study about the productivity of the reservoir. The result of the paragenetic sequence is obtained from a few analyses such as petrographic of microfacies, mineralogy, neomorphism or replacement, and relationship between porosity and cementation. The study is accomplished by combining observations using petrographical, epifluorescene (UVF), and cathodoluminescence (CL) microscopes. The diagenetic and paragenetic study indicate a sequence that includes three events of dissolution, four stages of calcite cement, and two last stage of dolomite formation that involve the 5 dolomite zones. From the result of UVF (fluorescence) the presence of oil is trapped in various carbonate cements suggesting that there are about two phases of oil charging (Figure 11) [19].

© 2023 R&D Centre for Oil and Gas Technology LEMIGAS All rights reserved

Figure 11. Paragenetic sequence of diagenetic events against time for the Arab-D study oil field. There are two phase of oil charging (H1-H2) that occurred in late cretaceous (H1) and oligocene (H2) (Rosales et al., 2018).

VOSviewer analysis also can be used to determine the bibliometric map with the author of the article which showed the range of impact from each article, the maximum numbers parameter set to 25 and from 37,848 authors only 537 authors meet the threshold and were grouped into 8 clusters with different colours that indicating each cluster. The red Clusters have the most citated authors with 133 author and the smallest group of nodes is cluster 8 colored brownish gray with 13 authors. (Figure 12).

This analysis of Co-citation from the country which the articles was from are grouped into nine clusters with different colours and there are 46 countries in total. The biggest nodes group are the cluster one with red nodes that has nine countries, and the smallest group of nodes are the cluster nine with brownish gray color and have around 2 countries which is India and Denmark. Indonesia is in cluster two (blue) with another six country such as Ireland, Japan, Malaysia, Singapore, Sweden, Switzerland, and United Kingdom. From this map we can also see the line on each node that representing connection between the country also the bigger the node shows that the country has made many contribution and publication in the paragenesis world and if we look closely on the Indonesia node, it showed that Indonesia had a close relation to some country which is United Kingdom, France, Sweden, and Iran but Indonesia represented by a small node which means that Indonesia still have a little publications in the paragenesis study (Figure 13).

D. CONCLUSIONS

This Article shows the overall summary about carbonate paragenesis obtained from secondary data which is a few articles published years before. Graphic from distribution of publications in paragenesis showed a stable and increasing trendline that indicates a consistent publications and study about this topic. From a few bibliometric analyses that were done resulted in a few bibliometric maps which is Co-occurrence bibliometric map of author keywords, Co-citation bibliometric map of cited author, and Co-citation bibliometric map from network of countries. Paragenesis study especially paragenetic sequence is very connected to many study fields. There are about 987 articles published with the topic of paragenesis since 1999 until 2022 with 537 authors from 46 different country. From the statistical analysis that were done according to the secondary data resulted a consistent trendline that is overall increasing even if there are drop in some year, but the graphic line keeps on increasing.

From this bibliographic analysis it can also showed the correlation between each key words in the co-occurrence map of keywords showing topics related to paragenesis especially paragenetic sequence. The map showed that paragenetic sequence connected to a few other key words such as Formation, Stage, fracture, mineralogy, deposit, phase, process, and many other key words. Another thing is that bibliometric analysis also can determine the mapping of authors co-citation analysis of each publication and from the country it originates from and divided into nine clusters with the biggest node which is United States, United Kingdom, China and Russia. Meanwhile Indonesia is in cluster two and mostly related with some country such as United Kingdom, France, Sweden, and Iran.

Based on the explanation above it shows that paragenesis is an important subject to continue study and research on. Paragenesis and diagenesis not only need to be studied in the ore deposits field, but it also needs to be used in the oil and gas fields, especially to determine when is the oil or hydrocarbon placing inside the pores and also from paragenesis we can know for sure about the geological history from the mineral deposits, formed fractures, compactions, and others products formed because of the diagenesis burials. Paragenesis also growing in many countries especially in Indonesia which is still small publications but have a promising growth. So, paragenesis study needs to be improved and much discussed because the future about paragenesis is still wide and promising.

E. ACKNOWLEDGEMENTS

The authors would like to thank you to the Faculty of Earth and Energy Technology, Trisakti University and to all the person involved in the making of this article.

GLOSARRY

- Paragenesis is a petrologic concept meaning an equilibrium sequence of mineral phases.
- **Carbonate** is a salt of carbonic acid, characterized by the presence of the carbonate ion, CO_2^{-3} .
- Diagenesis is changes to sediment or sedimentary rocks during and after rock formation, at temperatures and pressures less than that required for the formation of metamorphic rocks or melting. It does not include changes from wheatering.
- Bibliometric is a set of methods to quantitatively analyze scientific and technological literature.
- VOSviewer is a software tool for constructing and visualizing bibliometric networks. These networks may for instance include journals, researchers, or individual.
- **OpenRefine** is an open-source desktop application for data cleanup and transformation to other formats, an

activity commonly known as data wrangling. It is similar to spreadsheet applications, and can handle spreadsheet file formats such as CSV, but it behaves more like a database.

Hydrocarbon is an organic compound consisting entirely of hydrogen and carbon.

REFERENCES

[1] Birger Rasmussen, Janet R. Muhling; Evidence for widespread oil migration in the 1.88 Ga Gunflint Formation, Ontario, Canada. *Geology* 2019; 47 (10): 899–903. doi: https://doi.org/10.1130/G46469.1

[2] Ahr, Wayne M. 2008. Geology of carbonate reservoirs: the identification, description, and characterization of hydrocarbon reservoirs in carbonate rocks. Published by John Wiley & Sons, Inc., Hoboken, New Jersey. ISBN 978-0-470-16491-4

[3] Bruno Eustáquio Moreira Lima, Leonardo Ribeiro Tedeschi, André Luiz Silva Pestilho, Roberto Ventura Santos, Joselito Cabral Vazquez, Jarbas Vicente Poley Guzzo, Luiz Fernando De Ros. (2020). Deep-burial hydrothermal alteration of the Pre-Salt carbonate reservoirs from northern Campos Basin, offshore Brazil: Evidence from petrography, fluid inclusions, Sr, C and O isotopes. Marine and Petroleum Geology. Volume 113.

https://doi.org/10.1016/j.marpetgeo.2019.104143.

 [4] Craig, James R and Vaughan. (1994). Ore Microscopy and Ore Petrography. 2nd Edition.
 Published by John Wiley & Sons, Inc. USA. ISBN 0-471-55175-9

[5] Britannica, The Editors of Encyclopaedia."paragenesis". Encyclopedia Britannica, 20 Jul.1998.

[6] Dani, W.I. (2012). Studi Alterasi dan Mineralisasi Emas Berdasarkan Analisis Petrografi Conto Inti Pemboran Aaerah Arinem, kabupaten garut, Jawa Barat. [7] Widarsono, B. 2014. Porosity Versus Depth Characteristics of Some Reservoir Sandstones in Western Indonesia. Scientific Contribution Oil & Gas. 37(2):87-104

[8] Musu, J. 2007. To Explain The Nature of Core Porosity Using Results of Petrography Analysis. Lemigas Scientific Contributions. 30 (3): 43-51.

[9] Bruno Eustáquio Moreira Lima, Leonardo Ribeiro Tedeschi, André Luiz Silva Pestilho, Roberto Ventura Santos, Joselito Cabral Vazquez, Jarbas Vicente Poley Guzzo, Luiz Fernando De Ros. (2020). Deep-burial hydrothermal alteration of the Pre-Salt carbonate reservoirs from northern Campos Basin, offshore Brazil: Evidence from petrography, fluid inclusions, Sr, C and O isotopes. Marine and Petroleum Geology. Volume 113.

https://doi.org/10.1016/j.marpetgeo.2019.104143

[10] Dyah Hastuti, Endang Wiwik. (2017). The study of ore minerals parageneses in Ponorogo area, East Java. MATEC Web of Conferences. DOI: 101. 04018.

10.1051/matecconf/201710104018.

[11] Denyer, D.; Tranfield, D, 2009, Producing a Systematic Review. In The Sage Handbook of Organizational Research Methods; Buchanan, D.A., Bryman, A., Eds.; Sage Publications Ltd.: Thousand Oaks, CA, USA, 2009; pp. 671–689.

[12] Ellegaard, O., Wallin, J.A. The bibliometric analysis of scholarly production: How great is the impact?. *Scientometrics* **105**, 1809–1831 (2015). https://doi.org/10.1007/s11192-015-1645-z

[13] Hiatt, Eric. (2000). Links between Depositional and Diagenetic Processes in Basin Analysis: Porosity and Permeability Evolution in Sedimentary Rocks. In book: Fluids and Basin Evolution. Chapter: 4. Pp. 63-92. Publisher: Mineralogical Association of Canada.

[14] Melo, Gustavo & Monteiro, Lena & Xavier, Roberto & Moreto, Carolina & SANTIAGO, ERIKA & Dufrane, S.A. & Aires, Benevides & Santos, Antonio. (2017). Temporal

evolution of the giant Salobo IOCG deposit, Carajás Province (Brazil): constraints from paragenesis of hydrothermal alteration and U-Pb geochronology. Mineralium Deposita. 52. DOI: 10.1007/s00126-016-0693-5.

[15] Zupic, I., & Čater, T. (2015). Bibliometric Methods in Management and Organization. Organizational Research Methods, 18(3), 429– 472. https://doi.org/10.1177/1094428114562629

[16] Peng B, Li G, Li Z, et al. Discussion of multiple formation mechanisms of saddle dolomites—Comparison of geochemical data of Proterozoic–Paleozoic dolomites. Energy Exploration & Exploitation. 2018;36(1):66-96. doi: 10.1177/0144598717728464

[17] Hongwei Ping, Chunquan Li, Honghan Chen, Simon C. George, Se Gong; Overpressure release: Fluid inclusion evidence for a new mechanism for the formation of heavy oil. *Geology* 2020; 48 (8): 803–807. doi: https://doi.org/10.1130/G47227.1

[18] Blouet, Jean-Philippe; Imbert, Patrice; Foubert, Anneleen (2017). Mechanisms of biogenic gas migration revealed by seep carbonate paragenesis, Panoche Hills, California. AAPG Bulletin, 101(8), 1309– 1340. doi:10.1306/10171616021

[19] Rosales, I., Pomar, L., Al-Awwad, S.F., Microfacies, diagenesis and oil emplacement of the Upper Jurassic Arab-D carbonate reservoir in an oil field in central Saudi Arabia (Khurais Complex), Marine and Petroleum Geology (2018), doi: 10.1016/j.marpetgeo.2018.05.010

| Arti | cle Test | | | |
|--------|--|---|--|------------|
| ORIGIN | ALITY REPORT | | | |
| SIMIL | 2% 7% INTERNET SOURCES | 10% PUBLICATIONS | 6% STUDENT PA | PERS |
| PRIMAF | RY SOURCES | | | |
| 1 | Submitted to UPN Vete Student Paper | eran Yogyakart | а | 3% |
| 2 | Zornitsa Yordanova. "C of Firm's Innovation Pr Analysis", Springer Scie Media LLC, 2021 Publication | Chapter 12 Digi ocess - A Biblic ence and Busin | talization metric ess | 1 % |
| 3 | Idoia Rosales, Luis Pon Awwad. "Microfacies, c emplacement of the U carbonate reservoir in Saudi Arabia (Khurais C Petroleum Geology, 20 Publication | nar, Saad Fahd diagenesis and pper Jurassic A an oil field in c Complex)", Mar)18 | Al- oil rab-D entral rine and | 1 % |
| 4 | wikimili.com Internet Source | | | 1% |
| 5 | Bruno Eustáquio More Ribeiro Tedeschi, Andr Roberto Ventura Santo hydrothermal alteratio carbonate reservoirs fi | eira Lima, Leona é Luiz Silva Pes os et al. "Deep- on of the Pre-Sa rom northern (| ardo stilho, burial alt Campos | 1 % |

| | Basin, offshore Brazil: Evidence from petrography, fluid inclusions, Sr, C and O isotopes", Marine and Petroleum Geology, 2020 Publication | |
|----|---|-----|
| 6 | slideplayer.com Internet Source | 1% |
| 7 | Bo Peng, Guorong Li, Zongxing Li, Chenglin Liu, Yinhui Zuo, Wen Zhang, Li Yuan, Sha Zhao, Cheng Yao. "Discussion of multiple formation mechanisms of saddle dolomites— Comparison of geochemical data of Proterozoic–Paleozoic dolomites", Energy Exploration & Exploitation, 2017 Publication | <1% |
| 8 | api.crossref.org Internet Source | <1% |
| 9 | WWW.portergeo.com.au Internet Source | <1% |
| 10 | Jean-Philippe Blouet, Patrice Imbert, Anneleen Foubert. "Mechanisms of biogenic gas migration revealed by seep carbonate paragenesis, Panoche Hills, California", AAPG Bulletin, 2017 Publication | <1% |

11 Kevin Gerald Senduk, Moehammad Ali Jambak, Rizsa Rindira Sekar Ayu Heriadi,

<1%

Khoirul Arifin Syaifullah. "Biostratigraphic analysis and age determination, depositional environment and paleoclimate at well X section of the Kutai Basin, East Kalimantan region", AIP Publishing, 2021

Publication

| 12 | link.springer.com | <1 % |
|----|---------------------------------|------|
| 13 | repositorio.ual.es | <1 % |
| 14 | www.mdpi.com Internet Source | <1% |
| 15 | phys.org Internet Source | <1% |

| Exclude quotes | Off | Exclude matches | < 15 words |
|----------------------|-----|-----------------|------------|
| Exclude bibliography | On | | |