

Helminthologia_FULL Manuscript_HE-2023-0017 (1)

by Suriyani Tan

Submission date: 08-Aug-2023 11:21AM (UTC+0700)

Submission ID: 2142948578

File name: FULL_Manuscript_HE-2023-0017_2.docx (719.81K)

Word count: 9164

Character count: 54940

Fish and Food-Fatale: Food-borne Trematode *Opisthorchis viverrini* and Cholangiocarcinoma

Suriyani Tan^{1,2}, Machrumnizar Machrumnizar^{1,2*}

¹ Department of Parasitology, Faculty of Medicine, Universitas Trisakti, Jakarta 11440, Indonesia

² Tropical Diseases and Public Health Research Centre, Faculty of Medicine, Universitas Trisakti, Jakarta 11440, Indonesia

*Email: machrumnizar_md@trisakti.ac.id

Summary

Neglected Tropical Diseases (NTDs) are a group of communicable diseases with a long history with human beings. NTDs are the proxy of poverty since they affect those in low-income and extreme-poverty populations, as those populations lack access to proper health care, clean water, sanitary conditions, and hygiene. NTDs create losses for a nation that come from the health and the economic sectors as well since the costs of diagnosis, prevention, and treatment strain the national purse strings. One of the 20 different forms of NTDs on the list is food-borne trematodes, comprises of *Fasciola*, *Paragonimus*, *Clonorchis*, and *Opisthorchis*. Currently, it is estimated that food-borne trematodes can cause a devastating effect on mortality and morbidity. All of them are zoonotic, as humans become infected by ingestion of a second intermediate host, such as freshwater snails, fish, or water vegetables. *Opisthorchis viverrini*, one of the food-borne trematodes that can be found mostly in South East Asia regions, especially in the Mekong basin, is regarded as a group 1 carcinogen leading to cholangiocarcinoma (CCA). The aim of this study is to present the updated review of *Opisthorchis viverrini*. This study aims to present the updated review of *Opisthorchis viverrini* and CCA.

Keywords: Food-borne trematode, Neglected Tropical Diseases, *Opisthorchis viverrini*, Cholangiocarcinoma

Introduction

World Health Organization (WHO) stated that Neglected Tropical Diseases (NTDs) are a diverse group of infectious diseases, consisting of 20 diseases, mainly found in 149 tropical and subtropical countries (figure 1) (Mitra & Mawson, 2017). NTDs affect one-sixth of the global population, with the assumption that one in every six people will be affected by at least one type of NTDs. According to the WHO, the mortality and morbidity caused by NTDs are extremely high, as measured by disability-adjusted life years (DALYs). Globally, NTDs have resulted in about 534,000 fatalities and 57 million DALYs to date (Lin, Fang, Zheng, Wang, & Wu, 2022)

Neglected Tropical Diseases affect those who lack access to proper health care, clean water, sanitary conditions, and hygiene in low-income populations and even those who are extremely poor. NTDs create losses for a nation that come from both the health and the economic sectors as well, since the costs of diagnosis, prevention, and treatment place a strain on the national purse strings (Hotez, 2020; Redekop et al., 2017). Considerable population mobility contributes significantly to the transmission of NTDs thus more than a 70 percent of the cases are found in low-income or lower-middle-income countries. As a result of the increase in international traveling and climate change arising from global warming, NTDs are currently present in a number of European nations as well. Nonetheless, when the Covid-19 outbreak spread throughout the entire world, the global goals to fight NTDs has once again been disregarded (Adepoju, 2020; Tilli et al., 2021). NTDs lead to significant illness and mortality in addition to societal ignominy and prejudice; however, NTDs are effectively managed, diminished, or eradicated through the intervention of a variety of efficient public health measure. The current most cost-effective interventions in global health are those aimed at eradicating and controlling NTDs. By 2030, point 3.3 of the Sustainable

Development Goals (SDGs) on NTDs calls for the eradication of NTDs, malaria, tuberculosis (TB), and HIV/AIDS (Addisu et al., 2019).

Distribution of liver trematodes

One of the 20 different forms of NTDs on the list is food-borne trematodes. The term "food-borne trematodes" refers to a collection of illnesses brought on by a number of trematode worms, including *Clonorchis*, *Opisthorchis*, *Fasciola*, and *Paragonimus*. Two intermediary hosts serve as a broad overview of this category of trematodes. Aquatic vegetables, tiny and medium-sized fish, water snails, and crustaceans are examples of the first intermediate host and the second intermediate host, which are generally unique from one another. Humans as the definitive host will be infected if they consume food that has been contaminated by the second intermediate host (Robinson & Sotillo, 2022). The total number of people infected around the world may be more than the 2 million DALYs and deaths that the WHO estimates are lost globally. Food-borne trematodes is thought to have infected over 75 million individuals worldwide as of today, with 0.2 million new cases reported year. This information might not be accurate as there are many undetected cases (Chai & Jung, 2020).

The liver trematode worms in the *Opisthorchiidae* family that cause the food-borne trematode illnesses Opisthorchiasis and Clonorchiasis, which are *Opisthorchis viverrini* (*O. viverrini*), *Opisthorchis felinus* (*O. felinus*) and *Clonorchis sinensis* (*C. sinensis*). Over 700 million people globally, mostly in endemic regions of East and Southeast Asia, including Siberia, as well as numerous European countries, are at risk of acquiring one of these three tropical diseases (Suwannatrai, Saichua, & Haswell, 2018). *O. viverrini* is mainly restricted to Southeast Asia, *O. felinus* is indigenous to North Asia and Eastern Europe, and *C. sinensis* is prevalent in Korea, sections of Russia, China, and Vietnam (Maria Y Pakharukova et al., 2023; Sripa et al., 2021). According to China's national parasite inquiry, the prevalence of Clonorchiasis has increased by about 0.37 percent since the initial examination in 1988, and a recent study found that the prevalence of *C. sinensis* infection is 29.3 percent (Gao et al., 2020; Jiang et al., 2021). While nearly 1.2 million people reside with *O. felinus* in Western Siberia and other regions of the Russian Federation (Ovchinnikov, Mordvinov, & Fromm, 2017). In addition, it is thought that about 10 million people in Southeast Asia are affected with *O. viverrini* (Almanfaluthi, Widodo, Suttiapapa, Wongsaroj, & Sripa, 2022; Sripa & Echaubard, 2017). This liver parasite poses serious public health issues in Thailand, Democratic People's Republic of Lao, Cambodia, and Vietnam (Thinkhamrop, Khuntikeo, Sithithaworn, Thinkhamrop, Wangdi, Kelly Matthew, et al., 2019). There has been concern about these trematode worms in the western world due to reports of them in the United States (USA) as the surge in Asian immigrants or travelers, and the import of undercooked cyprinoid fish (Kaewkong et al., 2013; Lu et al., 2018). Additionally, some infections, like *Paragonimus westermani* (*P. westermani*) can be acquired locally in the US where infected first and second intermediate hosts may be prevalent (Fried & Abruzzi, 2010).

Life cycle and morphology of *O. viverrini*

The *O. viverrini* life cycle starts when the embryonic eggs laid by the adult worm enter the bile duct and are eliminated in the feces. These embryonated eggs do not hatch if they are not introduced to fresh water bodies (small ponds, rivers, flooded rice fields, reservoirs, or rivers), since an intermediate host snail *Bithynia*, such as the first intermediate host *O. viverrini*, will ingest them. The miracidia develop into sporocysts, rediae, and cercariae after hatching in the snail's digestive tract. Numerous thousands of cercariae are released every day as a result of snails' asexual reproduction. From the time eggs are consumed until cercaria are released, it takes roughly two

months. The free-swimming parasites then look for the cyprinoid fish, which serves as their next intermediate host. They encyst in the fins, skin, and muscles of the fish and develop into metacercaria after 21 days. The bile duct is where the metacercaria attach after swallowing and develop into adults, who lay eggs 3 to 4 weeks later (figure 2). The metacercaria first exist in the duodenum before ascending through the ampulla of Vater into the bile duct (Khieu et al., 2019; Sota et al., 2022; Suwannahitatorn, Webster, Riley, Mungthin, & Donnelly, 2019; Tangkawattana, Sereerak, Upontain, Tangkawattana, & Sripa, 2021).

Opisthorchis spp. eggs are frequently indistinguishable from *C. sinensis* eggs, measuring 19–30 μ m long by 10–20 μ m wide. The egg has been operated on, and the opercular "shoulder" and abopercular knob are both noticeable. As they are passed through excrement, embryonic eggs (figure 3). Adult worms are flattened on the abdomen, approximately 7 mm (5.4–10.2) long by 1.5 mm (0.8–1.9), lanceolate, thin, and transparent. The bile ducts and pancreas of the mammalian host are where adult worms live and connect to the mucosa. They may reside in the pancreatic duct, gallbladder, or larger bile duct in cases of serious illness. They produce 2000 to 4200 eggs per day (3160 eggs in average) (Charoensuk, Ribas, Chedtabud, & Prakobwong, 2022; P. Rachprakhon & W. Purivirojkul, 2021). The exact life span of *O. viverrini* in human is not clearly known, however, it may be over than 25 years (Kaewkes, 2003; Petney et al., 2018).

Genetic diversity of *O. viverrini* (*O. viverrini* sensu lato)

There had formerly been some investigations of the genetic diversity of *O. viverrini*, and the first publication utilizing the isoenzyme analysis method appeared in 1993 (Sueblinvong, Piyamputra, Priyanont, & Thaithong, 1993). Since then a number of molecular and genetic indicators studies have been carried out on *O. viverrini* using multilocus enzyme electrophoresis (MLEE), nuclear and mitochondrial DNA sequencing, and microsatellite marker fragment analysis (Kiatsopit et al., 2014; Namsanor et al., 2020; Opal Pitaksakulrat et al., 2018; W Saijuntha, 2007).

According to a thorough molecular study in 2007, *O. viverrini* sensu lato (s.l.) is a species complex consisting of numerous cryptic species (genetically different species that share a similar morphology) found in different wetland systems in Thailand and Laos (W Saijuntha, 2007). Bithynia snails serve as first intermediate vector, The first intermediate hosts in the life cycle are, a complex species that includes at least 11 cryptic species that are related to *O. viverrini* s.l. As a result, both species complexes are highly genetically variable in relation to geography (Hughes et al., 2017).

Despite this, a later study in 2010 that used AMOVA to assess the populations rejected *O. viverrini* s.l. as a species complex since a study didn't find any differences that were significant. Six populations of *O. viverrini* from Thailand, Laos, and Cambodia were examined in this genetic study using a single marker and a portion of the ND1 sequence. The limited variation that has been noted in a number of prior research also makes it crucial to emphasize that mitochondrial DNA appears to be inappropriate and untrustworthy for systematic and population genetic studies of *O. viverrini* (Thaenkham et al., 2010). Numerous research looking at genetic diversity and population structures have employed microsatellite DNA markers to bolster the idea that *O. viverrini* was a species complex (Namsanor et al., 2020; O. Pitaksakulrat et al., 2017). In Bangkok, Thailand, most of the *Opisthorchis viverrini* s.l. cercariae were detected in the snail vector, *Bithynia siamensis siamensis* (Phuphitchan Rachprakhon & Watchariya Purivirojkul, 2021). *O. viverrini* s.l. can be recognized from *C. sinensis* and *O. felinus* by differences in morphology, genetics, and geographic distribution (M. Y. Pakharukova & Mordvinov, 2022; Weerachai Saijuntha, Sithithaworn, Kaitsopit, Andrews, & Petney, 2014).

Significance of *O. viverrini* infection in public health

O. viverrini can cause hepatobiliary abnormalities, such as cholangitis, cholecystitis, cholelithiasis, advanced periductal fibrosis, and lastly, cholangiocarcinoma as the end point. Due to its connection to recurrent neglected tropical diseases whose effects on public health are advised to be underestimated, food-borne trematodes have come under increased scrutiny. The greatest incidence of CCA in the world and a hotspot for *O. viverrini* transmission can both be found in northeast Thailand. In this region, the annual incidence of CCA ranges from 93.8 to 317.6 per 100,000 people, with the majority of cases ending in death within one to two years of diagnosis (Steele et al., 2018). The widespread nature of helminth infection is a significant epidemiological trait. This means that while most infected people have relatively low worm burdens, some have substantial worm burdens. The risk of sickness is typically higher in those with the most worms, who also contribute disproportionately to environmental pollution. In *O. viverrini*, the usual pattern of overspread has a long history. It has been hypothesized that environmental exposure (host behavior shows considerable age dependence), individual vulnerability (genetic and immunogenic variables), and social factors are to blame for the overabundance of helminths (Echaubard, Sripa, Mallory, & Wilcox, 2016).

The *O. viverrini*'s complex life cycle, which involves a number of hosts and surroundings, makes it challenging to control using traditional techniques. Poor hygiene and sanitation may serve as the one of main factor resulting in stool contamination to water bodies, hence the reservoir hosts such as stray or domestic dogs and cats may also contribute to the transmission of this fluke. (Sripa, Tangkawattana, & Sangnikul, 2017; Tangkawattana & Sripa, 2018).

Risk factors of *O. viverrini* infections

The persistence of *O. viverrini* infection in endemic regions is influenced by a diverse array of risk factors (Suwannatrai et al., 2018). Age may be a factor in the increased burden of infection among working-age individuals, as well as the higher incidence of infection among men and those with lower levels of education. Cultural activities related to fishing, food preparation, and eating raw fish, which are deeply ingrained as parts of the region's native rice fish culture, are the major factors contributing to the high rate of *O. viverrini* infection in the endemic area (Prakobwong & Suwannatrai, 2020). Consuming freshwater fish that has already been salted, or smoked that is either raw or undercooked and has metacercaria on it can infect humans or other fish-eating animals. It is just the consumption of raw fish which caused the infection of *O. viverrini*, meanwhile the consumption of fish sauce doesn't have any impact as the fish sauce has undergone further process from the raw fish into fish sauce as the end product (Thaewngiew et al., 2014).

Food sharing among household and neighbor is a common practice in communities of local inhabitants (Isaan people) in Northeast Thailand, as this province bear the heaviest burden of *O. viverrini* infection (Phimpraphai, Tangkawattana, Kasemsuwan, & Sripa, 2018). Sharing raw fish dishes among neighbors may increase the prevalence of infection but also raise the danger of inflammation-related diseases because the average per-person infection intensity will be lower. Food sharing was tightly and favorably correlated with eating raw fish frequency and the dishes consumed variation (Saenna, Hurst, Echaubard, Wilcox Bruce, & Sripa, 2017). This implies that raw fish dishes are more likely to be accepted, offered, and consumed in households with higher food sharing propensities. Comparatively to homes that do not exchange food or do so only to a limited extent, this raises the risk of transmission. The smaller, less common raw fish is typically used in these communal dishes. The type of fish that is potentially infected with "*pla khao noi*" (Isan-Lao for small cyprinid fish) which is widely used in processing raw fish is usually found in local villages to be sold cheaply or distributed free of charge to fishermen's families and friends.

Other factors that promoting the everlasting transmission of *O. viverrine* including occupation factor where rice farmers and fishermen are more likely to consume raw fish dishes and defecate in or near wetlands and rice fields due to their daily activities, and repeated praziquantel (PZQ) treatment, as self- repeated used of PZQ without the medical supervision will lead to drug resistance (Suwannatrai et al., 2018; Thinkhamrop, Khuntikeo, Sithithaworn, Thinkhamrop, Wangdi, Kelly, et al., 2019).

The incidence of cholangiocarcinoma caused by liver trematodes

Fish-borne trematode infection and CCA have a well-established link. This infection can result in hepatobiliary diseases such as hepatomegaly, chronic cholangitis, cholecystitis, cholelithiasis, periportal stem fibrosis, gallstones, and bile duct cancer or cholangiocarcinoma (CCA) (Sripa et al., 2021). The World Health Organization's International Agency for Research on Cancer (WHO) categorized *C. sinensis* and *O. viverrini* as a Group 1A carcinogen (a metazoan parasite harmful to humans). It is estimated that 100 per 100,000 in men and 40 per 100,000 in women or approximately 1% of the individual with opisthorchiasis would develop into CCA and the disability-adjusted life years (DALYS) caused by the infection of *O. viverrine* and CCA as high as 150–235 × 103 DALYs (P. J. Brindley et al., 2021; Liau, Toh, & Shelat, 2023; W. Saijuntha, P. Sithithaworn, N. Kiatsovit, R. H. Andrews, & T. N. Petney, 2019).

Despite not being officially recognized as a human carcinogen, *O. felineus* is a member of Group III has potentially carcinogenic (Banks, Carbone, Blum, & Cesarman, 2012; Saltykova, Petrov, & Brindley, 2018; Tantrawatpan et al., 2020). In recent publication by Fedorova et al, they stated that a positive relationship between infection of *O. felineus* with the risk for CCA in the hospital-based study in Western Siberia, as the CCA patients had the history of *O. felineus* infection. This study revealed that this species might hold the possibility to be the Group 1A carcinogen, as the *O. viverrini* and *C. sinensis* (O. S. Fedorova et al., 2023).

In comparison to *O. felineus* area, long-term observation study indicated that CCA is more frequently documented in *O. viverrini* and *C. sinensis* endemic regions. Even though the carcinogenesis pathology caused by parasitic infection may be driven by a variety of events known as chronic inflammatory infectious processes, helminthiasis related to malignancy remain mainly unknown. Additional pathways involve genomic instability, induction of invasion, angiogenesis, and cancer metastases, deregulation of tumor suppressor genes, prolonged cell growth, cell death prevention, energy metabolism regulation, and immune evasion (Olga S Fedorova et al., 2020; Machicado & Marcos, 2016; Mariya Yurievna Pakharukova, da Costa, & Mordvinov, 2019). The three liver trematode species differ in how they become carcinogens, but there have been no comparative research studies, thus the information that is currently accessible is not sufficiently clear.

The role of parasite induces cholangiocarcinoma

Depending on its cellular source, hepatocellular carcinoma (HCC) and cholangiocarcinoma (CCA) are the two subtypes of primary malignant liver cancer that are distinguished histologically (El Jabbour, Lagana, & Lee, 2019). Hepatocellular carcinoma responsible for nearly 75 and 80 percent of cases worldwide. The rarest type of cancer is by far CCA constituting 10 to 25 percent of primary hepatic cancers. However, liver cancer incidence reveals 15 times higher in East Asian regions where Opisthorchiasis and/or Clonorchiasis are endemic. The fatality and morbidity of CCA are directly proportional and therefore have a shabby prognostic. The most prevalent kind of CCA is extrahepatic estimated 80–90% and about 50% of all CCA are perihilar (affecting the main

branches of the ducts). The 5-year survival rate for CCA patients after diagnosis has not increased is not good, as low as 5% to 17% of total cases (Chansitthichok et al., 2020). Cholangiocarcinoma can develop based on the anatomical location into intrahepatic (iCCA) or extrahepatic, while extrahepatic CCA are divided into perihilar (pCCA), and distal (dCCA); those may appear differently in terms of clinical findings, pathology, and epidemiology (figure 4) (Khan, Tavolari, & Brandi, 2019; Pratap, Raza, & Pratap, 2018; Singal, Lampertico, & Nahon, 2020). Various iCCA-specific risk factors are associated, although it is unclear how they cause iCCA to evolve. The prevalence of CCA varies globally due to variations in genetic and environmental risk factors. The liver fluke infestation is pathogenically linked to CCA throughout Asia, especially in regions where *O. viverrini*, *Clonorchis sinensis* and kindred flukes is endemic (Weerachai Saijuntha, Paiboon Sithithaworn, Nadda Kiatsopit, Ross H Andrews, & Trevor N Petney, 2019). A sizable previous amount of epidemiologic evidence supports the involvement of liver flukes in the emergence of CCA, particularly *Clonorchis sinensis* (Chinese liver fluke) and *Opisthorchis viverrini* (Thai liver fluke), both are significantly related to malignancy. Opisthorchiasis infection has been linked to an elevated risk of 2 to 5 times developing intra and extra-hepatic CCA as a result of adult flukes residing in the biliary tract over years and triggering an immunological response from the host and inducing prolonged inflammation (Massarweh & El-Serag, 2017).

Pathogenesis of *O. viverrini* induced cholangiocarcinoma

The mechanism of *O. viverrini* induced CCA is uncertain, but chronic inflammation is thought to play a role as results from the flat worm infecting the host's biliary tract. (Chaiyadet et al., 2015; Zimmermann, 2019). The most frequent cause of CCA is malignant transformation of cholangiocyte cells, while it is also possible for peribiliary gland epithelial cells and/or biliary stem cells to evolve into cancer. Moreover, hepatic stem/progenitor cells are confirmed to be the source of some subgroups of CCA and combined HCC/CCA (Nakagawa et al., 2018). There are at least three different pathways embroiled in the opisthorchiasis driven carcinogenesis. First, once the parasites mechanically injured the biliary epithelium, the tissue healing has begun. In order to build a cellular covering for the injury, tissue healing reactions cause the release of enzymes that breakdown the extracellular matrix, impede with epithelium adhesion, and stimulate cell proliferation. Normally, a sum of released substances reacts to cease the process after the injury has healed. Healing and recovery from injury is hampered by chronic epithelial breakage caused by the grazing and passage activity of the liver fluke. DNA damage and oncogenesis are presumed to be caused by the continuous cell division involved with tissue injury that is mechanically driven in the presence of external co-factors. The wound - healing reaction apparently causes cell proliferation, which, as in existence of co-factors, considerably aids in the growth of cancer (Sawanyawisuth, Sashida, & Sheng, 2021; Sripa, Tangkawattana, & Brindley, 2018).

Chronic immunological responses are the second pathway that can increase the risk of malignancy. During cholestasis and inflammation, it is discovered that the quantities of growth factors, cytokines, and bile acids are enhanced. These substances contribute to these molecular changes and promote the proliferation and viability of changed cells. The immune reaction is initiated when Nitric Oxide (NO) and oxygen radical spurt from activated phagocytes often eradicate various antigens. This reaction may have unfavorable harmful side effects as NO is carcinogenic and induces endogenous carcinogenic nitrosamine production (CCA primary catalyst) (Chaiyadet et al., 2015; Smout et al., 2011).

The immune reaction in those with opisthorchiasis infection unaffected the invading worms resulting in a protracted reaction and NO buildup. Inducible nitric oxide synthase (iNOS) is upregulated in inflammatory cholangiopathies and CCA, where cytokines promote iNOS production in epithelial

cells. Plasma and urine nitrite contents increase although nitrosamine and their precursors' contents from diet are not rising in opisthorchis infection. As the excretory metabolite of nitrosamines, nitrites are a good indicator of significant endogenous nitrosamine synthesis. The eradication of the parasite, which is easily achieved by using the anthelmintic medication praziquantel, should make this fluke infection outcome reversible (Jaiswal, LaRusso, & Gores, 2001; Smout et al., 2011; Sripa, Jumnainsong, Tangkawattana, & Haswell, 2018).

Further research established that the areas of fluke-induced tumorigenesis contain DNA breakage caused by nitrite generation ((Kawanishi, Hiraku, Pinlaor, & Ma, 2006). Oxidative DNA breakage and nitrates including 8-oxo-7,8-hydro-2'-deoxiguanine (8-oxodG) and 8-nitroguanine (8-NG), both have been connected to inflammation-related tumorigenesis. Nitric oxide (NO) and reactive nitrogen oxide species (RNOS) synthesis are known to interact with cytoplasmic DNA and proteins, both of which are the results of increased inducible nitric oxide synthase (iNOS) activity. Additionally, interaction between reactive nitrogen oxide species (RNOS) and cellular genome trigger in DNA strands breakage and transmutation. Nitric oxide (NO) and RNOS as well interact with DNA repair enzymes which is promptly rendered inactive by S-nitrosylation of the cysteine residue in its active site, to further enhance mutagenesis. During Opisthorchiasis elicited an inflammatory response and released ROS/RNS with high oxidative potential. Afterward, biomolecules in the inflammatory region, including *O. viverrini* in the bile duct, will react with ROS/RNS. Opisthorchiasis has also been linked to higher rates of nitrite and nitrate which are indicators of endogenous NO production. A study linking urinary nitrite levels and infection severity with a higher risk of CCA, demonstrated that nitrosamines are involved in the pathogenesis of opisthorchiasis. Nevertheless, the source of the rise in nitrite content is unknown, which may be related to modifications in the host's endogenous nitrite synthesis or a rising in external nitrites from fermentation products (Jaiswal, LaRusso, Burgart, & Gores, 2000; Vale et al., 2013) Long-lasting inflammation can also come from the eggs that the flukes lay and potential for cancer, even has not been thoroughly investigated, it may considerably impact toward disease as observed in the other primary cancer-causing fluke.

Finally, various soluble proteins and other mediators that fulfill numerous of function at the host-parasite boundary are excreted and secreted by liver fluke to endure for an extended of time in a hostile environment (Drurey & Maizels, 2021). It has long been assumed that this interaction affects host cellular homeostasis and aids in the development of cancer. In vitro studies demonstrated that the ES product of *O. viverrini* to mammalian cell lines causes morphological changes because the ES product is mitogenic and can be absorbed by host bile cells. This finding is supported by in vivo studies, the growth of CCA in a hamster opisthorchiasis model was faster when the model was given ES products before being infected with *O. viverrini* indicating that immunological memory can speed up carcinogenesis (Maria Y Pakharukova, Zaparina, Baginskaya, & Mordvinov, 2022).

Diagnosis and management of Opisthorchiasis

Diagnosis

O. viverrini infection diagnosis can be quite challenging because the majority of chronic human cases show few distinct signs or symptoms, with the exception of a palpable rise in liver frequency. Even in severe infections, hematologic and biochemical characteristics are typically unremarkable. Although ultrasound is not frequently used in the community, it can be useful in making the diagnosis since it can detect abnormalities in the gallbladder. The most accurate way to determine *O. viverrini* infection is to look for eggs in the bile, duodenal fluid, or feces/stool. Stool samples are examined for parasites, which is a common practice that provides a regular diagnosis. However,

sensitivity and accuracy depend on the technique used for the fecal inspection and the microscopist's skill. The community-level persistence of *O. viverrini* infection is also a result of this constraint (Charoensuk, Subrungruang, Mungthin, Pinlaor, & Suwannahitatorn, 2019).

Management

Praziquantel (PZQ) is an oral anthelmintic medication used to treat Opisthorchiasis. The recommended dosage for adults is 75 mg/kg taken orally for two days, split three times daily. The same dosage applies to children (CDC – National Center for Health Statistics – Homepage. <https://www.cdc.gov/nchs/>. Accessed on June 22, 2023). Food safety is a high priority for disease control due to traditional habits in the impacted regions. Prevention strategies for the disease include programs to raise awareness of food safety, a better approach for identifying and treating opisthorchiasis, and school-based health program for longer-term goals.

To overcome the continuous transmission of *O. viverrini*, a novel control method known as the "Lawa model" employing the EcoHealth / One Health approach was implemented to the Khon Kaen Province's in Lake Lawa area, Thailand where liver flukes are endemic. The program comprises anthelmintic medicine therapy, innovative approaches to intense health education in the community and in schools, ecosystem monitoring, and engaged community involvement. Due to its success stories, this technique began in 2007 and has continued for more than ten years, as this control method has been adopted by WHO

Diagnosis and management of Cholangiocarcinoma

Diagnosis

The diagnosis of CCA is frequently complicated and necessitates a combination of several diagnostic techniques to pinpoint the stricture's anatomical location, differentiate within malignancy and benign strictures, define CCA coming from mixed hepatocellular-CCA, the bile or other primary hepatic tumors. As addition, the diagnosis of CCA must take into consideration of the tumor's stage and grade in order to determine the best course of therapy (Paul J. Brindley et al., 2021; Shin, Moon, & Kim, 2023). Until nowadays, CCA is confirmed by the patient history taking, clinical and laboratory (tumor biomarkers) also imaging diagnosis. As a result, numerous efforts have been undertaken to develop novel CCA biomarkers that could help with diagnosis as well as forecast prognosis and therapeutic response.

There are several biomarkers that can be used for the diagnosis of cholangiocarcinoma (CCA) depends on the type of samples. Talking about the samples from serum, carbohydrate antigen 19-9 (CA19-9) and carcinoembryonic antigen (CEA) are the most useful biomarkers in the since both of these biomarkers have good sensitivity and specificity in clinical usage (Tshering, Dorji, Chaijaroenkul, & Na-Bangchang, 2018). Biomarkers used from the tissue samples include gene methylations and DNA-related markers. It is important to highlight, however, that no one biomarker can be used alone to accurately diagnose CCA. Combining these possible biomarkers with other laboratory procedures is likely to considerably improve diagnostic accuracy.

Various investigations indicated possible non-invasive biomarkers for cholangiocarcinoma (CCA) found in body fluid, such as blood (serum), bile, urine and or other body fluids which are used until current such as:

1. Carbohydrate antigen (CA): CA19-9 is an antigen found in normal epithelial cells, including biliary ducts. CA 19-9 (carbohydrate antigen 19-9) is a cell surface glycoprotein complex that has been linked to pancreatic ductal adenocarcinoma (PDAC), and routinely used as tumor-marker in CCA. Elevated levels of CA19-9 correlate with tumor mass and degree of tumor differentiation. (Lee, Teng, & Shelat, 2020; Liang et al., 2015). Several

376 studies found out that combination of CA19-9 with other biomarkers increasing the
377 sensitivity and specificity. Study by Lapitz A, et al revealed that combination of serum
378 extravesicles (EVs) contain protein biomarkers with CA19-9 has remarkable outcomes, as
379 the sensitivity and specificity is higher compared than only CA 19-9 alone (Lapitz et al.,
380 2023). However, CA19-9 is not specific to CCA and tumor of hepatobiliary, but can also
381 be elevated in another tumor outside the gastrointestinal tract is one of the limitations.
382 Another condition such as inflammatory condition or obstruction of biliary and respiratory
383 tracts may also result in the elevation of CA19-9 level (Lee et al., 2020). Another limitation
384 of CA19-9 as this biomarker has low positive predictive value, thus not useful for screening
385 purpose.

386 2. Carcinoembryonic antigen (CEA): CEA is oncofetal glycoprotein produced by the
387 gastrointestinal system (colon) that has been linked to tumor metastasis. This biomarker
388 has high sensitivity and specificity, like CA-19-9. Ample researches have been published
389 that highlight the diagnostic value of elevated CEA levels in bodily fluids in primary
390 malignant tumors of the stomach, colon, biliary system, and pancreas (Joo et al., 2003;
391 Loosen et al., 2017). However, like CA19-9, CEA is not specific to CCA and can be
392 elevated in other types of tumors, and CEA has low sensitivity and specificity as prognostic
393 marker (Kim et al., 2021).

394 3. Micro-RNA (miRNA): miRNA is small non-coding RNA that regulate the gene expression,
395 as nowadays have been explored extensively as biomarkers of tumor, as miRNAs can
396 circulate throughout the whole body, making these molecules obtainable in wide range of
397 body fluid. miRNAs can play dual role, either as protective or oncogenic factors to tumor.
398 The level of miRNAs varies, either increasing or decreasing, according to the tumor types.
399 The level of several miRNAs, such as miR-26a, miR-141, miR-210, miR-31, miR-21, and
400 miR-421 are increasing in CCA, through cell proliferation modulation (Haga, Yan,
401 Takahashi, Wood, & Patel, 2014). The accuracy of miRNAs as potential biomarker with
402 another biomarker has also been studied. miRNAs-150, expressed in immune cells, such
403 as B and T cells, if combined with CA19-9 results in more accurate CCA diagnosis. Level
404 of CCA-19 inversely proportional with miRNA-150, as CCA 19-9 level is high in CCA,
405 whereas miRNA-150 level is low (Salem, Ghazala, El Gendi, Emara, & Ahmed, 2020).

406 4. Long non-coding RNA (lncRNA): lncRNA is type of non-coding RNA with more than 200
407 nucleotides, which can be found inside in the EVs that circulate through human body. Like
408 miRNA, lncRNA also play dual role, as tumor suppressor role or oncogenic role. Some
409 specific lncRNAs that have been studied in relation to Cholangiocarcinoma, such as
410 Prostate cancer-associated transcript 1(PCAT1), Neighboring enhancer of FOXA2 (NEF),
411 Metastasis-associated lung adenocarcinoma transcript 1(MALAT1), H19, Colon cancer-
412 associated transcript 1 (CCAT1), and ZEB1 Antisense RNA 1(ZEB-AS1) (Wu, Hayat, Hu,
413 & Yang, 2022; Yang et al., 2020).

414 5. Exostosin1 (EXT1): EXT1 is a gene encodes an endoplasmic reticulum-resident type II
415 transmembrane glycosyltransferase involved in the chain elongation step of heparan sulfate
416 biosynthesis. This gene is believed as tumor suppressor gene and has been investigated as
417 a potential biomarker for CCA. However, its diagnostic accuracy and clinical utility are still
418 being evaluated (Zeng & Tao, 2015).

419 It is important to note that while these biomarkers have been studied for their potential in
420 diagnosing CCA, none of them have been considered adequate for clinical application on their
421 own. To improve the accuracy of CCA diagnosis, the combination of biomarkers and clinical
422 finding should be considered. In the immediate term, the key problem of effective CCA biomarkers

is not to identify new molecules and pathways, but to validate or disprove the significance of present approaches in bridging the gap between research findings and clinical application. This implies that the emphasis should be on confirming current biomarkers and study methodologies in order to facilitate research translation into clinical practice.

Recent breakthroughs in "Omics" technologies provide great prospects for the identification of biomarkers connected to diseases. Next-generation biomarkers for cholangiocarcinoma propose several potential advantages, as these biomarkers provide early and non-invasive diagnostic that offer reassurance for the patient, as the CCA can be detected at the early phase and prompt treatment could be applied as soon as possible, leading to decreasing mortality rate; improving patients outcomes due to more accurate and reliable diagnostic information; better prognostic and information. These biomarkers also reliable for non-invasive monitoring of disease progression, and in return, reducing the needs for the invasive procedures, such as tumor biopsies, and allows for more frequent and convenient monitoring of patients (Chang, Chen, Yeh, & Hsiao, 2020; Rodrigues et al., 2021; Silsirivanit, Sawanyawisuth, Riggins, & Wongkham, 2014).

Management

Management of CCA differs according to the anatomical location of the disease, similar to how it is diagnosed. iCCA frequently receives therapy with surgical resection and estimated 60 percent of patients continue to live for five years after the procedure. But the majority of iCCA patients have big, incurable tumors when they first show. For patients in the beginning phases of pCCA or dCCA, surgery is also a choice. Surgery combined with neoadjuvant chemoradiation is the sole curative treatment for pCCA attended by liver transplantation. The overall five-year survival rate of pCCA patients who receive surgery is approximately 10% - 40%. While pCCA patients who received neoadjuvant treatment attended by liver transplantation survived for 5 years around 65 percent. The management of dCCA is also surgical resection, yet the Whipple operation technique is often used. After surgical, a mere 27 percent of patients remain alive for five years (Bao et al., 2022; Paul J. Brindley et al., 2021). Additionally, a preventative program for undiagnosed populations with initial-stage CCA is necessary to undergo surgery. The only effective treatment for CCA is a resection surgery (Alsaleh et al., 2019; Mariya Yurievna Pakharukova et al., 2019; Rizvi & Gores, 2013).

ACKNOWLEDGMENT

None

ETHICAL APPROVAL AND CONFLICT OF INTEREST

This review article is not accompanied by an ethical statement because it does not involve humans or animals as research subjects. All of the authors have read and agree with the manuscript. The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

AUTHORS CONTRIBUTION

ST and MM both contributed equally in all of the process, from conceptualization, drafting, writing, review and editing.

469 REFERENCES

- 470 ADDISU, A., ADRIAENSEN, W., BALEW, A., ASFAW, M., DIRO, E., DJIRMAY, A. G., . . . HARRIES, A. D.
471 (2019): Neglected tropical diseases and the sustainable development goals: an urgent call for action from the front line.
472 *BMJ Glob Health*, 4(1): e001334. DOI: 10.1136/bmjgh-2018-001334
- 473 ADEPOJU, P. (2020): NTDs in the time of COVID-19. *Lancet Microbe*, 1(6): e244. DOI:10.1016/S2666-
474 5247(20)30153-1
- 475 ALMANFALUTHI, M., WIDODO, S., SUTTIPIRAPA, S., WONGSAROJ, T., & SRIPA, B. (2022): The burden of
476 opisthorchiasis and leptospirosis in Thailand: A nationwide syndemic analysis. *Acta Tropica*, 226: 106227.
477 DOI:10.1016/j.actatropica.2021.106227
- 478 ALSALEH, M., LEFTLEY, Z., BARBERA, T. A., SITHITHAWORN, P., KHUNTIKEO, N., LOILOME, W., . . .
479 TAYLOR-ROBINSON, S. D. (2019): Cholangiocarcinoma: a guide for the nonspecialist. *Int J Gen Med*, 12: 13-23.
480 DOI:10.2147/ijgm.s186854
- 481 BANKS, L., CARBONE, A., BLUM, H., & CESARMAN, E. (2012): Biological agents: A review of human
482 carcinogens. *IARC Monogr*, 100: 49.
- 483 BAO, F., LIU, J., CHEN, H., MIAO, L., XU, Z., & ZHANG, G. (2022): Diagnosis Biomarkers of Cholangiocarcinoma
484 in Human Bile: An Evidence-Based Study. *Cancers*, 14(16): 3921. DOI:10.3390/cancers14163921
- 485 BRINDLEY, P. J., BACHINI, M., ILYAS, S. I., KHAN, S. A., LOUKAS, A., SIRICA, A. E., The, B.T., GORES, G.
486 J. (2021): Cholangiocarcinoma. *Nat Rev Dis Primers*, 7(1): 65. DOI:10.1038/s41572-021-00300-2
- 487 CHAI, J.-Y., & JUNG, B.-K. (2020): Foodborne intestinal flukes: A brief review of epidemiology and geographical
488 distribution. *Acta Tropica*, 201: 105210. DOI:10.1016/j.actatropica.2019.105210
- 489 CHAIYADET, S., SOTILLO, J., SMOUT, M., CANTACESSI, C., JONES, M. K., JOHNSON, M. S., . . .
490 LAOHAVIROJ, M. (2015): Carcinogenic liver fluke secretes extracellular vesicles that promote cholangiocytes to
491 adopt a tumorigenic phenotype. *The Journal of Infectious Diseases*, 212(10): 1636-1645. DOI:10.1093/infdis/jiv291
- 492 CHANG, Y.-C., CHEN, M.-H., YEH, C.-N., & HSIAO, M. (2020): Omics-Based Platforms: Current Status and
493 Potential Use for Cholangiocarcinoma. *Biomolecules*, 10(10): 1377. DOI:10.3390/biom10101377
- 494 CHANSITTHICHOK, S., CHAMNAN, P., SARKHAMPEE, P., LERTSAWATVICHAN, N., VORAVISUTTHIKUL,
495 P., & WATTANARATH, P. (2020): Survival of Patients with Cholangiocarcinoma Receiving Surgical Treatment in
496 an O. viverrini Endemic Area in Thailand: A Retrospective Cohort Study. *Asian Pac J Cancer Prev*, 21(4): 903-909.
497 DOI:10.31557/apjcp.2020.21.4.903
- 498 CHAROENSUK, L., RIBAS, A., CHEDTABUD, K., & PRAKOBWONG, S. (2022): Infection rate of Opisthorchis
499 viverrini metacercariae in cyprinoid fish from the markets and its association to human opisthorchiasis in the local
500 community in the Northeast Thailand. *Acta Tropica*, 225: 106216. DOI:10.1016/j.actatropica.2021.106216
- 501 CHAROENSUK, L., SUBRUNGRUANG, I., MUNGTHIN, M., PINLAOR, S., & SUWANNAHITATORN, P.
502 (2019): Comparison of stool examination techniques to detect Opisthorchis viverrini in low intensity infection. *Acta*
503 *Tropica*, 191: 13-16. DOI:10.1016/j.actatropica.2018.12.018
- 504 DRUREY, C., & MAIZELS, R. M. (2021): Helminth extracellular vesicles: Interactions with the host immune system.
505 *Molecular Immunology*, 137: 124-133. DOI:10.1016/j.molimm.2021.06.017
- 506 ECHAUBARD, P., SRIPA, B., MALLORY, F. F., & WILCOX, B. A. (2016): The role of evolutionary biology in
507 research and control of liver flukes in Southeast Asia. *Infection, Genetics and Evolution*, 43: 381-397.
508 DOI:10.1016/j.meegid.2016.05.019
- 509 EL JABBOUR, T., LAGANA, S. M., & LEE, H. (2019): Update on hepatocellular carcinoma: Pathologists' review.
510 *World J Gastroenterol*, 25(14): 1653-1665. DOI:10.3748/wjg.v25.i14.1653
- 511 FEDOROVA, O.S., FEDOTOVA, M.M., ZVONAREVA, O.I., MAZEINA, S.V., KOVSHIRINA, Y.V.,
512 SOKOLOVA, T.S., GOLOVACH, E.A., KOVSHIRINA, A.E., KONOVALOVA, U.V., KOLOMEETS, I.L.,
513 GUTOR, S.S., PETROV, V.A., HATTENDORF, J., OGORODOVA, L.M., ODERMATT, P. (2020): Opisthorchis
514 felinus infection, risks, and morbidity in rural Western Siberia, Russian Federation. *PLoS Neglected Tropical*
515 *Diseases*, 14(6): e0008421. DOI: 10.1371/journal.pntd.0008421
- 516 FEDOROVA, O.S., KOVSHIRINA, A.E., KOVSHIRINA, Y.V., HATTENDORF, J., ONISHCHENKO, S.V.,
517 KATANAKHOVA, L.L., TASLICKI, S.S., CHIZHIKOV, A.V., TATAUROV, I.A., VTORUSHIN, S.V., SRIPA, B.,
518 OGORODOVA, L.M., ODERMATT, P. (2023): Opisthorchis Felinus Infection is a Risk Factor for
519 Cholangiocarcinoma in Western Siberia: A Hospital-based Case-control Study. *Clin Infect Dis*, 76(3): e1392-e1398.
520 DOI:10.1093/cid/ciac497
- 521 FRIED, B., & ABRUZZI, A. (2010): Food-borne trematode infections of humans in the United States of America.
522 *Parasitol Res*, 106(6): 1263-1280. DOI:10.1007/s00436-010-1807-0

- GAO, Y., LI, Y., LIU, X., ZHANG, T., YU, G., WANG, Y., SHI, Y., CHI, X., WANG, X., GAO, X., WU, R., ZHANG, Y., HANG, L., SUN, S., GUAN, Y., XU, Y., MENG, J., LIU, X., JIANG, C., MA, H., LUO, L., YANG, Q., YIN, X., PENG, F., ZHI, Y., QU, W., ZHANG, X., REN, T., LIU, L., ZHAO, J., ZHANG, F., YAKEPU, A., PAN, Y., XU, H., NIU, J. (2020): High prevalence of *Clonorchis sinensis* infections and coinfection with hepatitis virus in riverside villages in northeast China. *Sci Rep*, 10(1): 11749. DOI:10.1038/s41598-020-68684-x
- HAGA, H., YAN, I., TAKAHASHI, K., WOOD, J., & PATEL, T. (2014): Emerging insights into the role of microRNAs in the pathogenesis of cholangiocarcinoma. *Gene Expr*, 16(2): 93-99. DOI:10.3727/105221614X13919976902174
- HOTEZ, P. J. (2020): NTDs in the 2020s: An epic struggle of effective control tools versus the Anthropocene. *PLOS Neglected Tropical Diseases*, 14(9): e0007872. DOI:10.1371/journal.pntd.0007872
- HUGHES, T., O'CONNOR, T., TECHASEN, A., NAMWAT, N., LOILOME, W., ANDREWS, R. H., . . . TAYLOR-ROBINSON, S. D. (2017): Opisthorchiasis and cholangiocarcinoma in Southeast Asia: an unresolved problem. *International Journal of General Medicine*, 10: 227-237. DOI:10.2147/IJGM.S133292
- JAISWAL, M., LARUSSO, N. F., BURGART, L. J., & GORES, G. J. (2000): Inflammatory cytokines induce DNA damage and inhibit DNA repair in cholangiocarcinoma cells by a nitric oxide-dependent mechanism. *Cancer Res*, 60(1): 184-190. DOI:10.1053/gast.2001.20875
- JAISWAL, M., LARUSSO, N. F., & GORES, G. J. (2001): Nitric oxide in gastrointestinal epithelial cell carcinogenesis: linking inflammation to oncogenesis. *Am J Physiol Gastrointest Liver Physiol*, 281(3): G626-G634. DOI:10.1152/ajpgi.2001.281.3.G626
- JIANG, Z.-H., WAN, X.-L., LV, G.-L., ZHANG, W.-W., LIN, Y., TANG, W.-Q., . . . CHAN, C. H. F. (2021): High prevalence of *Clonorchis sinensis* infection in Guangxi, Southern China. *Trop Med Health*, 49(1): 6. DOI:10.1186/s41182-021-00297-0
- JOO, K. R., KIM, D. H., PARK, J. H., BANG, S.-J., SHIN, J. W., PARK, N. H., & PARK, J. H. (2003): The role of bile carcinoembryonic antigen in diagnosing bile duct cancer. *J Korean Med Sci*, 18(6): 855-858. DOI:10.3346/jkms.2003.18.6.855
- KAEWKES, S. (2003): Taxonomy and biology of liver flukes. *Acta Tropica*, 88(3): 177-186. DOI:10.1016/j.actatropica.2003.05.001
- KAEWKONG, W., INTAPAN, P.M., SANPOOL, O., JANWAN, P., THANCHOMNANG, T., LAUMMAUNWAI, P., LULITANOND, V., DOANH, P.N., MALEEWONG, W. (2013): Molecular differentiation of *Opisthorchis viverrini* and *Clonorchis sinensis* eggs by multiplex real-time PCR with high resolution melting analysis. *Korean J Parasitol*, 51(6): 689-694. doi:10.3347/kjp.2013.51.6.689. DOI:10.3347/kjp.2013.51.6.689
- KAWANISHI S, HIRAKU Y, PINLAOR S, MA N. (2006): Oxidative and nitrative DNA damage in animals and patients with inflammatory diseases in relation to inflammation-related carcinogenesis. *Biol Chem*, 387(4):365-72. DOI: 10.1515/BC.2006.049.
- KHAN, S. A., TAVOLARI, S., & BRANDI, G. (2019): Cholangiocarcinoma: Epidemiology and risk factors. *Liver International*, 39(S1): 19-31. DOI:10.1111/liv.14095
- KHIEU, V., FÜRST, T., MIYAMOTO, K., YONG, T.-S., CHAI, J.-Y., HUY, R., SINUON, M., ODERMATT, P. (2019): Is *Opisthorchis viverrini* emerging in Cambodia?. *Adv Parasitol*, 103: 31-73. DOI:10.1016/bs.apar.2019.02.002
- KIATSOPIT, N., SITHITHAWORN, P., SAIJUNTHA, W., PITAKSAKULRAT, O., PETNEY, T. N., WEBSTER, J. P., & ANDREWS, R. H. (2014): Analysis of the population genetics of *Opisthorchis viverrini* sensu lato in the Nam Ngum River wetland, Lao PDR, by multilocus enzyme electrophoresis. *Parasitol Res*, 113(8): 2973-2981. DOI:10.1007/s00436-014-3959-9
- KIM, H. S., HAN, Y., KANG, J. S., KANG, Y. H., LEE, M., SOHN, H. J., KIM, H., KWON, W., JANG, J. Y. (2021): Serum carcinoembryonic antigen and carbohydrate antigen 19-9 as preoperative diagnostic biomarkers of extrahepatic bile duct cancer. *BJS Open*, 5(6). DOI:10.1093/bjsopen/zrab127
- LAPITZ, A., AZKARGORTA, M., MILKIEWICZ, P., OLAIZOLA, P., ZHURAVLEVA, E., GRIMSRUD, M. M., SCHRAMM, C., ARBELAIZ, A., O'ROURKE, C.J., LA CASTA, A., MILKIEWICZ, M., PASTOR, T., VESTERHUS, M., JIMENEZ-AGÜERO, R., DILL, M.T., LAMARCA, A., VALLE, J.W., MACIAS, RIR., IZQUIERDO-SANCHEZ, L., PÉREZ CASTAÑO, Y., CABALLERO-CAMINO, F.J., RIAÑO, I., KRAWCZYK, M., IBARRA, C., BUSTAMANTE, J., NOVA-CAMACHO, L.M., FALCON-PEREZ, J.M., ELORTZA, F., PERUGORRIA, M.J., ANDERSEN, J.B., BUJANDA, L., KARLSEN, T.H., FOLSERAAAS, T., RODRIGUES, P.M., BANALES, J. M. (2023): Liquid biopsy-based protein biomarkers for risk prediction, early diagnosis, and prognostication of cholangiocarcinoma. *J Hepatol*, 79(1): 93-108. DOI:10.1016/j.jhep.2023.02.027
- LEE, T., TENG, T. Z. J., & SHELAT, V. G. (2020): Carbohydrate antigen 19-9 - tumor marker: Past, present, and future. *World J Gastrointest Surg*, 12(12): 468-490. DOI:10.4240/wjgs.v12.i12.468

LIANG, B., ZHONG, L., HE, Q., WANG, S., PAN, Z., WANG, T., & ZHAO, Y. (2015): Diagnostic Accuracy of Serum CA19-9 in Patients with Cholangiocarcinoma: A Systematic Review and Meta-Analysis. *Med Sci Monit*, 21: 3555-3563. DOI:10.12659/msm.895040

LIAU, M. Y. Q., TOH, E. Q., & SHELAT, V. G. (2023): *Opisthorchis viverrini*-Current Understanding of the Neglected Hepatobiliary Parasite. *Pathogens*, 12(6). DOI:10.3390/pathogens12060795

LIN, Y., FANG, K., ZHENG, Y., WANG, H.-L., & WU, J. (2022): Global burden and trends of neglected tropical diseases from 1990 to 2019. *Journal of Travel Medicine*, 29(3): taac031. DOI:10.1093/jtm/taac031

LOOSEN, S. H., RODERBURG, C., KAUERTZ, K. L., KOCH, A., VUCUR, M., SCHNEIDER, A. T., BINNEBÖSEL, M., ULMER, T.F., LURJE, G., SCHOENING, W., TACKE, F., TRAUTWEIN, C., LONGERICH, T., DEJONG, C.H., NEUMANN, U.P., LUEDDE, T. (2017): CEA but not CA19-9 is an independent prognostic factor in patients undergoing resection of cholangiocarcinoma. *Sci Rep*, 7(1): 16975. DOI:10.1038/s41598-017-17175-7

LU, X.-T., GU, Q.-Y., LIMPANONT, Y., SONG, L.-G., WU, Z.-D., OKANURAK, K., & LV, Z.-Y. (2018): Snail-borne parasitic diseases: an update on global epidemiological distribution, transmission interruption and control methods. *Infectious Diseases of Poverty*, 7(1): 28. DOI:10.1186/s40249-018-0414-7

MACHICADO, C., & MARCOS, L. A. (2016): Carcinogenesis associated with parasites other than *Schistosoma*, *Opisthorchis* and *Clonorchis*: A systematic review. *Int J Cancer*, 138(12): 2915-2921. DOI:10.1002/ijc.30028

MASSARWEH, N. N., & EL-SERAG, H. B. (2017): Epidemiology of hepatocellular carcinoma and intrahepatic cholangiocarcinoma. *Cancer control*, 24(3): 1073274817729245. DOI:10.1177/1073274817729245

MITRA, A. K., & MAWSON, A. R. (2017): Neglected Tropical Diseases: Epidemiology and Global Burden. *Trop. Med. Infect. Dis*, 2(3). DOI:10.3390/tropicalmed2030036

NAKAGAWA, H., HAYATA, Y., YAMADA, T., KAWAMURA, S., SUZUKI, N., & KOIKE, K. (2018): Peribiliary Glands as the Cellular Origin of Biliary Tract Cancer. *Int. J. Mol. Sci*, 19(6): 1745. DOI:10.3390/ijms19061745

NAMSANOR, J., PITAKSAKULRAT, O., KOPOLRAT, K., KIATSOPIT, N., WEBSTER, B. L., GOWER, C.M., WEBSTER, J.P., LAHA, T., SAIJUNTHA, W., LAOPROM, N., ANDREWS, R.H., PETNEY, T.N., BLAIR, D., SITHITHAWORN, P. (2020): Impact of geography and time on genetic clusters of *Opisthorchis viverrini* identified by microsatellite and mitochondrial DNA analysis. *Int J Parasitol*, 50(14): 1133-1144. DOI:10.1016/j.ijpara.2020.06.011

OVCHINNIKOV, V. Y., MORDVINOV, V. A., & FROMM, B. (2017): Extreme conservation of miRNA complements in opisthorchiids. *Parasitol Int*, 66(6): 773-776. DOI:10.1016/j.parint.2017.09.006

PAKHARUKOVA, M. Y., DA COSTA, J. M. C., & MORDVINOV, V. A. (2019): The liver fluke *Opisthorchis felinus* as a group III or group I carcinogen. *Aopen*, 2: 23. DOI:10.1051/fopen/2019016

PAKHARUKOVA, M. Y., LISHAI, E. A., ZAPARINA, O., BAGINSKAYA, N. V., HONG, S.-J., SRIPA, B., & MORDVINOV, V. A. (2023): *Opisthorchis viverrini*, *Clonorchis sinensis* and *Opisthorchis felinus* liver flukes affect mammalian host microbiome in a species-specific manner. *PLoS Neglected Tropical Diseases*, 17(2): e0011111. DOI:10.1371/journal.pntd.0011111

PAKHARUKOVA, M. Y., & MORDVINOV, V. A. (2022): Similarities and differences among the *Opisthorchiidae* liver flukes: insights from *Opisthorchis felinus*. *Parasitology*, 149(10): 1306-1318. DOI:10.1017/s0031182022000397

PAKHARUKOVA, M. Y., ZAPARINA, O., BAGINSKAYA, N. V., & MORDVINOV, V. A. (2022): Global changes in gene expression related to *Opisthorchis felinus* liver fluke infection reveal temporal heterogeneity of a mammalian host response. *Food and Waterborne Parasitology*, 27: e00159. DOI:10.1016/j.fawpar.2022.e00159

PETNEY, T. N., ANDREWS, R. H., SAIJUNTHA, W., TESANA, S., PRASOPDEE, S., KIATSOPIT, N., & SITHITHAWORN, P. (2018): Chapter One - Taxonomy, Ecology and Population Genetics of *Opisthorchis viverrini* and Its Intermediate Hosts. In B. Sripa & P. J. Brindley (Eds.) *Advances in Parasitology*. Volume 101. Massachusetts, USA: Academic Press, pp. 1-39. DOI:10.1016/bs.apar.2018.05.001

PHIMPRAPHAI, W., TANGKAWATTANA, S., KASEMSUWAN, S., & SRIPA, B. (2018): Chapter Four - Social Influence in Liver Fluke Transmission: Application of Social Network Analysis of Food Sharing in Thai Isaan Culture. In B. Sripa & P. J. Brindley (Eds.) *Advances in Parasitology*. Volume 101. Massachusetts, USA: Academic Press, pp. 97-124. DOI:10.1016/bs.apar.2018.05.004

PITAKSAKULRAT, O., KIATSOPIT, N., LAOPROM, N., WEBSTER, B. L., WEBSTER, J. P., LAMBERTON, P. H. L., LAHA, T., ANDREWS, R.H., PETNEY, T.N., BLAIR, D., CARLTON, E.J., SPEAR, R.C., SITHITHAWORN, P. (2017): Preliminary genetic evidence of two different populations of *Opisthorchis viverrini* in Lao PDR. *Parasitol Res*, 116(4): 1247-1256. DOI:10.1007/s00436-017-5401-6

PITAKSAKULRAT, O., WEBSTER, B. L., WEBSTER, J. P., LAHA, T., SAIJUNTHA, W., LAMBERTON, P. H., KIATSOPIT, N., ANDREWS, R.H., PETNEY, T.N., SITHITHAWORN, P. (2018): Phylogenetic relationships within the *Opisthorchis viverrini* species complex with specific analysis of *O. viverrini sensu lato* from Sakon Nakhon,

Thailand by mitochondrial and nuclear DNA sequencing. *Infect Genet Evol*, 62: 86-94. DOI:10.1016/j.meegid.2018.04.022

PRAKOBWONG, S., & SUWANNATRAI, K. (2020): Reduction of Reinfection Rates with *Opisthorchis viverrini* through a Three-Year Management Program in Endemic Areas of Northeastern Thailand. *Korean J Parasitol*, 58(5): 527-535. DOI:10.3347/kjp.2020.58.5.527

PRATAP, P., RAZA, S.T., PRATAP, S. (2018). Cholangiocarcinoma: Etiology, Pathogenesis, Diagnosis, and Management. In: Waly, M., Rahman, M. (eds) *Bioactive Components, Diet and Medical Treatment in Cancer Prevention*. Volume 1. Berlin, Germany: Springer, Cham, pp. 201-213. DOI:10.1007/978-3-319-75693-6_16

RACHPRAKHON, P., & PURIVIROJKUL, W. (2021): Very low prevalence of *Opisthorchis viverrini* s.l. cercariae in *Bithynia siamensis siamensis* snails from the canal network system in the Bangkok Metropolitan Region, Thailand. [Très faible prévalence des cercaires d'*Opisthorchis viverrini* s.l. chez les mollusques *Bithynia siamensis siamensis* du réseau de canaux dans la région métropolitaine de Bangkok, Thaïlande]. *Parasite*, 28: 2-2. DOI:10.1051/parasite/2020072

REDEKOP, W. K., LENK, E. J., LUYENDIJK, M., FITZPATRICK, C., NIESSEN, L., STOLK, W. A., . . . HONTELEZ, J. A. (2017): The socioeconomic benefit to individuals of achieving the 2020 targets for five preventive chemotherapy neglected tropical diseases. *PLOS Neglected Tropical Diseases*, 11(1): e0005289. DOI:10.1371/journal.pntd.0005289

RIZVI, S., & GORES, G. J. (2013): Pathogenesis, diagnosis, and management of cholangiocarcinoma. *Gastroenterology*, 145(6): 1215-1229. DOI:10.1053/j.gastro.2013.10.013

ROBINSON, M. W., & SOTILLO, J. (2022): Foodborne trematodes: old foes, new kids on the block and research perspectives for control and understanding host-parasite interactions. *Parasitology*, 149(10): 1257-1261. DOI:10.1017/S0031182022000877

RODRIGUES, P. M., VOGEL, A., ARRESE, M., BALDERRAMO, D. C., VALLE, J. W., & BANALES, J. M. (2021). Next-Generation Biomarkers for Cholangiocarcinoma. *Cancers (Basel)*, 13(13):3222. DOI:10.3390/cancers13133222

SAENNA, P., HURST, C., ECHAUBARD, P., WILCOX BRUCE, A., & SRIPA, B. (2017): Fish sharing as a risk factor for *Opisthorchis viverrini* infection: evidence from two villages in north-eastern Thailand. *Infect Dis Poverty*, 6(2): 48-56. DOI:10.1186/s40249-017-0281-7

SAIJUNTHA, W. (2007): A comprehensive investigation of genetic variation among natural populations of *Opisthorchis viverrini* in Thailand and Lao PDR. Ph. D Thesis, Khon Kaen: Khon Kaen University.

SAIJUNTHA, W., SITHITHAWORN, P., KATSOPIT, N., ANDREWS, R. H., & PETNEY, T. N. (2014). Liver Flukes: Clonorchis and Opisthorchis. In R. Toledo & B. Fried (Eds.) *Digenetic Trematodes*. New York, NY: Springer New York, pp. 153-199. DOI:10.1007/978-1-4939-0915-5_6

SAIJUNTHA, W., SITHITHAWORN, P., KATSOPIT, N., ANDREWS, R. H., & PETNEY, T. N. (2019). Liver Flukes: Clonorchis and Opisthorchis. *Adv Exp Med Biol*, 1154: 139-180. DOI:10.1007/978-3-030-18616-6_6

SALEM, P. E. S., GHAZALA, R. A., EL GENDI, A. M., EMARA, D. M., & AHMED, N. M. (2020). The association between circulating MicroRNA-150 level and cholangiocarcinoma. *J Clin Lab Anal*, 34(11): e23397. DOI:10.1002/jcla.23397

SALTYKOVA, I. V., PETROV, V. A., & BRINDLEY, P. J. (2018). Chapter One - Opisthorchiasis and the Microbiome. In B. Sripa & P. J. Brindley (Eds.) *Advances in Parasitology*. Volume 102. Massachusetts, USA: Academic Press, pp. 1-23. DOI:10.1016/bs.apar.2018.07.001

SAWANYAWISUTH, K., SASHIDA, G., & SHENG, G. (2021). Epithelial-Mesenchymal Transition in Liver Fluke-Induced Cholangiocarcinoma. *Cancers*, 13(4): 791. DOI:10.3390/cancers13040791

SHIN, D. W., MOON, S.-H., & KIM, J. H. (2023). Diagnosis of Cholangiocarcinoma. *Diagnostics*, 13(2): 233. DOI:10.3390/diagnostics13020233

SILSIRIVANIT, A., SAWANYAWISUTH, K., RIGGINS, G. J., & WONGKHAM, C. (2014). Cancer biomarker discovery for cholangiocarcinoma: the high-throughput approaches. *J Hepatobiliary Pancreat Sci*, 21(6): 388-396. DOI:10.1002/jhbp.68

SINGAL, A. G., LAMPERTICO, P., & NAHON, P. (2020). Epidemiology and surveillance for hepatocellular carcinoma: New trends. *J Hepatol*, 72(2): 250-261. DOI:10.1016/j.jhep.2019.08.025

SMOUT, M. J., SRIPA, B., LAHA, T., MULVENNA, J., GASSER, R. B., YOUNG, N. D., BETHONY, J.M., BRINDLEY, P.J., LOUKAS, A. (2011). Infection with the carcinogenic human liver fluke, *Opisthorchis viverrini*. *Mol Biosyst*, 7(5): 1367-1375. DOI:10.1039/c0mb00295j

SOTA, P., SUTTIPRAPA, S., TANGKAWATTANA, S., SRIPA, M., BLAIR, D., & SRIPA, B. (2022). Does *Opisthorchis viverrini* circulate between humans and domestic cats in an endemic area in Thailand? *Parasitology*, 149(10): 1334-1338. DOI:10.1017/S0031182022000646

SRIPA, B., & ECHAUBARD, P. (2017). Prospects and Challenges towards Sustainable Liver Fluke Control. *Trends Parasitol*, 33(10): 799-812. DOI:10.1016/j.pt.2017.06.002

SRIPA, B., JUMNAINSONG, A., TANGKAWATTANA, S., & HASWELL, M. R. (2018). Immune response to *Opisthorchis viverrini* infection and its role in pathology. *Adv Parasitol*, 102: 73-95. DOI:10.1016/bs.apar.2018.08.003

SRIPA, B., SUWANNATRAI, A. T., SAYASONE, S., DO, D. T., KHIEU, V., & YANG, Y. (2021). Current status of human liver fluke infections in the Greater Mekong Subregion. *Acta Tropica*, 224: 106133. DOI:10.1016/j.actatropica.2021.106133

SRIPA, B., TANGKAWATTANA, S., & BRINDLEY, P. J. (2018). Chapter Five - Update on Pathogenesis of Opisthorchiasis and Cholangiocarcinoma. In B. Sripa & P. J. Brindley (Eds) *Advances in Parasitology*. Volume 102. Massachusetts, USA: Academic Press, pp. 97-113. DOI:10.1016/bs.apar.2018.10.001

SRIPA, B., TANGKAWATTANA, S., & SANGNIKUL, T. (2017). The Lawa model: A sustainable, integrated opisthorchiasis control program using the EcoHealth approach in the Lawa Lake region of Thailand. *Parasitol Int*, 66(4): 346-354. DOI:10.1016/j.parint.2016.11.013

STEELE, J. A., RICHTER, C. H., ECHAUBARD, P., SAENNA, P., STOUT, V., SITHITHAWORN, P., & WILCOX, B. A. (2018). Thinking beyond Opisthorchis viverrini for risk of cholangiocarcinoma in the lower Mekong region: a systematic review and meta-analysis. *Infect Dis Poverty*, 7(1): 44. DOI:10.1186/s40249-018-0434-3

SUEBLINVONG, T., PIYAMPUTRA, A., PRIYANONT, S., & THAITHONG, S. (1993). Isoenzyme characterization of *Opisthorchis viverrini* from man. *Chula Med J*, 37(6): 375-385.

SUWANNAHITATORN, P., WEBSTER, J., RILEY, S., MUNGTIN, M., & DONNELLY, C. A. (2019). Uncooked fish consumption among those at risk of *Opisthorchis viverrini* infection in central Thailand. *PLoS One*, 14(1): e0211540. DOI:10.1371/journal.pone.0211540

SUWANNATRAI, A., SAICHUA, P., & HASWELL, M. (2018). Chapter Two - Epidemiology of *Opisthorchis viverrini* Infection. In B. Sripa & P. J. Brindley (Eds.) *Advances in Parasitology*, Volume 101. Massachusetts, USA: Academic Press, pp. 41-67. DOI:10.1016/bs.apar.2018.05.002

TANGKAWATTANA, S., SEREERAK, P., UPONTAIN, S., TANGKAWATTANA, P., & SRIPA, B. (2021). Investigation of possible alternate animal reservoir hosts of *Opisthorchis viverrini*. *Acta Tropica*, 217: 105850. DOI:10.1016/j.actatropica.2021.105850

TANGKAWATTANA, S., & SRIPA, B. (2018). Chapter Six - Integrative EcoHealth/One Health Approach for Sustainable Liver Fluke Control: The Lawa Model. In B. Sripa & P. J. Brindley (Eds.) *Advances in Parasitology*, Volume 101. Massachusetts, USA: Academic Press, pp. 115-139. DOI:10.1016/bs.apar.2018.07.002

TANTRAWATPAN, C., SAIJUNTHA, W., BUNCHOM, N., SUKSAVATE, W., PILAP, W., WALALITE, T., AGATSUMA, T., TAWONG, W., SITHITHAWORN, P., ANDREWS, R.H., PETNEY, T.N. (2020). Genetic structure and geographical variation of *Bithynia siamensis goniomphalos sensu lato* (Gastropoda: Bithyniidae), the snail intermediate host of *Opisthorchis viverrini sensu lato* (Digenea: Opisthorchiidae) in the Lower Mekong Basin revealed by mitochondrial DNA sequences. *Int J Parasitol*, 50(1): 55-62. DOI:10.1016/j.ijpara.2019.10.007

THAENKHAM, U., NUAMTANONG, S., SA-NGUANKIAT, S., YOONUAN, T., TOUCH, S., MANIVONG, K., VONGHACHACK, Y., SATO, M., WAIKAGUL, J. (2010). Monophyly of *Opisthorchis viverrini* populations in the lower Mekong Basin, using mitochondrial DNA nad1 gene as the marker. *Parasitol Int*, 59(2): 242-247. DOI:10.1016/j.parint.2010.02.009

THAEWNONGIEW, K., SINGTHONG, S., KUTCHAMART, S., TANGSAWAD, S., PROMTHET, S., SAILUGKUM, S., & WONGBA, N. (2014). Prevalence and risk factors for *Opisthorchis viverrini* infections in upper Northeast Thailand. *Asian Pac J Cancer Prev*, 15(16): 6609-6612. DOI:10.7314/apjcp.2014.15.16.6609

THINKHAMROP, K., KHUNTIKEO, N., SITHITHAWORN, P., THINKHAMROP, W., WANGDI, K., KELLY, M.J., SUWANNATRAI, A.T., GRAY, D.J. (2019). Repeated praziquantel treatment and *Opisthorchis viverrini* infection: a population-based cross-sectional study in northeast Thailand. *Infect Dis Poverty*, 08(02): 34-42. DOI:10.1186/s40249-019-0529-5

TILLI, M., OLLIARO, P., GOBBI, F., BISOFFI, Z., BARTOLONI, A., & ZAMMARCHI, L. (2021). Neglected tropical diseases in non-endemic countries in the era of COVID-19 pandemic: the great forgotten. *J Travel Med*, 28(1): taaa179. DOI:10.1093/jtm/taaa179

TSHERING, G., DORJI, P. W., CHAIJAROENKUL, W., & NA-BANGCHANG, K. (2018). Biomarkers for the Diagnosis of Cholangiocarcinoma: A Systematic Review. *Am J Trop Med Hyg*, 98(6): 1788-1797. DOI:10.4269/ajtmh.17-0879

VALE, N., GOUVEIA, M. J., BOTELHO, M., SRIPA, B., SUTTIPRAPA, S., RINALDI, G., GOMES, P., BRINDLEY, P. J., DA COSTA, J. M. C. (2013). Carcinogenic liver fluke *Opisthorchis viverrini* oxysterols detected by LC-MS/MS survey of soluble fraction parasite extract. *Parasitol Int*, 62(6): 535-542. DOI:10.1016/j.parint.2013.08.001

745 WU, Y., HAYAT, K., HU, Y., & YANG, J. (2022). Long Non-Coding RNAs as Molecular Biomarkers in
746 Cholangiocarcinoma. *Front Cell Dev Biol*, 10: 890605-890605. DOI:10.3389/fcell.2022.890605
747 YANG, Y., DENG, X., LI, Q., WANG, F., MIAO, L., & JIANG, Q. (2020). Emerging roles of long noncoding RNAs
748 in cholangiocarcinoma: Advances and challenges. *Cancer Commun (Lond)*, 40(12): 655-680.
749 DOI:10.1002/cac2.12109
750 ZENG, X., & TAO, H. (2015). Diagnostic and prognostic serum marker of cholangiocarcinoma (Review). *Oncol Lett*,
751 9(1): 3-8. DOI:10.3892/ol.2014.2696
752 ZIMMERMANN, A. (2019): Tumors and tumor-like lesions of the hepatobiliary tract. 1st Edition, Switzerland,
753 Springer International Publishing, 3559pp.
754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770

771

772

773

774

775

<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/> Taeniasis/cysticercosis	<input type="checkbox"/> Leishmaniasis
<input type="checkbox"/> Food-borne trematode	<input type="checkbox"/> Buruli ulcer
<input type="checkbox"/> Lymphatic filariasis	<input type="checkbox"/> Frambusia
<input type="checkbox"/> Schistosomiasis	<input type="checkbox"/> Lepra
<input type="checkbox"/> Helminthiasis	<input type="checkbox"/> Trachoma
<input type="checkbox"/> Onchocerciasis	<input type="checkbox"/> Chromoblastomycosis, mycetoma and other deep fungal infections
<input type="checkbox"/> Dracunculiasis	<input type="checkbox"/> Arbovirus infection
<input type="checkbox"/> Echinococcosis	<input type="checkbox"/> Rabies
<input type="checkbox"/> Chagas disease	<input type="checkbox"/> Scabies and other ectoparasites
<input type="checkbox"/> African Trypanosomiasis	<input type="checkbox"/> Venomous snake bite

Figure 1. List of 20 NTDs according to WHO

776

777

778

779

780

781

782

783

784





Figure 2. Life cycle of *Opisthorchis viverrini*

798



799

Figure 3. The eggs of *O. viverrine* (A) and adult worms (B)
(Taken from cdc.gov/dpdx/opisthorchiasis/index.html)

809

810

811

812

813

814

815

816

817

818




Figure 4. Classification based on anatomical location of cholangiocarcinoma

ORIGINALITY REPORT

6%

SIMILARITY INDEX

5%

INTERNET SOURCES

4%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

1

coek.info

Internet Source

1%

2

link.springer.com

Internet Source

1%

3

Weerachai Saijuntha, Ross H. Andrews,
Paiboon Sithithaworn, Trevor N. Petney.

"Current assessment of the systematics and
population genetics of *Opisthorchis viverrini*
sensu lato (Trematoda: Opisthorchiidae) and
its first intermediate host *Bithynia siamensis*
sensu lato (Gastropoda: Bithyniidae) in
Thailand and Southeast Asia", *Infection,
Genetics and Evolution*, 2021

Publication

1%

4

pubmed.ncbi.nlm.nih.gov

Internet Source

1%

5

www.ajtmh.org

Internet Source

1%

6

ugspace.ug.edu.gh

Internet Source

1%

7

"Digenetic Trematodes", Springer Science and
Business Media LLC, 2019

Publication

1 %

8

www.science.gov

Internet Source

1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On